

OXYTETRACYCLINE  
 TYPE B  
 BLUE BIRD  
 CATTLE FEED B  
 MEDICATED

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

**ACTIVE DRUG INGREDIENT**

Oxytetracycline.....20 g/lb

INDICATIONS FOR USE	OXYTETRACYCLINE Concentration in Type C Medicated Feed	lb of Type B/ton of Feed
For calves and beef and non-lactating dairy cattle for treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline.	1000 - 5000 g/ton*	50 - 250

\* The Type C medicated feed should be adequate to provide 10 mg oxytetracycline per pound of body weight daily.

**GUARANTEED ANALYSIS**

Crude Protein (Min) ..... %  
 Equivalent crude protein from Non-Protein Nitrogen (NPN) when added (max) ..... %  
 Crude Fat (Min) ..... %  
 Crude Fiber (Max) ..... %  
 Calcium, if added (Min) ..... %  
 Calcium, if added (Max) ..... %  
 Phosphorus, if added (Min) ..... %  
 Salt, if added (Min) ..... %  
 Salt, if added (Max) ..... %  
 Sodium (Min)\* ..... %  
 Sodium (Max)\* ..... %  
 Potassium, if added (Min) ..... ppm  
 Vitamin A, if added (Min) ..... IU/lb

\* Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Ingredients as defined by AAFCO

**WARNING: WHEN USED IN MILK REPLACERS, THE TREATMENT CLAIM (10 MG/LB) IS LIMITED TO BACTERIAL ENTERITIS CAUSED BY *ESCHERICHIA COLI* ONLY.**

\_\_\_ lbs (\_\_\_ kg) NET WEIGHT

BLUE BIRD FEED MILL  
 Robin, IN 00000