

BLUE BIRD CTC*
Type C Cattle Feed B/Bacterial enteritis and Bacterial pneumonia
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Calves, Beef and Non-Lactating Dairy Cattle: For treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to Chlortetracycline.

ACTIVE DRUG INGREDIENT

Chlortetracycline..... _____ g/ton **

GUARANTEED ANALYSIS

Crude Protein, Min..... _____ %
 NPN, Max _____ %
 Crude Fat, Min..... _____ %
 Crude Fiber, Min..... _____ %
 Calcium, Min..... _____ %
 Calcium, Max..... _____ %
 Phosphorus, Min..... _____ %
 Potassium, Min _____ %
 Salt, Min (if added) _____ %
 Salt, Max (if added) _____ %
 Sodium, Min (if exceeding salt guarantee)..... _____ %
 Sodium, Max (if exceeding salt guarantee)..... _____ %
 Vitamin A, Min..... _____ IU/lb

INGREDIENTS

Ingredients as defined by AAFCO

FEEDING DIRECTIONS

Treat for not more than 5 days. Feed approximately 400 g/t, varying with body weight and feed consumption to provide 10 mg/lb body weight per day. The following table gives examples of feeding rates and drug levels:

Drug Level in Feed (g/ton)	Drug Level in Feed (mg/lb)	Type C Medicated Feed Consumption (lb of feed per 100 lb bodyweight per day)
500	250	4
667	333	3
1,000	500	2
2,000	1,000	1

Withdrawal Periods and Residue Warnings: Withdraw 10 days prior to slaughter. A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

BLUE BIRD FEED COMPANY
CITY. STATE. ZIP

NET WT ON BAG OR BULK

* This Bluebird label refers to Chloratet™ brand of Chlortetracycline.

**The final printed feed label must show the actual concentration of chlortetracycline. This drug level is equivalent to approximately 400 g/ton, depending on body weight and feed consumption, but cannot be greater than 2000 g/ton.

Lot No.