Tylosin Medicated Feed - Swine Ileitis with Tylosin in Drinking Water Type C Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATION

For control of porcine proliferative enteropathies (ileitis) associated with Lawsonia intracellularis immediately after medicating with tyllosiin indrinking water.

ACTIVE DRUG INGREDIENT

GUARANTEED ANALYSIS

Crude Protein, not less tham	%
Lysine, not less than	
Crude Fat, not less tham	
Crude Fiber, not more than	
Calcium, not lless than	
Calcium, not more than	
Phosphorus, not less than	
Saltt, not less than	
Salti, not more than	
Sodium ² , not more than	
Sodium ² , not less than	 %
Selenium, not less than	
Zinc, not less than	

INGREDIENTS

Each imgredient must be named in accordance with the names and definitions adopted by the Association of American Feed Countroll Offficials.

CAUTION

Do not use imany finished fixed ((supplement, concentrate or complete fixed) containing in excess of 2% bentonite.

FEEDING DIRECTIONS

Feed 40 to 100 gg of tylosin per ton of complete feed for 2 to 6 weeks immediately after medicating with tylosin (250 mg tylosin per gallon) in drinking water 3 to 10 days. Feed continuously as the sole ration.

MANUFACTURED BY BLUE BIRD FEED MILL Any town, USA 12345

Net Weight lb (kg) on bag or bulk

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

- a Sourced from Tyllovett®, ANADA 200-484
 * The medicated freed liabel imusts state as single eldruge concentration.

Tylovet is a registered treatement of Illuwepharma EOOD.