

BLUE BIRD CTC*
Type C Turkey Feed/Bluecomb
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline.

ACTIVE DRUG INGREDIENT

Chlortetracycline..... g/ton**

GUARANTEED ANALYSIS

Crude Protein, Min %
 Lysine, Min %
 Methionine, Min %
 Crude Fat, Min %
 Crude Fiber, Min %
 Calcium, Min %
 Calcium, Max %
 Phosphorus, Min %
 Salt, Min (if added) %
 Salt, Max (if added) %
 Sodium, Min (if exceeding salt guarantee)..... %
 Sodium, Max (if exceeding salt guarantee)..... %

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Feed continuously for 7 to 14 days to provide 25 mg chlortetracycline per pound of body weight per day. The following table gives examples of feeding rates and drug levels:

Drug level in Feed (g/ton)	Drug level in Feed (mg/lb)	Bodyweight	Type C Medicated Feed Consumption (lbs. per bodyweight per day)
1,000	500	5	0.25
1,250	625	15	0.6
1,667	833	20	0.6
1000	500	30	1.5

Withdrawal Periods and Residues Warnings: No withdrawal period is required when used according to label. Do not feed to turkeys producing eggs for human consumption.

BLUE BIRD FEED COMPANY
CITY. STATE. ZIP
NET WEIGHT ON BAG OR BULK

* This Bluebird label refers to Chloratet™ brand of Chlortetracycline.

**Final label for Type B medicated feed must state a single concentration of drug Lot No.