

BLUE BIRD CTC*
Type B Turkey Feed/Mycoplasma synoviae
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to chlortetracycline in turkeys.

ACTIVE DRUG INGREDIENT

Chlortetracycline..... 201-80,000 g/ton**

GUARANTEED ANALYSIS

Crude Protein, Min %
 Lysine, Min %
 Methionine, Min %
 Crude Fat, Min %
 Crude Fiber, Min %
 Calcium, Min %
 Calcium, Max %
 Phosphorus, Min %
 Salt, Min (if added) %
 Salt, Max (if added) %
 Sodium, Min (if exceeding salt guarantee)..... %
 Sodium, Max (if exceeding salt guarantee)..... %

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed containing 200 grams chlortetracycline per ton.

The following table provides examples of mixing rates:

Type B Medicated Feed Concentration (g/ton)	Type B Medicated Feed to make one ton of Type C Medicated Feed (lb)	Drug Level in Type C Medicated Feed (g/ton)
8,000	50	200
16,000	25	200
20,000	20	200

Feed the resulting Type C medicated feed continuously for 7 to 14 days.

Withdrawal Periods and Residues Warnings: No withdrawal period is required when used according to label. Do not feed to turkeys producing eggs for human consumption.

BLUE BIRD FEED COMPANY
CITY. STATE. ZIP
NET WEIGHT ON BAG OR BULK

* This Bluebird label refers to Chloratet™ brand of Chlortetracycline.

**Final label for Type B medicated feed must state a single concentration of drug Lot No.