

OXYTETRACYCLINE
TYPE C
BLUE BIRD
CHICKEN FEED A1
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For chickens for control of infectious synovitis caused by *Mycoplasma synoviae*; control of fowl cholera caused by *Pasteurella multocida* susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENT

Oxytetracycline 100 to 200 g/ton

GUARANTEED ANALYSIS

Crude Protein (Min) %
Lysine (Min) %
Methionine (Min) %
Crude Fat (Min) %
Crude Fiber (Max) %
Calcium (Min) %
Calcium (Max) %
Phosphorus (Min) %
Salt, if added (Min) %
Salt, if added (Max) %
Total Sodium (Min)* %
Total Sodium (Max)* %

* Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed 100 - 200 g oxytetracycline per ton of feed continuously for 7 - 14 days.

WARNING: ZERO-DAY WITHDRAWAL. DO NOT USE IN LOW-CALCIUM FEED CONTAINING LESS THAN 0.55% DIETARY CALCIUM. USE IN SUCH FEEDS MAY RESULT IN VIOLATIVE RESIDUES. DO NOT ADMINISTER TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION.

_____ lbs (_____ kg) NET WEIGHT

BLUE BIRD FEED MILL
Robin, IN 00000