

OXYTETRACYCLINE
TYPE C
BLUE BIRD
TURKEY FEED B
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For turkeys for control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENT

Oxytetracycline200 g/ton

GUARANTEED ANALYSIS

Crude Protein (Min).....%
Lysine (Min)%
Methionine (Min).....%
Crude Fat (Min)%
Crude Fiber (Max)%
Calcium (Min).....%
Calcium (Max)%
Phosphorus (Min).....%
Salt, if added (Min)%
Salt, if added (Max)%
Total Sodium (Min)*%
Total Sodium (Max)*%

*Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed 200 g oxytetracycline per ton of feed continuously for 7-14 days.

WARNING: WITHDRAW 5 DAYS BEFORE SLAUGHTER. DO NOT ADMINISTER TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION.

____lbs (____kg) NET WEIGHT

BLUE BIRD FEED MILL
Robin, IN 00000