

OXYTETRACYCLINE
 TYPE B
 BLUE BIRD
 TURKEY FEED C
 MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

ACTIVE DRUG INGREDIENT

Oxytetracycline.....20 g/lb

INDICATIONS FOR USE	OXYTETRACYCLINE Concentration in Type C Medicated Feed	lb of Type B/ton of Feed
For turkeys for control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline.	400 – 1000 g/ton*	20 - 50

* The Type C medicated feed should be adequate to provide 25 mg oxytetracycline per pound of body weight daily.

GUARANTEED ANALYSIS

Crude Protein (Min) %
 Lysine (Min)..... %
 Methionine (Min) %
 Crude Fat (Min)..... %
 Crude Fiber (Max)..... %
 Calcium (Min) %
 Calcium (Max) %
 Phosphorus (Min) %
 Salt, if added (Min) %
 Salt, if added (Max)..... %
 Total Sodium (Min)* %
 Total Sodium (Max)* %

* Minimum and maximum percentage of total sodium shall be guaranteed only when sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

WARNING: WITHDRAW 5 DAYS BEFORE SLAUGHTER. DO NOT ADMINISTER TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION.

___ lbs (___ kg) NET WEIGHT

BLUE BIRD FEED MILL
 Robin, IN 00000