

**Erythromycin Chicken Ration III
Type C Medicated Feed
(Erythromycin Type C Medicated Feed)**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATION

As an aid in the prevention of chronic respiratory disease during periods of stress.

ACTIVE DRUG INGREDIENT

Erythromycin 92.5 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than	%
Crude Fat, not less than	%
Crude Fiber, not more than	%
Lysine (min).....	%
Methionine (min).....	%
Calcium (min)	%
Calcium (max)	%
Phosphorus (min).....	%
Salt ¹ (min).....	%
Salt ¹ (max).....	%
Sodium ² (min).....	%
Sodium ² (max).....	%

¹ If added

² Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO

FEEDING DIRECTIONS

Feed 92.5 g erythromycin per ton of feed for 2 days before stress and 3 to 6 days after stress.

WITHDRAWAL PERIODS

Withdrawal 24 hours before slaughter.

Medicated Feed Expiration Date: _____ (not to exceed 2 weeks after the date of manufacture)

Net Weight _____ lb. (_____ kg.)

Manufactured by
Blue Bird Feed Company
Blue bird, MD 00000