

CHLORTETRACYCLINE
TYPE C
BLUE BIRD
TURKEY FEED D
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For turkeys for control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline.

ACTIVE DRUG INGREDIENT

Chlortetracycline calcium complex400 to 1000 g/ton

GUARANTEED ANALYSIS

Crude Protein (Min)	_____%
Lysine (Min).....	_____%
Methionine (Min)	_____%
Crude Fat (Min).....	_____%
Crude Fiber (Max).....	_____%
Calcium (Min)	_____%
Calcium (Max)	_____%
Phosphorus (Min).....	_____%
Salt, if added (Min)	_____%
Salt, if added (Max).....	_____%
Total Sodium (Min)*	_____%
Total Sodium (Max)*	_____%

*Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed 25 mg of chlortetracycline per pound of body weight daily continuously for 7-14 days. Consumption rates may vary and the chlortetracycline concentration in the feed should be adjusted based on consumption rates. For example, if a bird weighs 15 lb, consuming 0.75 lb of complete feed per day, the feed should contain 1000 g chlortetracycline per ton.

RESIDUE WARNINGS AND WITHDRAWAL PERIODS: ZERO-DAY WITHDRAWAL PERIOD. DO NOT ADMINISTER TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION.

_____lbs (____kg) NET WEIGHT

BLUE BIRD FEED MILL
Robin, IN 00000