

CHLORTETRACYCLINE
TYPE C
BLUE BIRD
TURKEY FEED A
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For turkeys for control of infectious synovitis caused by *M. synoviae* susceptible to chlortetracycline.

ACTIVE DRUG INGREDIENT

Chlortetracycline calcium complex 200 g/ton

GUARANTEED ANALYSIS

Crude Protein (Min)	_____ %
Lysine (Min).....	_____ %
Methionine (Min)	_____ %
Crude Fat (Min).....	_____ %
Crude Fiber (Max).....	_____ %
Calcium (Min)	_____ %
Calcium (Max)	_____ %
Phosphorus (Min).....	_____ %
Salt, if added (Min)	_____ %
Salt, if added (Max).....	_____ %
Total Sodium (Min)*.....	_____ %
Total Sodium (Max)*	_____ %

*Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed 200 g chlortetracycline per ton of feed continuously for 7-14 days.

RESIDUE WARNINGS AND WITHDRAWAL PERIODS: ZERO-DAY WITHDRAWAL PERIOD. DO NOT ADMINISTER TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION.

_____ lbs (_____ kg) NET WEIGHT

BLUE BIRD FEED MILL
Robin, IN 00000