## NEOMYCIN - OXYTETRACYCLINE TYPE C BLUE BIRD SWINE FEED A MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

## ACTIVE DRUG INGREDIENT<sup>1</sup>

Oxytetracycline (from oxytetracycline dihydrate) equivalen	t to oxytetracycline hydrochloride400 to 600 g/ton
Neomycin Sulfate	400 to 600 g/ton
<sup>1</sup> In a one to one ratio.	

INDICATIONS FOR USE	OXYTETRACYCLINE and NEOMYCIN SULFATE AMOUNT
Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> susceptible to oxytetracycline and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by <i>E. coli</i> susceptible to neomycin.	10 mg/lb of body weight daily* Feed continuously for 7-14 days
For breeding swine for control and treatment of Leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to oxytetracycline	10 mg/lb of body weight daily* Feed continuously for not more than 14 days

<sup>\*</sup>Consumption rates may vary and the oxytetracycline and neomycin sulfate concentrations in the feed should be adjusted based on consumption rates. For example, at the consumption rate of 4% of an animal's body weight, feed has to contain 500 g/ton oxytetracycline and 500 g/ton neomycin sulfate.

## **GUARANTEED ANALYSIS**

Crude Protein (Min)	%
Lysine (Min)	
Crude Fat (Min)	
Crude Fiber (Max)	
Calcium (Min)	
Calcium (Max)	
Phosphorus (Min)	
Salt (Min)	
Salt (Max)	
Selenium (Min)	
Zinc (Min)	ppm

(Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee)

## **INGREDIENTS**

Ingredients as defined by AAFCO

WARNING: WITHDRAW 5 DAYS BEFORE SLAUGHTER.

\_\_\_\_lbs (\_\_\_kg) NET WEIGHT

BLUE BIRD FEED MILL Robin, IN 00000