

Melengestrol Acetate + Lasalocid - HEIFER DRY Feed TD

Type C Medicated Feed

(melengestrol acetate and lasalocid Type C top-dress medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS

For increased rate of weight gain, improved feed efficiency, and suppression of estrus (heat) in heifers being fed in confinement for slaughter.

ACTIVE DRUG INGREDIENTS

Melengestrol acetate¹ (0.0000276 – 0.00022%) or 0.25 to 2.0 g/ton*
Lasalocid (as lasalocid sodium) 100 to 1440 g/ton*
Each pound contains (0.125-1.0) mg melengestrol acetate, (50-720) mg lasalocid.

GUARANTEED ANALYSIS

Crude Protein, not less than	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than	_____	%
Salt ² , not less than	_____	%
Salt ² , not more than.....	_____	%
Sodium ³ , not less than.....	_____	%
Sodium ³ , not more than.....	_____	%
Potassium, not less than	_____	%

²If added.

³ Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials. This includes trace minerals and vitamins.

DIRECTIONS FOR USE

Feed to heifers at the rate of 0.5 to 2.0 pound(s) per head per day (specify one level) to provide 0.25-0.5 mg melengestrol acetate per head per day (specify one level), 100 to 360 mg lasalocid per head per day (specify one level). Mix into a complete feed prior to feeding. When mixing this supplement into a complete ration, average daily feed intake (air dry or ~90% DM) must be estimated prior to mixing and feeding so that the specified per head per day level of melengestrol acetate is consumed.

Melengestrol acetate and lasalocid may be fed continuously throughout the time the heifers are being grown and finished for slaughter.

ADDITIONAL INFORMATION

May be fed to heifers in combination with a complete feed containing 8 to 10 grams of tylosin per ton of

air dried feed (~90% DM) for reduction of incidence of liver abscesses associated with *Fusobacterium necrophorum* and *Arcanobacterium pyogenes*. When tylosin is fed in combination with melengestrol acetate and lasalocid, the level of tylosin is limited to 90 mg/head/day.

IMPORTANT: Melengestrol acetate is only for use in heifers being fed in confinement for slaughter. Not effective in steers and spayed heifers. This combination may only be marketed as a pelleted product.

CAUTION

Safety of lasalocid for use in unapproved species has not been established. Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle and sheep. Do not allow horses or other equines access to Type B feeds containing lasalocid as ingestion may be fatal.

WARNINGS

Warning: No withdrawal period is required. A withdrawal time has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

When mixing and handling lasalocid, use protective clothing, impervious gloves and a dust mask, avoid contact with eyes. Operators should wash thoroughly with soap and water after handling. Use melengestrol acetate only as directed.

Lot Number (if applicable): _____

Expiration date: _____

(Must be used within 90 days of manufacture)

MANUFACTURED BY

Blue Bird Feed Company
Anytown, USA 11111

Net weight lb (kg) on bag or bulk

*The medicated feed label must state a single drug concentration.

¹ Sourced from HeifermaxTM, ANADA# 200-343

Elanco, Heifermax and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

Last Revision: November 2016