

Melengestrol Acetate + Lasalocid – HEIFER DRY TD
Type C Medicated Feed
(melengestrol acetate and lasalocid Type C top-dress medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For Increased Rate of Weight Gain, Improved Feed Efficiency, and Suppression of Estrus (Heat) in heifers fed in confinement for slaughter.

ACTIVE INGREDIENTS¹

Melengestrol acetate^a.....(0.0000276-0.00022%) or 0.25 to 2.0 g/ton
Lasalocid (as lasalocid sodium)^b.....(100-1440) g/ton
Each pound contains (0.125-1.0) mg melengestrol acetate, (50-720) mg lasalocid.

GUARANTEED ANALYSIS

Crude protein, not less than.....	%
Crude fat, not less than.....	%
Crude fiber, not more than.....	%
Calcium, not less than.....	%
Calcium, not more than.....	%
Phosphorus, not less than.....	%
Potassium, not less than.....	%
Salt ² , not less than.....	%
Salt ² , not more than.....	%
Sodium ³ , not less than.....	%
Sodium ³ , not more than.....	%

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials. This includes trace minerals and vitamins.

DIRECTIONS FOR USE

Feed to heifers at the rate of 0.5 to 2.0 pound(s) per head per day (specify one level) to provide 0.25-0.5 mg melengestrol acetate per head per day (specify one level) and 100 to 360 mg lasalocid per head per day (specify one level). This Type C product may be top dressed onto or mixed into a complete feed prior to feeding.

Melengestrol acetate and lasalocid may be fed continuously throughout the time the heifers are being grown and finished for slaughter.

ADDITIONAL INFORMATION

May be fed to heifers in combination with a complete feed containing 8 to 10 grams of tylosin per ton for reduction of incidence of liver abscesses associated with *Fusobacterium necrophorum* and *Arcanobacterium pyogenes*. When tylosin is fed in combination with melengestrol acetate and lasalocid, the level of tylosin is limited to 90 mg/head/day.

IMPORTANT: Melengestrol acetate is only for use in heifers being fed in confinement for slaughter. Not effective in steers and spayed heifers. This combination may only be marketed as a pelleted product.

CAUTION

Do not allow horses or other equines access to premixes or supplements containing lasalocid, as ingestion may be fatal. The safety of lasalocid for use in unapproved species has not been established. Mixing errors resulting in excessive lasalocid concentration which could be fatal to cattle and sheep.

Warning: No withdrawal period is required. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

When mixing and handling lasalocid, use protective clothing, impervious gloves and a dust mask, avoid contact with eyes. Operators should wash thoroughly with soap and water after handling.

Use melengestrol acetate only as directed.

Lot Number (if applicable): _____

Expiration date: _____

(Must be used within 90 days of manufacture)

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

Final printed label on formulated Type C medicated feed must bear a single concentration for each drug.

®MGA® is the proprietary name of melengestrol acetate Type A medicated article (NADA 039-402).

®Bovatec® is the proprietary name of lasalocid sodium Type A medicated article (NADA 096-298).

Huvepharma Inc. December 2016