

**DCX + Chlortetracycline-AUREO Type C Cattle Supplement Medicated  
Type C Medicated Feed**  
(decoquinatate and chlortetracycline Type C top-dress medicated feed)

**Caution:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Calves, beef, and non-lactating dairy cattle. For the prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii*; for treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to chlortetracycline.

**ACTIVE DRUG INGREDIENTS\***

Decoquinatate <sup>a</sup> .....	90.9 to 535.7 g/ton*
Chlortetracycline <sup>b</sup> (as hydrochloride).....	4000 to 20,000 g/ton*

**GUARANTEED ANALYSIS**

Crude protein, not less than .....	_____ %
Crude fat, not less than .....	_____ %
Crude fiber, not more than.....	_____ %
Calcium, not less than.....	_____ %
Calcium, not more than .....	_____ %
Phosphorus, not less than.....	_____ %
Potassium, not less than.....	_____ %
Salt <sup>1</sup> , not less than.....	_____ %
Salt <sup>1</sup> , not more than .....	_____ %
Sodium <sup>2</sup> , not less than .....	_____ %
Sodium <sup>2</sup> , not more than .....	_____ %

<sup>1</sup>If added.

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials. This includes trace minerals and vitamins.

**FEEDING DIRECTIONS**

Administer as a top dress supplement or mix into the daily ration to provide 22.7 mg decoquinatate per 100 lb of body weight per day and 1 g chlortetracycline per 100 lb body weight/day for not more than 5 days. \_\_\_\_\_\*\* pounds of this supplement will medicate 100 lbs of body weight. If top-dressed, spread uniformly on top of ration and allow sufficient space so that all cattle can eat at the same time. When it is fully consumed, resume feeding 22.7 mg decoquinatate per 100 lb of body weight/day for a total of 28 days to prevent coccidiosis.

**Caution:** Bentonite should not be used in decoquinatate feeds.

**Warning:** No withdrawal period is required. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Do not feed to animals producing milk for food.

Lot Number (if applicable): \_\_\_\_\_

**MANUFACTURED BY**  
Blue Bird Feed Company  
Robin, Indiana 11111

**NET WEIGHT ON BAG OR BULK**

---

\*Final printed label on formulated Type C Medicated feed must bear a single concentration for each drug. The ratio between chlortetracycline and decoquinatate must be 44:1.

\*\*Actual concentration of chlortetracycline and decoquinatate will vary with body weight and feed consumption to provide 22.7 mg decoquinatate per 100 lb of body weight per day and 1 g chlortetracycline per 100 lb body weight/day for not more than 5 days. The following table provides examples of supplementation rates and amount of drug per ton of feed.

Decoquinatate Level in Supplement		Chlortetracycline Level in Supplement		Pounds of Supplement per 100 pounds of body weight
mg/lb	grams/ton	mg/lb	grams/ton	
45.45	90.9	2000	4000	0.5
<b>56.75</b>	<b>113.5</b>	<b>2500</b>	<b>5000</b>	<b>0.4</b>
113.5	227	5000	10,000	0.2
227	454	10,000	20,000	0.1

<sup>a</sup>Deccox® is the proprietary name of decoquinatate Type A medicated article (NADA 039-417).

<sup>b</sup>Aureomycin® is the proprietary name of chlortetracycline Type A medicated article (NADA 048-761).

Zoetis Inc. [9 November 2016]