

**Chlortetracycline-AUREO (10 mg/lb BW) + Laidlomycin (5-10 g/ton) CATTLE FEED  
TYPE C Medicated Complete Feed**  
(chlortetracycline and laidlomycin propionate potassium Type C medicated feed)

**Caution:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline, and for improved feed efficiency in cattle fed in confinement for slaughter.

**ACTIVE DRUG INGREDIENTS\***

Chlortetracycline<sup>a</sup> ..... 500 to 4000 g/ton  
Laidlomycin propionate potassium<sup>b</sup> ..... 5 to 10 g/ton

**GUARANTEED ANALYSIS**

Crude protein, not less than ..... %  
Crude fat, not less than ..... %  
Crude fiber, not more than ..... %  
Calcium, not less than ..... %  
Calcium, not more than ..... %  
Phosphorus, not less than ..... %  
Potassium, not less than ..... %  
Salt<sup>1</sup>, not less than ..... %  
Salt<sup>1</sup>, not more than ..... %  
Sodium<sup>2</sup>, not less than ..... %  
Sodium<sup>2</sup>, not more than ..... %  
Vitamin A<sup>1</sup> (not less than) ..... IU/lb.

<sup>1</sup>If added.

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials. This includes trace minerals and vitamins.

**FEEDING DIRECTIONS**

Provide this Type C medicated feed to confined cattle to treat bacterial enteritis and/or pneumonia while improving feed efficiency. Feed continuously at a rate of 10 mg chlortetracycline per lb. body weight per day and 30 to 150 mg laidlomycin propionate potassium per head per day for not more than 5 days. The following table provides examples of feeding rates and amount of drug per ton of feed.

Body Wt. Lb.	Feed Consumption		Chlortetracycline		Laidlomycin	
	% Body Wt.	lb./head/day	Dose, mg/lb./day	Feeding Rate, g/ton	Dose, mg/head/day	Feeding Rate g/ton
500	2	10	10	1000	30	6
	2	10	10	1000	50	10
	3	15	10	667	38	5
	3	15	10	667	75	10
1000	2	20	10	1000	50	5
	2	20	10	1000	100	10
	3	30	10	667	75	5
	3	30	10	667	150	10

**Caution**

Do not allow horses or other equines access to feeds containing laidlomycin propionate potassium. The safety of laidlomycin propionate potassium in unapproved species has not been established. Not for use in animals intended for breeding. If manufactured using liquid Type B laidlomycin propionate potassium feed, this dry Type C medicated feed expires 7 days after the date of manufacture.

**Warning:** No withdrawal period is required. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.



Lot Number (if applicable): \_\_\_\_\_

**MANUFACTURED BY**

Blue Bird Feed Company  
Robin, Indiana 11111

**NET WEIGHT ON BAG OR BULK**

\*Final printed label on formulated Type C Medicated feed must bear a single concentration for each drug.

<sup>a</sup>Aureomycin® is the proprietary name of chlortetracycline Type A medicated article (NADA 048-761).

<sup>b</sup>Cattlyst® is the proprietary name of laidlomycin propionate potassium Type A medicated article (NADA 141-025).

Zoetis Inc. [7 September 2016]