

Sulfadimethoxine and Ormetoprim

For catfish and salmonids

Type B Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the control of enteric septicemia of catfish caused by *Edwardsiella ictaluri* strains susceptible to the sulfadimethoxine and ormetoprim combination.

For the control of furunculosis in salmonids (trout and salmon) caused by *Aeromonas salmonicida* strains susceptible to the sulfadimethoxine and ormetoprim combination.

Active Drug Ingredients

Sulfadimethoxine75.8 g/pound
Ormetoprim15.2 g/pound

Guaranteed Analysis

Crude Protein, not less thanXX.X%
Crude Fat, not less than.....XX.X%
Crude Fiber, not more thanXX.X%
Phosphorus, not less thanXX.X%

Ingredients

Ingredients as defined by AAFCO

Mixing Directions

IMPORTANT: Must be thoroughly mixed in feeds before use.

Medication of feed before pelletizing or extruding. Thoroughly mix sulfadimethoxine-ormetoprim Type B medicated feed into the appropriate amount of non-medicated mash feed prior to pelletizing or extruding.

Medication of feed after pelletizing. Thoroughly mix sulfadimethoxine-ormetoprim Type B medicated feed into a liquid slurry of water. Coat the non-medicated pelleted fish feed with the slurry which should be constantly agitated to ensure uniform suspension of the sulfadimethoxine-ormetoprim Type B medicated feed during the addition. Generally, one gallon of liquid slurry is needed for each 200 lbs of pellets.

<i>Feed Intake of Fish as Percent (%) of Body Weight</i>	<i>Lbs of Sulfadimethoxine-Ormetoprim Type B Medicated Feed</i>	<i>Lbs of Non-Medicated Feed</i>
1%	50	1950
2%	25	1975
3%	16.6	1983.4
4%	12.5	1987.5
5%	10	1990
6%	8.3	1991.7

Administer the resulting Type C medicated feed daily for five consecutive days to provide approximately 50 mg of active ingredients per kg (23 mg/lb) of live body weight of fish per day.

CAUTION: If fish show no improvement within 2 to 3 days, or if signs of disease reappear after termination of treatment, reevaluate management practices, diagnosis of outbreak, and establish susceptibility of the bacterial isolate(s) to the drug.

WARNING: For catfish, withdraw 3 days before release as stocker fish or slaughter. For salmonids, withdraw 42 days before release as stocker fish or slaughter.

Manufactured by

Blue Bird Feed Mill

Any Town, USA 11111

Net weight on bag or bulk