

VIRGINIAMYCIN/SALINOMYCIN
TYPE C MEDICATED FEED

BLUE BIRD
BROILER FEED
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For prevention of necrotic enteritis caused by *Clostridium perfringens* susceptible to virginiamycin and for the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

ACTIVE DRUG INGREDIENT

Virginiamycin 20 g/ton
Salinomycin 40 - 60 g/ton*

GUARANTEED ANALYSIS

Crude Protein (Min) %
Crude Fat (Min) %
Crude Fiber (Max) %
Calcium (Min) %
Calcium (Max) %
Phosphorus (Min) %
Salt (Min, if added) %
Salt (Max, if added) %
Lysine (Min) %
Methionine (Min) %
Sodium (Min, if exceeds salt min) %
Sodium (Max, if exceeds salt max) %

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed continuously as the sole ration for broiler or fryer chickens.

CAUTION: Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age. Not approved for use with pellet binders. May be fatal if accidentally fed to adult turkeys or horses.

_____ lbs (_____ kg) NET WEIGHT

BLUEBIRD FEED COMPANY
ANYTOWN, USA 12345

Lot #: _____

* Final printed label should contain one concentration of salinomycin.