

Medicated Broiler Chicken Feed

(avilamycin and narasin Type C medicated feed)

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS FOR USE

For the prevention of mortality caused by necrotic enteritis associated with *Clostridium perfringens*; and for the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima* in broiler chickens.

ACTIVE INGREDIENTS

Avilamycin ¹	13.6 to 40.9 g/ton*
Narasin ²	54 to 90 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____ %
Lysine, not less than.....	_____ %
Methionine, not less than.....	_____ %
Crude Fat, not less than.....	_____ %
Crude Fiber, not more than.....	_____ %
Calcium, not less than.....	_____ %
Calcium, not more than.....	_____ %
Phosphorus, not less than.....	_____ %
Salt ³ , not less than.....	_____ %
Salt ³ , not more than.....	_____ %
Sodium ⁴ , not less than.....	_____ %
Sodium ⁴ , not more than.....	_____ %

³If added.

⁴Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed as the sole ration for 21 consecutive days. Feed to chickens that are at risk of developing, but not yet showing clinical signs of, necrotic enteritis associated with *Clostridium perfringens* in broiler chickens.

CAUTIONS

To assure responsible antimicrobial drug use in broiler chickens, treatment administration must begin on or before 10 days of age. The safety of avilamycin has not been established in chickens intended for breeding purposes. Avilamycin has not been demonstrated to be effective in broiler chickens showing clinical signs of necrotic enteritis prior to the start of medication.

The Veterinary Feed Directive (VFD) expiration date must not exceed 90 days from the VFD issuance date. VFDs for avilamycin shall not be refilled.

WARNINGS

Withdrawal Period and Residue Warnings



No withdrawal period is required when used according to labeling.



User Safety Warnings

Not for use in humans. Keep out of reach of children. To report adverse effects in users, to obtain more information, or to obtain a Safety Data Sheet(s), contact Elanco at 1-800-428-4441.

Animal Safety Warnings

Do not allow adult turkeys, horses or other equines access to narasin formulations. Ingestion of narasin by these species has been fatal.

MANUFACTURED BY
BLUE BIRD FEED
MILL Any town, USA
12345

Net Weight lb (kg) on bag or bulk

Lot # _____

NADA #141-467, Approved by FDA
Last Revision October 2017

*The medicated feed label must state a single drug concentration.

¹ Sourced from Integrity™, NADA# 141-439

² Sourced from Monteban™, NADA# 118-980

Elanco, Integrity, Monteban and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.