

DERA/BOV COCCI — CATTLE FEED
TYPE C Medicated Feed
(chlortetracycline and lasalocid Type C medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* and for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline in cattle weighing up to 800 pounds.

ACTIVE DRUG INGREDIENTS*

Chlortetracycline^a500 to 4000 g/ton**
Lasalocid^b (as lasalocid sodium).....30 to 181.8 g/ton**

GUARANTEED ANALYSIS

Crude protein, not less than%
(This includes not more than _____% equivalent crude protein from non-protein nitrogen)¹
Crude fat, not less than%
Crude fiber, not more than%
Calcium, not less than%
Calcium, not more than%
Phosphorus, not less than%
Potassium, not less than%
Salt¹, not less than%
Salt¹, not more than%
Sodium², not less than%
Sodium², not more than%
Vitamin A¹, not less than IU/lb

¹If added

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

FEEDING DIRECTIONS

Hand feed continuously for not more than 5 days to provide 10 mg chlortetracycline per lb. body weight per day and 1 mg lasalocid per 2.2 lb. body weight per day to cattle weighing up to 800 lb. with a maximum of 360 mg of lasalocid per head per day. The following table provides examples of feeding rates and amounts:

Body Wt. Lb.	Feed Consumption		Chlortetracycline		Lasalocid	
	% Body Wt.	lb/head/day	Amount mg/lb BW/day	Feeding Rate g/ton	Amount mg/head/day	Feeding Rate g/ton
200	0.5	1	10	4000	90.9	181.8
	1	2	10	2000	90.9	90.9
	2	4	10	1000	90.9	45.5
	3	6	10	667	90.9	30.3
800	0.5	4	10	4000	360	180
	1	8	10	2000	360	90
	2	16	10	1000	360	45
	3	24	10	667	360	30

Caution

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

Warning: No withdrawal period is required. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Lot Number (if applicable): _____

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

*Final printed label on formulated Type C medicated feed must bear a single concentration for each drug.

**The ratio between chlortetracycline and lasalocid must be 22:1.

^aDeracin® is the proprietary name of chlortetracycline Type A medicated article (ANADA 200-510).

^bBOVATEC is the proprietary name of lasalocid sodium Type A medicated article (NADA 096-298).

Pharmgate LLC [March 2017]