

Salinomycin and Lincomycin CHICKEN Feed
Type C Medicated Complete Feed
(salinomycin sodium and lincomycin Type C medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Indications for Use

For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, and for the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to lincomycin in broiler chickens.

Active Ingredients

Salinomycin sodium activity^a40-60 g/ton (0.0044-0.0066%)*
Lincomycin^b (as lincomycin hydrochloride agricultural grade) 2 g/ton

Guaranteed Analysis

Crude Protein, not less than		%
Lysine, not less than		%
Methionine, not less than		%
Crude Fat, not less than		%
Crude Fiber, not more than		%
Calcium, not less than		%
Calcium, not more than		%
Phosphorus, not less than		%
Salt ¹ , not less than		%
Salt ¹ , not more than		%
Sodium ² , not less than		%
Sodium ² , not more than		%

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Each ingredient must be specifically named in accordance with the names and definitions adopted by the AAFCO.

Feeding Directions

Feed as the sole ration to broiler chickens.

Warnings

Withdrawal Periods and Residue Warnings

No withdrawal period is required when used according to labeling.
Do not feed to laying hens producing eggs for human consumption.



User Safety Warnings

Not for use in humans. Keep out of reach of children.

To obtain a Safety Data Sheet(s), contact Zoetis, Inc. at 1-888-963-8471 or www.zoetis.com.

Animal Safety Warnings

Not for use in laying hens, breeding chickens, or turkeys. May be fatal if accidentally fed to adult turkeys or to horses. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

Other Warnings

Caution: Do not use in Type C medicated feeds containing pellet binders.

Questions/Comments

To report suspected adverse events or side effects, or for technical assistance contact Zoetis Inc. at 1-888-963-8471 or www.zoetis.com. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/AnimalVeterinary/SafetyHealth>.

Lot Number (if applicable): _____

Manufactured by
Blue Bird Feed Company
Robin, Indiana 11111

Net Weight on Bag or Bulk

*Final printed label on formulated Type C medicated feed must bear single concentration.

^aBio-Cox® is the proprietary name of salinomycin sodium Type A medicated article (NADA 128-686).

^bLincomix® is the proprietary name of lincomycin Type A medicated article (NADA 097-505).

NADA #141-484, Approved by FDA
Zoetis Inc. [29 January 2018]