

# Monensin and Bacitracin Methylene Disalicylate Medicated Broiler Chicken Feed Type C Medicated Feed

To aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*, and for increase in rate of weight gain and improved feed efficiency.

## ACTIVE DRUG INGREDIENT

Monensin, USP.....	90 to 110 g/ton*
Bacitracin Methylene Disalicylate.....	5 to 25 g/ton*

## GUARANTEED ANALYSIS

Crude Protein, not less than .....	_____ %
Lysine, not less than .....	_____ %
Methionine, not less than .....	_____ %
Crude Fat, not less than .....	_____ %
Crude Fiber, not more than .....	_____ %
Calcium, not less than .....	_____ %
Calcium, not more than.....	_____ %
Phosphorus, not less than .....	_____ %
Salt <sup>1</sup> , not less than .....	_____ %
Salt <sup>1</sup> , not more than .....	_____ %
Sodium <sup>2</sup> , not less than .....	_____ %
Sodium <sup>2</sup> , not more than .....	_____ %

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

## INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

## FEEDING DIRECTIONS

Feed continuously as the sole ration.

## CAUTION

Do not feed to laying chickens. The use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain. Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal.

\*Final printed label on formulated Type C medicated feed must bear a single drug concentration.

**MANUFACTURED BY**  
**BLUE BIRD FEED MILL**  
Any town, USA 12345  
**Net Weight lb (kg) on bag or bulk**