

**Monensin and Lincomycin CHICKEN Feed
Type C Medicated Complete Feed
(monensin and lincomycin Type C medicated feed)**

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Indication for Use

As an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*, and for the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to lincomycin in broiler chickens.

Active Ingredients

Monensin^a 90 to 110 g/ton*
Lincomycin^b (as lincomycin hydrochloride agricultural grade) 2 g/ton

Guaranteed Analysis

Crude Protein, not less than	_____ %
Lysine, not less than	_____ %
Methionine, not less than	_____ %
Crude Fat, not less than	_____ %
Crude Fiber, not more than	_____ %
Calcium, not less than	_____ %
Calcium, not more than	_____ %
Phosphorus, not less than	_____ %
Salt ¹ , not less than	_____ %
Salt ¹ , not more than	_____ %
Sodium ² , not less than	_____ %
Sodium ² , not more than	_____ %

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Each ingredient must be specifically named in accordance with the names and definitions adopted by the AAFCO.

Feeding Directions

Feed as the sole ration to broiler chickens.



Withdrawal Periods and Residue Warnings.

No withdrawal period is required when used according to labeling.
Do not feed to laying hens producing eggs for human consumption.
Do not feed to chickens over 16 weeks of age.



User Safety Warnings

Not for use in humans. Keep out of reach of children.

To obtain a Safety Data Sheet(s), contact Zoetis, Inc. at 1-888-963-8471 or www.zoetis.com.

Animal Safety Warnings

Not for use in laying hens, breeding chickens, or turkeys. Not for replacement chickens intended to become broiler breeding chickens. Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

Other Warnings

Caution: In the absence of coccidiosis in broiler chickens, the use of monensin with no withdrawal period may limit feed intake, resulting in reduced weight gain.

Questions/Comments

To report suspected adverse events or side effects, or for technical assistance contact Zoetis Inc. at 1-888-963-8471 or www.zoetis.com. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/AnimalVeterinary/SafetyHealth>.

Do not use this medicated feed after: (90 days from the date of manufacture).
Lot Number (if applicable): _____

Manufactured by
Blue Bird Feed Company
Robin, Indiana 11111

Net Weight on Bag or Bulk

* Final printed label on formulated Type C medicated feed must bear single concentration.

^aCoban™ is the proprietary name of monensin Type A medicated article (NADA 038-878).

^bLincomix® is the proprietary name of lincomycin Type A medicated article (NADA 097-505).

NADA #141-491, Approved by FDA.
Zoetis Inc. [5 February 2018]