

**BLUE BIRD
BOV - COCCI
Type C Medicated Confined Sheep Feed**

For prevention of coccidiosis caused by *Eimeria ovina*, *E. crandallis*, *E. ovinoidalis* (*E. ninakohlyakimovae*), *E. parva*, and *E. intricata* in sheep maintained in confinement.

Active Drug Ingredient

Lasalocid (as Lasalocid sodium).....20 - 30 g/ton*

Guaranteed Analysis

Crude Protein (Min).....	_____	%
NPN ^a (Max).....	_____	%
Crude Fat (Min).....	_____	%
Crude Fiber (Max).....	_____	%
Calcium ^a (Min).....	_____	%
Calcium ^a (Max).....	_____	%
Phosphorus ^a (Min).....	_____	%
Salt ^a (Min).....	_____	%
Salt ^a (Max).....	_____	%
Sodium ^b (Min).....	_____	%
Sodium ^b (Max).....	_____	%
Copper ^c (Min).....	_____	ppm
Copper ^c (Max).....	_____	ppm
Selenium ^a (Min).....	_____	ppm
Vitamin A ^a (Min).....	_____	IU/lb

^a Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

^b Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

^c Copper minimum and maximum guarantee for sheep required when added or level exceeds 20 ppm

Ingredients

Ingredients as defined by AAFCO.

Feeding Directions

Feed continuously to provide not less than 15 mg or more than 70 mg lasalocid sodium activity per head per day depending on body weight.

The following table provides examples of feeding rates and amount of drug in feed.

Drug Level in Complete Feed		Pounds of Type C Feed/Head/Day	mg Lasalocid/head/day
grams/ton	mg/lb		
20	10	1.5	15
20	10	3	30
25	12.5	3.6	45
25	12.5	4.8	60
30	15	4.66	70

**Manufactured by
Blue Bird Feed Mill
City, State, Zip**

NET WEIGHT ON BAG OR BULK

*The final printed feed label should list a drug concentration that is adequate to provide not less than 15 mg or more than 70 mg of lasalocid per head per day

096298_04012010S

BEST COPY AVAILABLE