

**Clopidol and Lincomycin CHICKEN Feed**  
**Type C Medicated Complete Feed**  
(clopidol and lincomycin Type C medicated feed)

**Caution:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

**Indication for Use**

As an aid in the prevention of cecal and intestinal coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, and for the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to lincomycin in broiler chickens.

**ACTIVE INGREDIENTS**

Clopidol<sup>a</sup>.....113.5 g/ton (0.0125%)  
Lincomycin<sup>b</sup> (as lincomycin hydrochloride agricultural grade) .....2 g/ton

**GUARANTEED ANALYSIS**

Crude Protein, not less than.....	_____	%
Lysine, not less than .....	_____	%
Methionine, not less than.....	_____	%
Crude Fat, not less than .....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than .....	_____	%
Phosphorus, not less than .....	_____	%
Salt <sup>1</sup> , not less than.....	_____	%
Salt <sup>1</sup> , not more than .....	_____	%
Sodium <sup>2</sup> , not less than .....	_____	%
Sodium <sup>2</sup> , not more than .....	_____	%

<sup>1</sup> If added.

<sup>2</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be specifically named in accordance with the names and definitions adopted by the AAFCO.

**Feeding Directions**

Feed as the sole ration to broiler chickens.



**Withdrawal Periods and Residue Warnings**

No withdrawal period is required when used according to labeling.  
Do not feed to chickens over 16 weeks of age.



**User Safety Warnings**

Not for use in humans. Keep out of reach of children.

To obtain a Safety Data Sheet(s), contact Zoetis, Inc. at 1-888-963-8471 or [www.zoetis.com](http://www.zoetis.com).

**Animal Safety Warnings**

Not for use in laying hens, breeding chickens, or turkeys. Do not allow rabbits, hamsters, guinea pigs, horses or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

**Questions/Comments**

To report suspected adverse events or side effects, or for technical assistance contact Zoetis Inc. at 1-888-963-8471 or [www.zoetis.com](http://www.zoetis.com). For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/AnimalVeterinary/SafetyHealth>.

Lot Number (if applicable): \_\_\_\_\_

**Manufactured By**  
Blue Bird Feed Company  
Robin, Indiana 11111

**Net Weight on Bag or Bulk**

<sup>a</sup>Coyden® is the proprietary name of clopidol Type A medicated article (NADA 034-393).

<sup>b</sup>Lincomix® is the proprietary name of lincomycin Type A medicated article (NADA 097-505).

Approved by FDA under NADA #141-485.  
Zoetis Inc. [6 September 2018]