

**BLUE BIRD  
LPK  
Type B Liquid Cattle Feed – FE  
Medicated**

For improved feed efficiency in cattle fed in confinement for slaughter.

**Active Drug Ingredient:** Laidlomycin propionate potassium..... 100 – 2,000 g/ton\*

**Guaranteed Analysis**

Crude Protein (Min).....	_____	%
NPN <sup>a</sup> (Max).....	_____	%
Crude Fat (Min).....	_____	%
Crude Fiber (Max).....	_____	%
Calcium <sup>a</sup> (Min).....	_____	%
Calcium <sup>a</sup> (Max).....	_____	%
Phosphorus <sup>a</sup> (Min).....	_____	%
Potassium <sup>a</sup> (Min).....	_____	%
Salt <sup>a</sup> (Min).....	_____	%
Salt <sup>a</sup> (Max).....	_____	%
Sodium <sup>b</sup> (Min).....	_____	%
Sodium <sup>b</sup> (Max).....	_____	%
Vitamin A <sup>a</sup> (Min).....	_____	IU/lb

<sup>a</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>b</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

pH must be within the range of 6.0 - 8.0 and dry matter must be within the range of 62% - 75%.

**Ingredients**

Ingredients as defined by AAFCO

**Mixing Directions**

Thoroughly mix this liquid Type B medicated feed with non-medicated feed ingredients to manufacture a Type C medicated feed containing 5-10 grams laidlomycin propionate potassium per ton on a 90% dry matter basis. The resulting Type C medicated feed should be fed continuously to provide not less than 30 mg or more than 150 mg laidlomycin propionate potassium per head per day. The Type C feed made with this Type B liquid feed must be fed within 7 days of manufacture. The following table provides examples of mixing rates:

Type B laidlomycin concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C laidlomycin concentration (g/ton)
500	33.33	1966.67	8.33
1000	20	1980	10
2000	10	1990	10

*For liquid Type B feeds stored in recirculating tank systems:* Recirculate daily (even when not used) and immediately prior to use for not less than 10 minutes, moving not less than one (1) percent of the tank contents per minute from the bottom of the tank to the top. *For liquid Type B feeds stored in mechanical, air or other agitation-type tank systems:* Agitate daily (even when not used) and immediately prior to use for not less than 10 minutes, creating turbulence at the bottom of the tank that is visible at the top.

**Caution**

Do not allow horses or other equines access to feeds containing laidlomycin propionate potassium.  
The safety of laidlomycin propionate potassium in unapproved species has not been established.  
Not for use in animals intended for breeding.

**Manufactured By:**

Blue Bird Feed Mill  
City, State, Zip

**NET WEIGHT ON BAG OR BULK**

**Expiration Date: [21 days from date of manufacture]**

---

\*The final printed feed label must show actual concentration of laidlomycin propionate potassium.