

Desktop 300x250



Frame 1



Frame 2



Frame 3



Frame 4



Frame 5



Frame 6

Links to: <https://www.rexulti.com/depression-symptom-questionnaire>

Desktop 300x250 | Ad Choice




Frame 1



Frame 2



Frame 3



Frame 4



Frame 5



Frame 6

Important Safety Information, 300x250

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressants to treat major depressive disorder (MDD) in adults.
It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.
Increased risk of suicidal thoughts and actions. REXULTI and antidepressants may increase suicidal thoughts and actions in some people 24 years of age and younger, especially within the first few months of treatment or when the dose is changed. Depression and other mental illnesses are the most important causes of suicidal thoughts and actions. Patients on antidepressants and their families or caregivers should watch for new or worsening depression symptoms, especially sudden changes in mood, behaviors, thoughts, or feelings. Report any change in these symptoms immediately to the doctor.

Do not take REXULTI if you are allergic to brexpiprazole or any of the ingredients in REXULTI.

REXULTI may cause serious side effects, including:

- Cerebrovascular problems, including stroke, in elderly people with dementia-related psychosis that can lead to death.**
- Neuroleptic malignant syndrome (NMS) is a serious condition that can lead to death. Call your healthcare provider or go to the nearest hospital emergency room right away if you have some or all of the following signs and symptoms of NMS: high fever; changes in your pulse, blood pressure, heart rate, and breathing; stiff muscles; confusion; increased sweating.**
- Uncontrolled body movements (tardive dyskinesia). REXULTI can cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop taking REXULTI. Tardive dyskinesia may also start after you stop taking REXULTI.**
- Problems with your metabolism such as:**
 - high blood sugar (hyperglycemia) and diabetes.** REXULTI may cause high blood sugar. Some people who take REXULTI. Extremely high blood sugar can lead to coma or death. Your healthcare provider should check your blood sugar before you start, or soon after you start REXULTI and then regularly during long term treatment with REXULTI.
Call your healthcare provider if you have any of these symptoms of high blood sugar during treatment with REXULTI:
 - feel very thirsty
 - feel very hungry
 - feel sick to your stomach
 - need to urinate more than usual
 - feel weak or tired
 - feel confused, or your breath smells fruity
 - increased fat levels (cholesterol and triglycerides)** In your blood. Your healthcare provider should check the fat levels in your blood before you start, or soon after you start REXULTI, and then periodically during treatment with REXULTI.
 - weight gain.** You and your healthcare provider should check your weight before you start and often during treatment with REXULTI.
- Unusual and uncontrollable (compulsive) urges.** Some people taking REXULTI have had strong unusual urges, to gamble and gambling that cannot be controlled (compulsive gambling). These urges may include sexual urges, shopping, and eating or binge eating. If you or your family members notice that you are having new or unusual strong urge or behaviors, talk to your healthcare provider.
- Low white blood cell count.** Your healthcare provider may do blood tests during the first few months of treatment with REXULTI.
- Decreased blood pressure (orthostatic hypotension) and fainting.** You may feel dizzy, lightheaded or pass out (faint) when you rise too quickly from a sitting or lying position.
- Falls.** REXULTI may make you sleepy or dizzy, may cause a decrease in your blood pressure when changing position (orthostatic hypotension), and can slow your thinking and motor skills which may lead to falls that can cause fractures or other injuries.
- Seizures (convulsions).**
- Problems controlling your body temperature so that you feel too warm.** Do not become too hot or dehydrated during treatment with REXULTI. Do not exercise too much. In hot weather, stay inside in a cool place if possible. Stay out of the sun. Do not wear too much clothing or heavy clothing. Drink plenty of water.
- Difficulty swallowing that can cause food or liquid to get into your lungs.**
- Sleepiness, drowsiness, feeling tired, difficulty thinking and doing normal activities.** Do not drive a car, operate machinery, or do other dangerous activities until you know how REXULTI affects you. REXULTI may make you feel drowsy.

Before taking REXULTI, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had heart problems or a stroke
- have or have had low or high blood pressure
- have or have had diabetes or high blood sugar or a family history of diabetes or high blood sugar. Your healthcare provider should check your blood sugar before you start REXULTI and during treatment with REXULTI.
- have or have had high levels of total cholesterol, LDL cholesterol, or triglycerides, or low levels of HDL cholesterol
- have or have had seizures (convulsions)
- have or have had kidney or liver problems
- have or have had a low white blood cell count
- are pregnant or plan to become pregnant. REXULTI may harm your unborn baby. Taking REXULTI during your third trimester of pregnancy may cause your baby to have abnormal muscle movements, such as fetal拳形手 (fetal拳形手) after birth. Talk to your healthcare provider about the risk to your unborn baby if you take REXULTI during pregnancy.
 - Tell your healthcare provider if you become pregnant or think you are pregnant during treatment with REXULTI.
 - There is a pregnancy exposure registry for women who are exposed to REXULTI during pregnancy. If you become pregnant during treatment with REXULTI, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics. You can register by calling 1-866-961-2388 or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>.
- are breastfeeding or plan to breastfeed. It is not known if REXULTI passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with REXULTI.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. REXULTI and other medicines may affect each other causing possible serious side effects. REXULTI may affect how other medicines work, and other medicines may affect how REXULTI works. Your healthcare provider can tell you if it is safe to take REXULTI with your other medicines. Do not start or stop any medicines during treatment with REXULTI without first talking to your healthcare provider.

The most common side effects of REXULTI include weight gain, sleepiness, dizziness, common cold symptoms, and restlessness or feeling like you need to move (akathisia). These are not all the possible side effects of REXULTI. For more information, ask your healthcare provider or pharmacist. You are encouraged to report side effects of REXULTI (brexpiprazole). Please contact Otsuka America Pharmaceutical, Inc. at 1-800-438-9927 or FDA at 1-800-FDA-1088 (www.fda.gov/medwatch).

Please read FULL PRESCRIBING INFORMATION, including BOXED WARNING, and MEDICATION GUIDE.

© 2023 Otsuka America Pharmaceutical, Inc.
May 2023 11US23EBC0168



WHEN ADDED TO AN ANTIDEPRESSANT,

ANTIDEPRESSANT

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.

WHEN ADDED TO AN ANTIDEPRESSANT, REXULTI WAS PROVEN TO PROVIDE A

62% GREATER REDUCTION IN DEPRESSION SYMPTOMS

ANTIDEPRESSANT + REXULTI
brexpiprazole 2mg tablets

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.

WHEN ADDED TO AN ANTIDEPRESSANT, REXULTI WAS PROVEN TO PROVIDE A

62% GREATER REDUCTION IN DEPRESSION SYMPTOMS VS. THE ANTIDEPRESSANT ALONE.*

*6 week study in adults, with ongoing symptoms of depression, still taking an antidepressant. Individual results may vary.

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.

When depression sets you back, Keep Moving Forward.

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.

REXULTI
brexpiprazole 2mg tablets

Ask your doctor today about adding REXULTI to your antidepressant.

ACTOR PORTRAYAL

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.

Frame 1

Frame 2

Frame 3

Frame 4

Frame 5

Frame 6



WHEN ADDED TO AN ANTIDEPRESSANT,

ANTIDEPRESSANT

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.

Increased risk of suicidal thoughts and actions

WHEN ADDED TO AN ANTIDEPRESSANT, REXULTI WAS PROVEN TO PROVIDE A

62% GREATER REDUCTION IN DEPRESSION SYMPTOMS

ANTIDEPRESSANT + REXULTI® brexpiprazole 2mg tablets

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.

Increased risk of suicidal thoughts and actions

WHEN ADDED TO AN ANTIDEPRESSANT, REXULTI WAS PROVEN TO PROVIDE A

62% GREATER REDUCTION IN DEPRESSION SYMPTOMS VS. THE ANTIDEPRESSANT ALONE.*

*6 week study in adults, with ongoing symptoms of depression, still taking an antidepressant. Individual results may vary.

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.

Increased risk of suicidal thoughts and actions

When depression sets you back, Keep Moving Forward.

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.

Increased risk of suicidal thoughts and actions

REXULTI® brexpiprazole 2mg tablets

Ask your doctor today about adding REXULTI to your antidepressant.

ACTOR PORTRAYAL

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.

Increased risk of suicidal thoughts and actions

Frame 1

Frame 2

Frame 3

Frame 4

Frame 5

Frame 6

Important Safety Information, 300x600

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.
It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.
- Increased risk of suicidal thoughts and actions. REXULTI and antidepressant medicines may increase suicidal thoughts and actions in some people 24 years of age and younger, especially within the first few months of treatment or when the dose is changed. Depression and other mental illnesses are the most important causes of suicidal thoughts and actions. Patients, families, and caregivers should watch for new or worsening depression symptoms, especially sudden changes in mood, behaviors, thoughts, or feelings. Report any change in these symptoms immediately to the doctor.

Do not take REXULTI if you are allergic to brexpiprazole or any of the ingredients in REXULTI.

REXULTI may cause serious side effects, including:

- Cerebrovascular problems, including stroke, in elderly people with dementia-related psychosis that can lead to death.**
- Neuroleptic malignant syndrome (NMS) is a serious condition that can lead to death. Call your healthcare provider or go to the nearest hospital emergency room right away if you have some or all of the following signs and symptoms of NMS: high fever; changes in your pulse, blood pressure, heart rate, and breathing; stiff muscles; confusion; increased sweating.**
- Uncontrolled body movements (tardive dyskinesia). REXULTI may cause movements that you cannot control in your face, torso, and other body parts. Tardive dyskinesia may not go away, even if you stop taking REXULTI. Tardive dyskinesia may also start after you stop taking REXULTI.**
- Problems with your metabolism such as:**
 - increased fat levels (cholesterol and triglycerides) in your blood.** Your healthcare provider should check the fat levels in your blood before you start, or soon after you start REXULTI, and then periodically during treatment with REXULTI.
 - weight gain.** You and your healthcare provider should check your weight before you start and often during treatment with REXULTI.
- Unusual and uncontrollable (compulsive) urges.** Some people taking REXULTI have had strong unusual urges, to gamble and gambling that cannot be controlled (compulsive urges). Other uncontrollable urges include sexual urges, shopping, and eating or binge-eating. If you or your family members notice that you are having new or unusual strong urges or behaviors, talk to your healthcare provider.
- Low white blood cell count.** Your healthcare provider may do blood tests during the first few months of treatment with REXULTI.
- Decreased blood pressure (orthostatic hypotension) and fainting.** You may feel dizzy, lightheaded or pass out (faint) when you rise too quickly from a sitting or lying position.
- Falls.** REXULTI may make you sleepy or dizzy, may cause a decrease in your blood pressure when changing position (orthostatic hypotension), and can slow your thinking and motor skills which may lead to falls that can cause fractures or other injuries.
- Seizures (convulsions).**
- Problems controlling your body temperature so that you feel too warm or too cold.** Some people are hot or hydrated during treatment with REXULTI. Do not exercise too much. In hot weather, stay inside in a cool place if possible. Stay out of the sun. Do not wear too much clothing or heavy clothing. Drink plenty of water.
- Difficulty swallowing that can cause food or liquid to get into your lungs.**
- Sleepiness, drowsiness, feeling tired, difficulty thinking and doing normal activities.** Do not drive a car, operate machinery, or do other dangerous activities until you know how REXULTI affects you. REXULTI may make you feel drowsy.

Before taking REXULTI, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had heart problems or a stroke
- have or have had low or high blood pressure
- have or have had diabetes or high blood sugar or a family history of diabetes or high blood sugar. Your healthcare provider should check your blood sugar before you start REXULTI and during treatment with REXULTI.
- have or have had high levels of total cholesterol, LDL cholesterol, triglycerides, or low levels of HDL cholesterol
- have or have had seizures (convulsions)
- have or have had kidney or liver problems
- have or have had a low white blood cell count
- are pregnant or plan to become pregnant. REXULTI may harm your unborn baby. Taking REXULTI during your third trimester of pregnancy may cause your baby to have abnormal muscle movements or withdrawal symptoms after birth. Talk to your healthcare provider about the risk to your unborn baby if you take REXULTI during pregnancy.
- Tell your healthcare provider if you become pregnant or think you are pregnant during treatment with REXULTI.
- There is a pregnancy exposure registry for women who are exposed to REXULTI during pregnancy. If you become pregnant while taking treatment with REXULTI, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics. You can register by calling 1-866-961-2388 or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>.
- are breastfeeding or plan to breastfeed. It is not known if REXULTI passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with REXULTI.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Some other medicines may affect each other causing possible serious side effects. REXULTI may affect the way other medicines work, and other medicines may affect how REXULTI works. Your healthcare provider can tell you if it is safe to take REXULTI with your other medicines. Do not start or stop any medicine during treatment with REXULTI without first talking to your healthcare provider.

The most common side effects of REXULTI include weight gain, sleepiness, dizziness, common cold symptoms, and restlessness or feeling like you need to move (akathisia). These are not all the possible side effects of REXULTI. For more information, ask your healthcare provider or pharmacist. You are encouraged to report side effects of REXULTI (brexpiprazole) to the FDA. Otsuka America Pharmaceutical, Inc. at 1-800-437-9927 or FDA at 1-800-FDA-1088 www.fda.gov/medwatch.

Please read FULL PRESCRIBING INFORMATION, including BOXED WARNING, and MEDICATION GUIDE.

 © 2023 Otsuka America Pharmaceutical, Inc.
May 2023 11US23EBC0168

Desktop 160x600



On an antidepressant but still putting on a brave face?



WHEN ADDED TO AN ANTIDEPRESSANT,

ANTIDEPRESSANT

INDICATION and IMPORTANT SAFETY INFORMATION

REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise



WHEN ADDED TO AN ANTIDEPRESSANT, REXULTI WAS PROVEN TO PROVIDE A

62%
GREATER REDUCTION
IN DEPRESSION SYMPTOMS

ANTIDEPRESSANT + REXULTI[®] brexpiprazole 3mg tablets


INDICATION and IMPORTANT SAFETY INFORMATION

REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise



WHEN ADDED TO AN ANTIDEPRESSANT, REXULTI WAS PROVEN TO PROVIDE A

62%
GREATER REDUCTION
IN DEPRESSION SYMPTOMS
VS. THE ANTIDEPRESSANT ALONE.*

*6 week study in adults, with ongoing symptoms of depression, still taking an antidepressant. Individual results may vary.

INDICATION and IMPORTANT SAFETY INFORMATION

REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise



When depression sets you back,
Keep Moving Forward.



INDICATION and IMPORTANT SAFETY INFORMATION

REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise



REXULTI[®]
brexpiprazole 3mg tablets
Ask your doctor today about adding REXULTI to your antidepressant.



Take the symptom questionnaire to get started.

ACTOR PORTRAYAL

INDICATION and IMPORTANT SAFETY INFORMATION

REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- Increased risk of death in elderly people with dementia-related psychosis. Medicines like REXULTI can raise

Frame 1

Frame 2

Frame 3

Frame 4


Frame 5

Frame 6

Desktop 160x600 | Ad Choice




Frame 1



Frame 2



Frame 3



Frame 4



Frame 5



Frame 6

Important Safety Information, 160x600

**INDICATION and
IMPORTANT SAFETY
INFORMATION**
REXULTI is a prescription
medicine used with
antidepressant medicines to
treat major depressive disorder
(MDD). It is not known if REXULTI is
safe and effective in children
with MDD.

**IMPORTANT SAFETY
INFORMATION:**

- Increased risk of death
in elderly people with
dementia-related
psychosis.** Medicines
used to treat MDD may
increase the risk of death in
elderly people who have
dementia-related psychosis
(psychosis) due to
confusion and memory
loss. These medicines, including REXULTI,
are not approved for the
treatment of people with
dementia-related psychosis
without agitation that may
occur with Alzheimer's
disease. The increased risk of
death is not known if REXULTI
is used in elderly people
without dementia due to
Alzheimer's disease.
The increased risk of suicidal
thoughts and actions,
REXULTI and other
antidepressant
medicines may increase
suicidal thoughts and
actions in people 24 years of age and
younger, especially
within the first 4 to 12
months of treatment or
when the dose is
increased. Depression
and other mental
illnesses are the most
common causes of
suicidal thoughts and
actions. Patients on
REXULTI should watch
for new or worsening
depression symptoms,
especially sudden
changes in mood,
behaviors, thoughts, or
feelings. Report any
change in these
symptoms immediately
to the doctor.

**Do not take REXULTI if you
are allergic to brexpiprazole
or any other ingredients
in REXULTI.**

**REXULTI may cause serious
side effects, including:**

- Cerebrovascular
events, including
stroke, in elderly people
with dementia-related
psychosis that can lead
to death.**
- Neuroleptic malignant
syndrome, a serious
condition that
can lead to death. Call
your healthcare provider
or go to the nearest
hospital emergency room
right away if you have
some or all of the
following signs and
symptoms: headache, high
fever; changes in your
pulse, blood pressure,
heart rate, breathing;
stiff muscles; confusion;
increased sweating.**
- Uncontrollable
movements (tardive
dyskinesia). REXULTI may
cause movements that
you cannot control in
your face, tongue, or other
body parts. Tardive
dyskinesia may not go
away, even if you stop
taking REXULTI. Tardive
dyskinesia may also
start after you stop
taking REXULTI.**
- Problems with your
metabolism such as:
+ high blood sugar
(hyperglycemia) and
diabetes. An increase in
blood sugar can happen
in some people who take
REXULTI. High blood sugar
can lead to coma or death. Your
healthcare provider
should check your blood
sugar before you start,
or soon after you start
REXULTI, and then
regularly during
treatment with
REXULTI.**
- Call your healthcare
provider right away if
you have any of these symptoms of
high blood sugar during
treatment with REXULTI:
+ feel very thirsty
+ feel sick to your
stomach
+ need to urinate more
than usual
+ feel very tired
+ feel confused, or your
breath smells fruity
+ increased fat levels
(cholesterol and
triglycerides) in your
blood. Your healthcare
provider should check
the fat levels in your
blood before you start,
or soon after you start
REXULTI, and then
regularly during
treatment with
REXULTI.**
- + weight gain. You and
your healthcare provider
should check your
weight before you start
and often during
treatment with REXULTI.**
- Unusual and
uncontrollable
compulsive urges.
Some people taking
REXULTI have had strong
urges to gamble, sex, and
gambling that cannot
be controlled (compulsive
gambling). Other
compulsive urges include
sexual urges, shopping,
and eating uncontrollably.
If you or your family
members notice that you
are having strong
urges or behaviors,
talk to your healthcare
provider.**
- Low white blood cell
count. Your healthcare
provider should do blood
tests during the first few
months of treatment with
REXULTI.**
- Decreased blood
pressure (orthostatic
hypotension) and
fainting. You may feel
dizzy, lightheaded or pass
out (faint) when when you
rise from a sitting or
sitting or lying position.**
- Falls. REXULTI may make
you sleepy or dizzy, may
cause you to faint, or may
lower your blood pressure
when changing position
(orthostatic hypotension),
and can slow your thinking
(orthocognitive hypotension),
which may lead to falls that can
cause fractures or other injuries.**
- Seizures (convulsions).**
- Problems controlling
your body temperature
so that you feel too
warm. Do not become too
hot during treatment with
REXULTI. Do not exercise too much.
It is not known if REXULTI
can cause you to overheat
in a cool place if possible.
Stay out of the sun. Do
not wear many layers of
clothing or heavy clothing.
Drink plenty of water.**
- Difficulty breathing
that can cause fluid or liquid to
get into your lungs.**
- Sleepiness, drowsiness,
difficulty thinking and
difficulty doing
normal activities. Do not
drive a car, use machinery,
or do other
dangerous activities until
you know how REXULTI
affects you. REXULTI may
make you feel drowsy.**

**Before taking REXULTI, tell
your healthcare provider
about all of your medical
conditions, including if you:**

- have or have had
problems or a stroke.**
- have or have had low or
high blood sugar.**
- have or have had diabetes
or high blood sugar or a
family history of diabetes
or high blood sugar. Your
healthcare provider should
check your blood sugar
before you start REXULTI
and during treatment with
REXULTI.**
- have or have had high
cholesterol, LDL cholesterol,
triglycerides, or low levels
of HDL cholesterol.**
- have or have had
seizures (convulsions).**
- have or have had kidney
or liver problems.**
- have or have had a low
white blood cell count.**
- are pregnant or plan to
become pregnant.**

**REXULTI may affect your
unborn baby. Taking
REXULTI during your third
trimester of pregnancy
may cause your baby to
have abnormal muscle
movements and withdrawal
symptoms after birth. Talk
to your healthcare provider
about the risks of taking
REXULTI during pregnancy.**

**+ Tell your healthcare
provider if you become
pregnant while you are
pregnant during
treatment with REXULTI.**

**+ There is a pregnancy
registry for women who
are exposed to REXULTI during
pregnancy. If you are
pregnant during treatment with
REXULTI, talk to your healthcare
provider about registering
with the National
Pregnancy Registry for
Atypical Antipsychotics.
You can register by
calling 1-866-961-2388
or visit
[http://www.fda.gov/health
sciences/clinical-and-
research-programs/
pregnancyregistry](http://www.fda.gov/health
sciences/clinical-and-
research-programs/
pregnancyregistry)**

**+ Tell your healthcare provider
about all the medicines you
take, including prescription and
over-the-counter medicines,
vitamins, and dietary
supplements. REXULTI and
other medicines may affect
each other. This may cause
serious side effects. REXULTI
may affect the way other
medicines work. Your
healthcare provider can tell you
if it is safe to take REXULTI with
your other medicines. Do not
start or stop any medicine
during treatment with REXULTI
without talking to your
healthcare provider.**

**The most common side
effects of REXULTI include
drowsiness, dizziness, dry mouth,
dizziness, common cold
symptoms, and restlessness or
twitching that make you need to move
(akathisia).**

**These are not all the possible
side effects of REXULTI. For
more information, talk to your
healthcare provider or
pharmacist.**

**You are encouraged to report
side effects of REXULTI.
(brexpiprazole). Please contact
Otsuka America Pharmaceutical, Inc. at
1-800-338-9927 or FDA at
<http://www.fda.gov/medwatch>.**

**Please read FULL
PRESCRIBING
INFORMATION, including
BOXED WARNING, and
MEDICATION GUIDE.**

 © 2003 Otsuka America
Pharmaceuticals, Inc.
May 2003 11US3EBC0168


On an antidepressant but
still putting on a brave face?



Frame 1

WHEN ADDED TO AN ANTIDEPRESSANT,
ANTIDEPRESSANT

**WHEN ADDED TO AN ANTIDEPRESSANT,
REXULTI WAS PROVEN TO PROVIDE A** **62% GREATER REDUCTION**
IN DEPRESSION SYMPTOMS




REXULTI[®]
brexpiprazole
2mg tablets

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**,
and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with
antidepressant medicines to treat major depressive disorder
(MDD) in adults.

Frame 2

**WHEN ADDED TO AN ANTIDEPRESSANT,
REXULTI WAS PROVEN TO PROVIDE A** **62% GREATER REDUCTION**
IN DEPRESSION SYMPTOMS



REXULTI[®]
brexpiprazole
2mg tablets


FULL PRESCRIBING INFORMATION, including **BOXED WARNING**,
and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with
antidepressant medicines to treat major depressive disorder
(MDD) in adults.

Frame 3

**WHEN ADDED TO AN ANTIDEPRESSANT,
REXULTI WAS PROVEN TO PROVIDE A** **62% GREATER REDUCTION**
IN DEPRESSION SYMPTOMS
VS. THE ANTIDEPRESSANT ALONE.*

*6 week study in adults, with ongoing symptoms of depression,
still taking an antidepressant. Individual results may vary.




FULL PRESCRIBING INFORMATION, including **BOXED WARNING**,
and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with
antidepressant medicines to treat major depressive disorder
(MDD) in adults.

Frame 4



**When depression
sets you back,
Keep Moving
Forward.**



FULL PRESCRIBING INFORMATION, including **BOXED WARNING**,
and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with
antidepressant medicines to treat major depressive disorder
(MDD) in adults.

Frame 5




REXULTI[®]
brexpiprazole
2mg tablets

ACTOR PORTRAYAL



**Ask your doctor today about adding
REXULTI to your antidepressant.**



**Take the symptom
questionnaire to get started.**

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**,
and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with
antidepressant medicines to treat major depressive disorder
(MDD) in adults.

Frame 6

On an antidepressant but
still putting on a brave face?



Frame 1

WHEN ADDED TO AN ANTIDEPRESSANT,

ANTIDEPRESSANT

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

Frame 2

WHEN ADDED TO AN ANTIDEPRESSANT,
REXULTI WAS PROVEN TO PROVIDE A **62% GREATER REDUCTION**
IN DEPRESSION SYMPTOMS

ANTIDEPRESSANT + **REXULTI®**
brexpiprazole
2mg tablets

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

Frame 3

WHEN ADDED TO AN ANTIDEPRESSANT,
REXULTI WAS PROVEN TO PROVIDE A **62% GREATER REDUCTION**
IN DEPRESSION SYMPTOMS
VS. THE ANTIDEPRESSANT ALONE.*

*6 week study in adults, with ongoing symptoms of depression, still taking an antidepressant. Individual results may vary.

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

Frame 4



When depression
sets you back,
**Keep Moving
Forward.**

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

Frame 5



REXULTI®
brexpiprazole
2mg tablets
ACTOR PORTRAYAL

Ask your doctor today about adding
REXULTI to your antidepressant.

Take the symptom
questionnaire to get started.

FULL PRESCRIBING INFORMATION, including **BOXED WARNING**, and **MEDICATION GUIDE**.

INDICATION and IMPORTANT SAFETY INFORMATION
REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

Frame 6

Important Safety Information, 728x90

INDICATION and IMPORTANT SAFETY INFORMATION

REXULTI is a prescription medicine used along with antidepressant medicines to treat major depressive disorder (MDD) in adults.

It is not known if REXULTI is safe and effective in children with MDD.

IMPORTANT SAFETY INFORMATION:

- **Increased risk of death in elderly people with dementia-related psychosis.** Medicines like REXULTI can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). REXULTI is not approved for the treatment of people with dementia-related psychosis without agitation that may happen with dementia due to Alzheimer's disease.
- **Increased risk of suicidal thoughts and actions.** REXULTI and antidepressant medicines may increase suicidal thoughts and actions in some people 24 years of age and younger, especially within the first few months of treatment or when the dose is changed. Depression and other mental illnesses are the most important causes of suicidal thoughts and actions. Patients on antidepressants and their families or caregivers should watch for new or worsening depression symptoms, especially sudden changes in mood, behaviors, thoughts, or feelings. Report any change in these symptoms immediately to the doctor.

Do not take REXULTI if you are allergic to brexpiprazole or any of the ingredients in REXULTI.

REXULTI may cause serious side effects, including:

- **Cerebrovascular problems, including stroke, in elderly people with dementia-related psychosis that can lead to death.**
- **Neuroleptic malignant syndrome (NMS) is a serious condition that can lead to death.** Call your healthcare provider or go to the nearest hospital emergency room right away if you have some or all of the following signs and symptoms of NMS: high fever; changes in your pulse, blood pressure, heart rate, and breathing; stiff muscles; confusion; increased sweating
- **Uncontrolled body movements (tardive dyskinesia).** REXULTI may cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop taking REXULTI. Tardive dyskinesia may also start after you stop taking REXULTI.
- **Problems with your metabolism such as:**
 - **high blood sugar (hyperglycemia) and diabetes.** Increases in blood sugar can happen in some people who take REXULTI. Extremely high blood sugar can lead to coma or death. Your healthcare provider should check your blood sugar before you start, or soon after you start REXULTI and then regularly during long term treatment with REXULTI.
 - **increased fat levels (cholesterol and triglycerides) in your blood.** Your healthcare provider should check the fat levels in your blood before you start, or soon after you start REXULTI, and then periodically during treatment with REXULTI.
 - **weight gain.** You and your healthcare provider should check your weight before you start and often during treatment with REXULTI.
- **Unusual and uncontrollable (compulsive) urges.** Some people taking REXULTI have had strong unusual urges, to gamble and gambling that cannot be controlled (compulsive gambling). Other compulsive urges include sexual urges, shopping, and eating or binge eating. If you or your family members notice that you are having new or unusual strong urges or behaviors, talk to your healthcare provider.
- **Low white blood cell count.** Your healthcare provider may do blood tests during the first few months of treatment with REXULTI.
- **Decreased blood pressure (orthostatic hypotension) and fainting.** You may feel dizzy, lightheaded or pass out (faint) when you rise too quickly from a sitting or lying position.
- **Falls.** REXULTI may make you sleepy or dizzy, may cause a decrease in your blood pressure when changing position (orthostatic hypotension), and can slow your thinking and motor skills which may lead to falls that can cause fractures or other injuries.
- **Seizures (convulsions).**
- **Problems controlling your body temperature so that you feel too warm.** Do not become too hot or dehydrated during treatment with REXULTI. Do not exercise too much. In hot weather, stay inside in a cool place if possible. Stay out of the sun. Do not wear too much clothing or heavy clothing. Drink plenty of water.
- **Difficulty swallowing** that can cause food or liquid to get into your lungs.
- **Sleepiness, drowsiness, feeling tired, difficulty thinking and doing normal activities.** Do not drive a car, operate machinery, or do other dangerous activities until you know how REXULTI affects you. REXULTI may make you feel drowsy.

Before taking REXULTI, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had heart problems or a stroke
- have or have had low or high blood pressure
- have or have had diabetes or high blood sugar or a family history of diabetes or high blood sugar. Your healthcare provider should check your blood sugar before you start REXULTI and during treatment with REXULTI.
- have or have had high levels of total cholesterol, LDL cholesterol, or triglycerides, or low levels of HDL cholesterol
- have or have had seizures (convulsions)
- have or have had kidney or liver problems
- have or have had a low white blood cell count
- are pregnant or plan to become pregnant. REXULTI may harm your unborn baby. Taking REXULTI during your third trimester of pregnancy may cause your baby to have abnormal muscle movements or withdrawal symptoms after birth. Talk to your healthcare provider about the risk to your unborn baby if you take REXULTI during pregnancy.
 - Tell your healthcare provider if you become pregnant or think you are pregnant during treatment with REXULTI.
- There is a pregnancy exposure registry for women who are exposed to REXULTI during pregnancy. If you become pregnant during treatment with REXULTI, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics. You can register by calling 1-866-961-2388 or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>.
- are breastfeeding or plan to breastfeed. It is not known if REXULTI passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with REXULTI.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. REXULTI and other medicines may affect each other causing possible serious side effects. REXULTI may affect the way other medicines work, and other medicines may affect how REXULTI works. Your healthcare provider can tell you if it is safe to take REXULTI with your other medicines. Do not start or stop any medicines during treatment with REXULTI without first talking to your healthcare provider.

The most common side effects of REXULTI include weight gain, sleepiness, dizziness, common cold symptoms, and restlessness or feeling like you need to move (akathisia).

These are not all the possible side effects of REXULTI. For more information, ask your healthcare provider or pharmacist.

You are encouraged to report side effects of REXULTI (brexpiprazole). Please contact Otsuka America Pharmaceutical, Inc. at 1-800-438-9927 or FDA at 1-800-FDA-1088 (www.fda.gov/medwatch).

Please read **FULL PRESCRIBING INFORMATION**, including **BOXED WARNING**, and **MEDICATION GUIDE**.

