

#284

# Using Relative Supersaturation to Support “Urinary Tract Health” Claims for Adult Maintenance Cat Food

## Guidance for Industry

### Draft Guidance

*This guidance document is being distributed for comment purposes only.*

Submit comments on this draft guidance by the date provided in the *Federal Register* notice announcing the availability of the draft guidance. Submit electronic comments to <https://www.regulations.gov>. Submit written comments to the Dockets Management Staff (HFA-305), Food and Drug Administration, 5630 Fishers Lane, Rm. 1061, Rockville, MD 20852. All comments should be identified with docket number FDA-2023-D-4095.

For further information regarding this document, contact Karen Donnelly, Center for Veterinary Medicine (HFV-227), U.S. Food and Drug Administration, 12225 Wilkins Avenue, Rockville MD 20852, 240-402-9802, [karen.donnelly2@fda.hhs.gov](mailto:karen.donnelly2@fda.hhs.gov).

Additional copies of this draft guidance document may be requested from the Policy and Regulations Staff (HFV-6), Center for Veterinary Medicine, Food and Drug Administration, 7500 Standish Place, Rockville MD 20855, and may be viewed on the Internet at <https://www.fda.gov/animal-veterinary>, <https://www.fda.gov/regulatory-information/search-fda-guidance-documents>, or <https://www.regulations.gov>.

**U.S. Department of Health and Human Services  
Food and Drug Administration  
Center for Veterinary Medicine (CVM)  
November 2023**

## **Table of Contents**

|              |   |           |
|--------------|---|-----------|
| <b>I.</b>    | <b>Introduction.....</b>  | <b>1</b>  |
| <b>II.</b>   | <b>Background .....</b>   | <b>1</b>  |
|              | <b>A. Introduction to Urolithiasis in Cats.....</b>                               | <b>1</b>  |
|              | <b>B. Relative Supersaturation (RSS) Methodology .....</b>                        | <b>2</b>  |
|              | <b>C. Support for Using RSS Methodology in UTH Cat Food.....</b>                  | <b>3</b>  |
| <b>III.</b>  | <b>Recommended Information to Substantiate RSS-Based UTH Claims .....</b>         | <b>4</b>  |
| <b>IV.</b>   | <b>Utility.....</b>   | <b>4</b>  |
|              | <b>A. Target RSS Values .....</b>   | <b>4</b>  |
|              | <b>B. Study Duration and RSS Assessment Schedule .....</b>                        | <b>5</b>  |
| <b>V.</b>    | <b>Target Animal Safety.....</b>  | <b>5</b>  |
|              | <b>A. Safety Concerns.....</b>  | <b>5</b>  |
|              | <b>B. RSS Study Design.....</b>   | <b>6</b>  |
|              | <b>C. Combined Safety and Utility Study.....</b>                                  | <b>9</b>  |
| <b>VI.</b>   | <b>References.....</b>  | <b>9</b>  |
| <b>VII.</b>  | <b>Appendix 1 – Recommended Combined Safety and Utility Study Flowchart .....</b> | <b>12</b> |
| <b>VIII.</b> | <b>Appendix 2 – Glossary of Terms and Acronyms .....</b>                          | <b>13</b> |

*Contains Nonbinding Recommendations*

*Draft — Not for Implementation*

## **Using Relative Supersaturation to Support “Urinary Tract Health” Claims for Adult Maintenance Cat Food**

### **Draft Guidance for Industry**

*This draft guidance, when finalized, will represent the current thinking of the Food and Drug Administration (FDA, we, or Agency) on this topic. It does not establish any rights for any person and is not binding on FDA or the public. You can use an alternative approach if it satisfies the requirements of the applicable statutes and regulations. To discuss an alternative approach, contact the FDA staff responsible for this guidance as listed on the title page.*

#### **I. Introduction**

In response to requests from pet food manufacturers, the FDA’s Center for Veterinary Medicine (CVM) has evaluated the use of relative supersaturation (RSS) methodology to support urinary tract health (UTH) claims for certain adult maintenance cat food. RSS is a measurement that estimates the potential for crystal formation and bladder stone (urolith) growth, which is a common affliction in cats. This guidance provides recommendations for how pet food manufacturers can use RSS methodology to substantiate general structure or function claims that an adult maintenance cat food supports urinary tract health by promoting a healthy mineral content in the urinary tract.

The recommendations we provide in this guidance include: (1) the wording of UTH claims based on RSS methodology for use on adult maintenance cat food labeling; (2) the RSS criteria to substantiate UTH claims; and (3) the study data CVM requests from pet food manufacturers that demonstrate the utility and target animal safety of the cat food.

For purposes of this guidance, “you” means the manufacturer of adult maintenance cat food with UTH claims (UTH cat food) based on RSS methodology.

In general, FDA’s guidance documents do not establish legally enforceable responsibilities. Instead, guidances describe the Agency’s current thinking on a topic and should be viewed only as recommendations, unless specific regulatory or statutory requirements are cited. The use of the word *should* in Agency guidance means that something is suggested or recommended, but not required.

#### **II. Background**

##### **A. Introduction to Urolithiasis in Cats**

Urolithiasis (the presence of uroliths in the bladder) is a fairly common affliction in cats, with prevalence estimates from clinical studies ranging from 9-23% (Forester *et al.* 2010). The two most common types of uroliths in cats are magnesium ammonium

## ***Contains Nonbinding Recommendations***

*Draft — Not for Implementation*

phosphate (struvite), and calcium oxalate (CaOx), accounting for 50% and 37% of analyzed uroliths, respectively (Minnesota Urolith Center Global Data 2022).

In response to concerns about urolithiasis, pet food manufacturers use various formulation strategies to make UTH cat food. Historically, manufacturers of UTH cat food restricted the magnesium content and/or formulated their cat food to produce slightly acidic urine (pH of 5.9 to 6.4). The slight acidity and low magnesium content create a urinary environment that is unfavorable for struvite crystallization and urolith growth but may be favorable for CaOx urolith growth in predisposed cats (Lekcharoensuk *et al.* 2001). Formulating cat food based on RSS methodology is a more recent dietary strategy that some pet food manufacturers use to create UTH cat food. The principles of RSS apply to all urolith types, so UTH cat food based on RSS methodology has the potential to help protect cats from both struvite and CaOx urolithiasis.

### **B. Relative Supersaturation (RSS) Methodology**

RSS methodology originated in human medicine in the 1960's and is widely used for measuring the potential for urolith formation in humans. Researchers have also validated RSS methodology for use in dogs and cats (Robertson *et al.* 2002).

One of the primary conditions for urolith formation in any species is oversaturation of the urine with dissolved substances (solutes) that have the potential to precipitate out of solution and form crystals. These crystals can eventually grow into uroliths. RSS provides a numerical measurement of the degree of saturation of a specific urolith-forming substance, and thus a quantitative method to evaluate the risk of urolith formation. There is a unique RSS value for each possible urolith type in a urine sample. The RSS value for a specific urolith is determined by first measuring the concentration of multiple solutes, including magnesium, ammonium, phosphate, calcium, and oxalate, as well as the pH in a urine sample collected over at least 24 hours. These data are entered into a computer program that calculates the concentrations of the solutes that remain free to interact and form crystals, also known as the activity product (AP). An RSS value is obtained by dividing the AP for a given urolith-forming compound by its known thermodynamic solubility product (SP), which represents the maximum concentration at which a solute can be completely dissolved in solution.

$$\text{RSS} = \text{AP}_{\text{urolith of the urine}} / \text{SP}_{\text{urolith}}$$

There are three zones of urine saturation that describe the activity of a given crystal in a urine sample:

- (1) *Undersaturation* – crystals will not spontaneously form, and if present, will dissolve.
- (2) *Metastable supersaturation* – crystals will not spontaneously form but may form in the presence of solid particles (i.e., a nucleus) such as cells, an existing urolith, or other particulate matter. Existing crystals and uroliths will not dissolve and may grow.

***Contains Nonbinding Recommendations***  
*Draft — Not for Implementation*

(3) *Labile supersaturation* – crystals will spontaneously form and grow.

The calculated RSS value for a given crystal indicates which zone its concentration is in for that urine sample. An RSS value below the solubility product for a given crystal, i.e., RSS less than 1.0, puts it in the undersaturation zone. The concentration at which dissolved urolith material will spontaneously precipitate out of solution to form crystals is called the formation product, and the formation product defines the limit between the metastable and labile supersaturation zones. Studies suggest that the formation product for struvite uroliths is an RSS of 2.5, and for CaOx uroliths is an RSS of 12 (Houston and Elliot 2008; Bartges 2011), and these values are generally accepted and used by the veterinary community. The degree to which an RSS value falls above the solubility product and approaches or exceeds the formation product indicates the level of risk for urolith formation.

### **C. Support for Using RSS Methodology in UTH Cat Food**

We received requests to assess the use of RSS methodology to substantiate UTH claims on adult maintenance cat food not restricted to sale through or under the guidance of veterinarians. We therefore reviewed the available literature regarding RSS in cats, and we summarize our findings from that literature review below.

#### **1. RSS and Dietary Factors**

Many studies investigate the relationship between RSS values and various dietary factors that possibly influence UTH in cats, such as ingredients that alter urine pH (Bartges 2013; Queau *et al.* 2013; Stevenson *et al.* 2000; Markwell *et al.* 1999; Smith *et al.* 1998), dietary moisture (Buckley *et al.* 2011), and dietary sodium content (Queau *et al.* 2020; Hawthorne and Markwell 2004). These studies help pet food manufacturers, nutritionists, and diet formulators to make cat food that produce the desired RSS values.

#### **2. RSS and Urolith Activity**

There are numerous studies that evaluate the utility of RSS methodology that specifically assess the relationship between RSS values and struvite urolithiasis in cats. Several studies demonstrated that cat urine undersaturated for struvite ( $RSS_{\text{Struvite}} < 1$ ) successfully dissolved struvite uroliths *in-vitro*, and the smaller the RSS value, the faster the dissolution occurred (Tournier *et al.* 2008; Tournier and Biourge 2008; Sagols *et al.* 2015). Studies also show that RSS is better than pH for predicting struvite urolith activity *in-vitro* (Van Hoek *et al.* 2009, Sagols *et al.* 2015). Published *in-vivo* studies demonstrated that cats that consumed diets formulated to induce urine undersaturated for struvite experienced complete struvite urolith dissolution after 2 to 10 weeks of feeding (Houston *et al.* 2004, 2011; Tefft *et al.* 2021; Torres-Henderson *et al.* 2017).

Although researchers cannot perform dissolution studies for CaOx uroliths because this urolith is not able to be dissolved, several studies document a positive association

## ***Contains Nonbinding Recommendations***

*Draft — Not for Implementation*

between greater  $RSS_{CaOx}$  values and CaOx urolith formation in the bladder and/or kidney in humans (Ferraro *et al.* 2018; Prochaska *et al.* 2018; Parks *et al.* 1997; Thun and Schober 1991), rats (Bushinsky *et al.* 1995), and dogs (Stevenson *et al.* 2003, 2004). We conclude that this information about the relationship between RSS and CaOx uroliths in other species and the results from the struvite dissolution studies support the use of RSS methodology to substantiate claims that a cat food promotes UTH in cats.

### **III. Recommended Information to Substantiate RSS-Based UTH Claims**

If you use RSS methodology to substantiate general structure or function claims that an adult maintenance cat food supports UTH, we recommend that prior to marketing the product, you submit for our review:

- Empirical data demonstrating the utility and safety of the product’s mechanism(s) for maintaining urinary tract health (See sections [IV. Utility](#) and [V. Target Animal Safety](#)),
- A complete quantitative ingredient formulation of the product,
- A complete quantitative nutrient composition of the product, and
- A complete product label, which can be a pdf-file, printer’s proof, or an unused label.

If you include an RSS-supported UTH claim on the product label of the adult maintenance cat food, you also should include a mechanism of action statement on the product label and any additional labeling. We recommend you use the mechanism of action statement: “Formulated to promote a healthy mineral content in the urinary tract.”

The submitted information helps CVM ensure the UTH claim is substantiated, and helps CVM review the claim in the context of the entire product label and labeling as a whole.

### **IV. Utility**

#### **A. Target RSS Values**

To demonstrate utility of a UTH cat food based on RSS methodology, you should submit results to CVM from at least one feeding study that demonstrates cats eating the food achieve RSS values below the formation products for struvite and CaOx (2.5 and 12, respectively). The study should evaluate RSS for both struvite and CaOx since they are the two most common types of uroliths found in cats and preventing one urolith type while potentially promoting formation of the other is not supportive of a general UTH claim. To ensure that cats consuming the food are less likely to develop urinary crystals or uroliths, CVM’s recommended mean RSS for each urolith type is no greater than the midpoint of that urolith’s metastable supersaturation range. In addition, the upper bound of the 95% confidence interval (CI) around the mean should be no greater than the value of the formation product for the specific urolith. The values meeting the recommended mean RSS and 95% CI limits are:

## ***Contains Nonbinding Recommendations***

*Draft — Not for Implementation*

- Struvite: Mean RSS  $\leq 1.8$ ; Upper bound of 95% CI  $\leq 2.5$
- CaOx: Mean RSS  $\leq 6.0$ ; Upper bound of 95% CI  $\leq 12$

These values allow for some spread in individual cat RSS values and still provide an acceptable margin of safety against urolith formation. If the RSS analysis provides separate RSS values for each of the two forms of CaOx crystals – monohydrate and dihydrate – rather than providing an RSS value for total CaOx, then the values for recommended mean RSS and CI limit apply to each of those forms.

We also recommend you conduct RSS measurements on urine collected over at least 24 hours, which would likely provide more accurate results because it accounts for possible diurnal variation in mineral excretion and other parameters.

### **B. Study Duration and RSS Assessment Schedule**

Although we do not define a specific timepoint when a study group of cats should achieve the target RSS values, the study should demonstrate that once the cats do achieve the RSS targets, the cats continue to meet the RSS targets from that point of the study onward. Data from industry studies measuring RSS submitted to CVM demonstrate that RSS is stable from 5 to 26 weeks in cats. Therefore, we recommend you conduct utility studies for a minimum of 40 days with RSS for struvite and CaOx measured at prescribed intervals no less than every 10 days during the study. The first measurement should occur between day 10 and 21 of feeding the product. There should be at least four separate RSS measurements, but they need not occur at regularly spaced intervals. For example, the study could run for 40 days with RSS measurements on days 10, 20, 30, and 40, or the study could run for 120 days with measurements on days 10, 20, 30, 60, and 120. Other study durations and schedules are also possible, but we recommend you use the guidelines we have described in this paragraph when designing your study.

It is important to note that although we request a minimum of four RSS measurements that meet the criteria listed in section [IV.A. Target RSS Values](#) to demonstrate utility, this does not mean that meeting the utility criteria at any four times out of the total number of measurements performed would be acceptable. The study should demonstrate that cats can *consistently* produce urine that meets the RSS targets, and that consistency should be demonstrated by at least four consecutive measurements. Although it is your decision to determine how long a study and how many RSS measurements you wish to perform to meet that goal, you should make that decision prior to study initiation and adhere to that schedule regardless of the RSS results obtained.

## **V. Target Animal Safety**

### **A. Safety Concerns**

There may be several methods for formulating a cat food that will produce the desired RSS parameters, so safety concerns will vary with the formulation method(s) used. One safety concern is the degree to which the cat food acidifies the urine, because creation of

## ***Contains Nonbinding Recommendations***

*Draft — Not for Implementation*

an overly acidic urine can increase the risk of metabolic acidosis. Although acidification may not be the primary mechanism pet food manufacturers employ when formulating UTH cat food based on RSS methodology, urinary pH is one of the variables that affects RSS values, so adjusting a dietary formula to affect RSS may inherently affect urinary pH as well. Thus, urinary pH measurements are an important safety component for manufacturers of UTH cat food when relying on RSS methodology. Data from studies submitted to CVM by pet food manufacturers show a correlation between very low urine pH and negative balance of magnesium (Mg) and potassium (K). Thus, if urinary pH is low enough to cause metabolic acidosis, mineral balance data (calcium (Ca), phosphorus (P), Mg, and K) are needed to evaluate product safety. This is addressed in more detail in section [V.B. RSS Study Design](#).

Because manufacturers of UTH cat food based on RSS methodology may use various, potentially novel, formulation methods to achieve the RSS goals, nutritional adequacy of the cat food is another safety concern. UTH cat food based on RSS methodology should therefore meet at least one, and preferably both, of the Association of American Feed Control Officials (AAFCO) methods for substantiation of nutritional adequacy (AAFCO, 2022). These methods are: (1) formulate the cat food to meet the AAFCO Cat Food Nutrient Profile for Adult Maintenance, and/or (2) successfully complete and pass an appropriate AAFCO Protocol Feeding Study that demonstrates the cat food to be adequate for maintaining the nutritional status of adult cats.

### **B. RSS Study Design**

#### **1. Duration**

If the cat food is adequate for maintaining the nutritional status of adult cats because you formulated the cat food to meet the AAFCO Cat Food Nutrient Profile for Adult Maintenance or completed an appropriate AAFCO Protocol Feeding Study, we recommend you conduct at least a 40-day feeding study to evaluate product safety. A study of longer duration may be necessary if mineral balance data is needed (see section [V.B.2. Urinary pH Measurements and Mineral Balance Data](#)). If the nutritional adequacy of the cat food has not been substantiated using either of the AAFCO methods, then you should conduct a 26-week adult maintenance feeding study as described in the AAFCO *Official Publication* (AAFCO, 2022) and collect safety data during that time.

#### **2. Urinary pH Measurements and Mineral Balance Data**

We recommend safety studies include measurements of each cat's urinary pH at a minimum of four timepoints no less than 10 days apart. Thus, a 40-day study should include urine pH measurements on days 10, 20, 30, and 40, but other schedules could be acceptable if the study is intended to run longer than 40 days. You should perform urine pH measurements on 24-hour urine samples (rather than on single point-in-time urine collections), and then calculate the group mean pH and 95% CI for each time point. The CI values will indicate if the study needs to include collection of mineral



## ***Contains Nonbinding Recommendations***

*Draft — Not for Implementation*

balance data (Ca, P, Mg, K), which would necessitate a study of longer duration, or if the study can end after 40 days. We recommend the following guidelines:

- a. If the lower bound of the 95% CI for the group mean pH is *equal to or greater than* 6.0 for at least three time points, the study can end after a minimum of 40 days.
- b. If the lower bound of the 95% CI for the group mean pH is *less than* 6.0 for two or more of the time points, the study should continue with mineral balance collections.
- c. You should start mineral balance collections as soon as they are determined to be needed and you should collect mineral balance data at a minimum of three timepoints no less than 30 days apart. A baseline collection prior to feeding the test diet is optional. For this collection to serve as one of the three needed timepoints, it should be performed only after all cats have been consuming the same complete and balanced diet (e.g., standard colony diet) for a minimum of 7 days.

Additional urinary pH and/or RSS measurements could be made if the study is extended to allow for mineral balance collections. We do not consider these additional measurements to be necessary if the study already included the recommended four RSS measurements, but the additional pH and RSS data should be included in the final study report if they are collected.

### **3. Additional Clinical Data**

Regardless of the duration, studies that adequately address target animal safety should also include:

- a. Veterinary physical exams performed at the beginning and end of the study, and at the midpoint if the study duration is longer than 40 days,
- b. Body weights measured weekly and body condition scores determined at least monthly,
- c. Food consumption measured daily,
- d. Daily stool observations  $\pm$  periodic fecal scoring,
- e. Morbidity and mortality observations, with any animal that dies being necropsied and a cause of death determined, and

## ***Contains Nonbinding Recommendations***

*Draft — Not for Implementation*

- f. Serum chemistries,<sup>1</sup> hematologies,<sup>2</sup> and urinalyses<sup>3</sup> performed at the beginning and end of the study, and at the midpoint if the study is longer than 40 days.

### **4. Other Considerations**

- a. You should conduct appropriate statistical comparisons, using suitable methodology and numbers of animals to confidently (95%) detect statistically significant differences (10%) in safety parameters between baseline and later timepoints, should they exist.
- b. Successful studies should have no more than 25% of the cats starting the study removed for non-nutritional reasons or poor food intake, and cats may be removed for poor food consumption and their data excluded from the final analysis only within the first 14 days of the study. The reason for any cat's removal should be recorded and included in the final study report. A minimum of 10 cats should complete the study, so we recommend at least 12 cats start the study to ensure the minimum number of cats complete the study.
- c. Any medical treatment provided to cats on study and the reason for it should be recorded and included in the final study report.
- d. The final study report should include all data generated from individual animals as well as summary statistics for each day of measurement. The study report should also include a full discussion of the diet formulation, experimental methodology, statistical methods used to analyze the data, and interpretation of findings.
- e. If feasible, we recommend you keep animals on study for up to 26 weeks and continue to collect blood and urine samples while we review the submitted data, as we may request additional data and/or a longer duration of study based on the study findings. Aliquots from collected blood and urine samples should be frozen and stored for possible additional testing.
- f. This guidance document may not address all specifics important for the design of the study. We encourage you to submit protocols to us for review and comment prior to study execution. You should address our questions on details of the protocol before the study begins.

---

<sup>1</sup> A serum biochemistry profile should include glucose, albumin, total protein, total bilirubin, conjugated bilirubin, blood urea nitrogen, creatinine, alkaline phosphatase, alanine transaminase, aspartate aminotransferase, gamma-glutamyltransferase, calcium, magnesium, phosphorus, potassium, sodium, chloride, and tCO<sub>2</sub> or pCO<sub>2</sub>.

<sup>2</sup> Hematology profile should include a complete blood count and differential white cell count.

<sup>3</sup> Urinalysis should include color, protein, glucose, ketones, bilirubin, specific gravity (by refractometer), and microscopic sediment examination to look for crystals, casts, red and white blood cells, and epithelial cells.

***Contains Nonbinding Recommendations***  
*Draft — Not for Implementation*

### **C. Combined Safety and Utility Study**

Because we recommend you conduct both target animal safety and utility studies for a minimum of 40 days, you may opt to combine both studies in one feeding trial to maximize efficiency of time and use of resources. [Appendix 1](#) contains a flowchart that depicts the recommended timeline for a minimum 40-day combined safety and utility study. Please note that this flowchart applies only to cat food for which you have substantiated the nutritional adequacy for adult maintenance, as recommended above. UTH cat food based on RSS methodology that has not been substantiated for nutritional adequacy to maintain adult cats should undergo a 6-month safety study, during which nutritional adequacy for adult maintenance can be substantiated according to AAFCO protocols along with the other parameters discussed in this guidance.

### **VI. References**

1. Association of American Feed Control Officials. (2022). *2022 Official Publication of the Association of American Feed Control Officials*. Champaign, IL 61820. <https://www.aaftco.org/publications> (accessed October 31, 2023).
2. Bartges J. (2011) Urinary saturation testing. In: Bartges, J. W., Polzin, D. J., eds. *Nephrology and urology of small animals*. Chichester, UK: Blackwell Publishing Ltd.: 75-85.
3. Bartges JW, Kirk CA, Cox SK, Moyers TD. (2013) Influence of acidifying or alkalinizing diets on bone mineral density and urine relative supersaturation with calcium oxalate and struvite in healthy cats. *Am J Vet Res* 74: 1347-1352.
4. Buckley MF, Hawthorne A, Colyer A, Stevenson AE. (2011) Effect of dietary water intake on urinary output, specific gravity and relative supersaturation for calcium oxalate and struvite in the cat. *Br J Nutr* 106: S128-S130.
5. Bushinsky DA, Grynpas MD, Nilsson EL, Nakagawa Y, Coe FL. (1995) Stone formation in genetic hypercalciuric rats. *Kidney Int* 48: 1705-1713.
6. Ferraro PM, Ticinesi A, Meschi T, Rodgers A, Di Maio F, Fulignati P, Borghi L, Gambaro G. (2018) Short-term changes in urinary relative supersaturation predict recurrence of kidney stones: A tool to guide preventive measures in urolithiasis. *J Urology* 200: 1082-1087.
7. Forrester SD, Kruger JM, Allen TA. (2010) Feline Lower Urinary Tract Diseases. In: Hand.M.S., Thatcher, C.D., Remillard, R. L., et al, eds. in *Small Animal Clinical Nutrition*. 5<sup>th</sup> Ed. Topeka, Kan: Mark Morris Institute: 927-976.
8. Hawthorne AJ, Markwell PJ. (2004) Dietary sodium promotes increased water intake and urine volume in cats. *J Nutr* 134: 2128S-2129S.

## ***Contains Nonbinding Recommendations***

*Draft — Not for Implementation*


9. Houston DM, Rinkardt NE, Hilton J. (2004) Evaluation of the efficacy of a commercial diet in the dissolution of feline struvite bladder uroliths. *Vet Ther* 5 (3): 187-201.
10. Houston DM, Elliot DA. (2008) Nutritional management of feline lower urinary tract disorders, In *Encyclopedia of Feline Clinical Nutrition*, Pibot et al. eds.: 285-321.
11. Houston DM, Weese HE, Evason MD, Biourge V, van Hoek I. (2011) A diet with a struvite relative supersaturation less than 1 is effective in dissolving struvite stones *in vivo*. *Br J Nutr* 106: 90-92.
12. Lekcharoensuk C, Osborne CA, Lulich JP, Pusoonthornthum R, Kirk CA, Ulrich LK, Koehler LA, Carpenter KA, Swanson LL. (2001) Association between dietary factors and calcium oxalate and magnesium ammonium phosphate urolithiasis in cats. *J Am Vet Med Assoc* 219: 1228-1237.
13. Markwell PJ, Smith BHE, McCarthy KP. (1999) A non-invasive method for assessing the effect of diet on urinary calcium oxalate and struvite relative supersaturation in the cat. *An Tech* 50: 61-67.
14. Parks JH, Coward M, Coe FL. (1997) Correspondence between stone composition and urine supersaturation in nephrolithiasis. *Kidney Int* 51: 894-900.
15. Prochaska M, Taylor E, Perraro PM, Curhan G. (2018) Relative supersaturation of 24-hour urine and likelihood of kidney stones. *J Urology* 199 (5): 1262-1266.
16. Queau Y, Bijsmans ES, Feugier A, Biourge VC. (2020) Increasing dietary sodium chloride promotes urine dilution and decreases struvite and calcium oxalate relative supersaturation in healthy dogs and cats. *J Anim Physiol Anim Nutr* 104: 1524-1530.
17. Queau Y, Hoek I, Feugier A, Verger L, Soulard Y, Biourge V. (2013) Urinary pH Affects Urinary Calcium Excretion but Not Calcium Oxalate Relative Supersaturation in Health Cats. *ACVIM 2013 (VIN):1*.
18. Robertson WG, Jones S, Heaton MA, Stevenson AE, Markwell PJ. (2002) Predicting the Crystallization Potential of Urine from Cats and Dogs with Respect to Calcium Oxalate and Magnesium Ammonium Phosphate (Struvite). *J Nutr* 132:1637S-1641S.
19. Sagols EM, Cuchet-Subsol C, Billy H, Feugier A, Queau Y. (2015) *In vitro* dissolution of feline struvite stones with 2 diets showing different relative supersaturation values. In: *ESVNU European Society of Veterinary Nephrology and Urology, Proceedings*.
20. Smith BH, Stevenson AE, Markwell PJ. (1998) Urinary relative supersaturations of calcium oxalate and struvite in cats are influenced by diet. *J Nutr* 128: 2763S-2764S.
21. Stevenson AE, Robertson WG, Markwell P. (2003) Risk factor analysis and relative supersaturation as tools for identifying calcium oxalate stone-forming dogs. *J Sm An Prac* 44: 491-496.

## ***Contains Nonbinding Recommendations***

*Draft — Not for Implementation*

22. Stevenson AE, Blackburn JM, Markwell PJ, Robertson WG. (2004) Nutrient intake and urine composition in calcium oxalate stone-forming dogs: comparison with healthy dogs and impact on dietary modification. *Vet Ther* 5: 218-231.
23. Stevenson AE, Wrigglesworth DJ, Markwell PJ. (2000) Urine pH and urinary relative supersaturation in healthy adult cats. *Urolithiasis*, 818-820.
24. Tefft KM, Byron JK, Hostnik ET, Daristotle L, Carmella V, Frantz NZ. (2021) Effect of a struvite dissolution diet in cats with naturally occurring struvite urolithiasis. *J Feline Med Surg* 23 (4): 269-277.
25. Thun MJ and Schober S. (1991) Urolithiasis in Tennessee: An occupational window into a regional problem. *Am J Public Health* 81: 587-59.
26. Torres-Henderson C, Bunkers J, Contreras ET, Cross E, Lappin MR. (2017) Use of Purina Pro Plan Veterinary Diet UR Urinary St/Ox to dissolve struvite cysoliths. *Top Companion Anim Med* 32 (2): 49-54.
27. Tournier C, Biourge VC. (2008) Struvite relative supersaturation and dissolution kinetic. Investigation using an *in vitro* method to reproduce cat bladder environment. *Voorjaarsdagen* 253.
28. Tournier C, Malandain E, Abouhafs S, Aladenise S, Venet C, Ecochard C, Sergheraert R, Biourge V. (2008) Struvite relative supersaturation: a good predictor of struvite stones dissolution *in vitro*. In: Research Abstract Program of the 26th Annual ACVIM Forum. *J Vet Int Med* 22(3): 687-824.
29. Van Hoek I, Malandain E, Tournier C. (2009) RSS is a better predictor for struvite dissolution than urine pH. *Veterinary Focus* 19: 47-48.
30. "2022 Minnesota Urolith Center Global Data." *generated by* Minnesota Urolith Center, March 2023. <https://z.umn.edu/2022GlobalUrolith> (accessed October 31, 2023).

**VII. Appendix 1 – Recommended Combined Safety and Utility Study Flowchart**



+ PE = Physical Exam

\*CBC/Chem/UA = Complete Blood Count, Serum Chemistry, and Urinalysis

† Not needed if baseline collection performed

*Contains Nonbinding Recommendations*  
*Draft — Not for Implementation*

**VIII. Appendix 2 – Glossary of Terms and Acronyms**

**AAFCO** – Association of American Feed Control Officials

**Activity Product** – A numerical description for the concentration of a solute remaining free in solution that is free to interact with other solutes and form crystals

**CaOx** – Type of urolith composed of calcium and oxalate

**CI** – Confidence Interval

**CVM** – Center for Veterinary Medicine

**FDA** – Food and Drug Administration

**Formation Product** – The concentration at which solute will spontaneously precipitate out of solution to form crystals. Defines the limit between the metastable and labile supersaturation zones

***In-Vitro*** – Describes an experiment or test that takes place in a test tube, culture dish, or elsewhere outside a living organism

***In-Vivo*** – Describes an experiment or test that takes place inside a living organism.

**Labeling** – As defined in section 201(m) of the FD&C Act, “means all labels and other written, printed, or graphic matter (1) upon any article or any of its containers or wrappers, or (2) accompanying such article”

**Labile Supersaturation** – Describes the urinary zone in which crystals will spontaneously form and grow

**Metabolic acidosis** – The condition in which too much acid accumulates in the body, causing serious illness

**Metastable Supersaturation** – Describes the urinary zone in which crystals will not spontaneously form but may form in the presence of a nucleus (e.g., an existing urolith or other particulate matter). Existing crystals and uroliths will not dissolve and may grow

**RSS** – Relative Supersaturation

**Solute** – A substance dissolved in solution

**Struvite** – Type of urolith composed of magnesium, ammonia, and phosphate

**Thermodynamic Solubility Product** – The concentration of a solute at which it completely dissolves into solution. Defines the limit between the undersaturation and metastable supersaturation zones

***Contains Nonbinding Recommendations***  
*Draft — Not for Implementation*

**Urolith** – A bladder stone

**Urolithiasis** – The disease condition of having bladder stones

**Undersaturation** – Describes the urinary zone in which crystals will not spontaneously form, and if present, will dissolve

**UTH** – Urinary tract health

**UTH Cat Food** – Adult maintenance cat food with UTH claims