




VO:
For children struggling
every day

1 00:00



VO:
with ADHD, help turn
the tables

2 00:02



VO:
with Qelbree—a
nonstimulant
prescription medication
for children 6-17
with ADHD.

3 00:04



VO:
For challenging

4 00:11

GRAPHIC:
The first nonstimulant
approved for ADHD in
over a decade

SUPER:
For children 6 to 17 years
of age with attention-
deficit/hyperactivity
disorder (ADHD).

PERSISTENT GRAPHIC:
A nonstimulant ADHD
treatment

SUPER:
Individual results may
vary. Pediatric results
were measured after 6 or
8 weeks of therapy.

SUPER:
Individual results may vary. Pediatric results were measured after 6 or 8 weeks of therapy.



VO:
ADHD symptoms,

5

00:11

SUPER:
Individual results may vary. Pediatric results were measured after 6 or 8 weeks of therapy.




VO:
once-daily Qelbree helps provide significant relief.

6

00:13

SUPER:
For more information visit Qelbree.com, call 1-866-398-0833, or talk to your doctor and ask about the Qelbree patient brochure.



VO:
So, instead of their world revolving around ADHD,

7

00:15

SUPER:
For more information visit Qelbree.com, call 1-866-398-0833, or talk to your doctor and ask about the Qelbree patient brochure.



VO:
Qelbree helps make ADHD symptoms

8

00:18

GRAPHIC:
manageable

SUPER:
For more information visit Qelbree.com, call 1-866-398-0833, or talk to your doctor and ask about the Qelbree patient brochure.



VO:
manageable.

9 00:20


SUPER:
It is not known if Qelbree is safe and effective for children less than 6 years of age.



VO:
It's not known if Qelbree is safe and effective for children under 6.

10 00:22

SUPER:
Pay attention to changes in your child's mood, behavior, thoughts, or feelings within the first few months of treatment or when the dose is changed, as medications like Qelbree may increase the risk of suicidal thoughts or behaviors in some children.



VO:
Pay attention to changes in your child's mood, behavior, thoughts, or feelings

11 00:26

SUPER:
Pay attention to changes in your child's mood, behavior, thoughts, or feelings within the first few months of treatment or when the dose is changed, as medications like Qelbree may increase the risk of suicidal thoughts or behaviors in some children.




VO:
within the first few months of treatment or when the dose is changed,

12 00:29

SUPER:
Pay attention to changes in your child's mood, behavior, thoughts, or feelings within the first few months of treatment or when the dose is changed, as medications like Qelbree may increase the risk of suicidal thoughts or behaviors in some children.

Patient portrayals



A nonstimulant ADHD treatment


Pay attention to changes in your child's mood, behavior, thoughts, or feelings within the first few months of treatment or when the dose is changed, as medications like Qelbree may increase the risk of suicidal thoughts or behaviors in some children.

Qelbree
viloxazine

13 00:33

SUPER:
Tell your doctor about all your child's medications.

Patient portrayals



A nonstimulant ADHD treatment

Tell your doctor about all your child's medications.

Qelbree
viloxazine

14 00:38

SUPER:
Do not take MAOIs with or within 14 days of starting Qelbree.

Patient portrayals



A nonstimulant ADHD treatment

Do not take MAOIs with or within 14 days of starting Qelbree.

Qelbree
viloxazine

15 00:40

SUPER:
Qelbree can increase heart rate and blood pressure and may cause manic episodes in patients with bipolar disorder.

Patient portrayals



A nonstimulant ADHD treatment

Qelbree can increase heart rate and blood pressure and may cause manic episodes in patients with bipolar disorder.

Qelbree
viloxazine

16 00:44

VO:
as medications like Qelbree may increase the risk of suicidal thoughts or behaviors in some children.

VO:
Tell your doctor about all your child's medications.

VO:
Do not take MAOIs with or within 14 days of starting Qelbree.

VO:
Qelbree can increase heart rate and blood pressure and

SUPER:
Qelbree can increase heart rate and blood pressure and may cause manic episodes in patients with bipolar disorder.



VO:
may cause manic episodes in patients with bipolar disorder.

17 00:47

SUPER:
Qelbree may cause sleepiness or fatigue, so use caution before driving or operating heavy machinery.



VO:
Qelbree may cause sleepiness or fatigue, so use caution before driving or operating heavy machinery.

18 00:50



VO:
Visit Qelbree.com and ask their doctor if nonstimulant Qelbree

19 00:54

GRAPHIC:
[Qelbree Logo]
Helps make ADHD symptoms manageable
Commercially insured patients may pay as little as \$20
Terms and conditions apply.
QBE.2021-0148 V4



VO:
can help make your child's ADHD symptoms manageable.

20 00:58