

Heated tobacco product use, its correlates, and reasons for use among Mexican smokers

Lizeth Cruz-Jimenez, Inti Barrientos-Gutiérrez, Luis Zavala-Arciniega, Edna Arillo-Santillán, Katia Gallegos-Carrillo, Rosibel Rodríguez-Bolaños, Shannon Gravely, James F. Thrasher



PII: S0376-8716(22)00020-5

DOI: <https://doi.org/10.1016/j.drugalcdep.2022.109283>

Reference: DAD109283

To appear in: *Drug and Alcohol Dependence*

Received date: 18 June 2021

Revised date: 19 November 2021

Accepted date: 9 December 2021

Please cite this article as: Lizeth Cruz-Jimenez, Inti Barrientos-Gutiérrez, Luis Zavala-Arciniega, Edna Arillo-Santillán, Katia Gallegos-Carrillo, Rosibel Rodríguez-Bolaños, Shannon Gravely and James F. Thrasher, Heated tobacco product use, its correlates, and reasons for use among Mexican smokers, *Drug and Alcohol Dependence*, (2021) doi:<https://doi.org/10.1016/j.drugalcdep.2022.109283>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Heated tobacco product use, its correlates, and reasons for use among Mexican smokers

Lizeth Cruz-Jimenez ^a, Inti Barrientos-Gutiérrez ^a, Luis Zavala-Arciniega ^b, Edna Arillo-Santillán ^c, Katia Gallegos-Carrillo ^{a, d}, Rosibel Rodríguez-Bolaños ^c, Shannon Gravely ^e, James F. Thrasher ^{c, f, *}

^a Evaluation and Survey Research Center, National Institute of Public Health, Cuernavaca, Morelos, Mexico

^b University of Michigan, Ann Arbor, Michigan, USA

^c Tobacco Research Department, National Institute of Public Health, Cuernavaca, Morelos, Mexico

^d Epidemiology and Health Services Research Unit, Mexican Institute of Social Security, Cuernavaca, Morelos, Mexico

^e Department of Psychology, University of Waterloo, Waterloo, ON N2L 3G1, Canada

^f Department of Health Promotion, Education & Behavior, Arnold School of Public Health, University of South Carolina, Columbia, USA

* Correspondence: THRASHER@mailbox.sc.edu; +1.803.777.4862 and Lizeth Cruz-Jiménez; +52 55 5467 1000 ext.4336

Abstract

Background. Little is known about the use of novel heated tobacco products (HTPs) in low- and middle-income countries. We examined among smokers in Mexico the prevalence and correlates of HTP use, as well as reasons for using HTPs.

Methods: We analyzed data from five surveys (November 2019-March 2021) of an open cohort of adult smokers (n=6,500), including an oversample of those who also use e-cigarettes. Mixed-effects multinomial logistic models were used to estimate associations between study variables and current HTP use or prior HTP trial relative to never trying HTPs.

Results: The weighted prevalence of current HTP use was 1.1%. Independent correlates of current HTP use included greater smoking frequency, intention to quit, e-cigarette use, having partners/family-members who use e-cigarettes or HTPs, and exposure to HTP information inside/outside tobacco shops. Having partners/family members who smoke and not knowing about the harm of HTPs relative to cigarettes were associated with lower likelihood of current HTP use. Having tried HTPs was more likely among smokers with partners/family who use e-cigarettes or HTPs and exposure to HTP information outside shops and on newspapers/magazines. Among current users, the top two reasons for using HTPs were greater social acceptability (50.2%) and lower perceived harm (40.0%) relative to cigarettes.

Conclusions: Uptake of HTPs appears relatively low among Mexican smokers, and correlates of use are similar to those for e-cigarette use. Further research is needed to determine if HTPs use promotes or impedes smoking cessation, given current HTP users are also likely to use various nicotine products

Keywords: Heated tobacco products; new tobacco products; smokers; electronic cigarettes

1. Introduction

The tobacco industry increasingly markets a diverse array of nicotine products, including heated tobacco products (HTPs), which it claims will move smokers from cigarettes to these presumably less harmful products (Foundation for a Smoke-Free World, 2019). In 2014 Philip Morris International (PMI) launched its HTP “IQOS”, which instead of burning tobacco, heat a tobacco stick at a lower temperature (Philip Morris International, 2020a) to produce an aerosol that contains nicotine (Bitzer et al., 2020; Jackler et al., 2020). Their introduction to the market was followed by aggressive global expansion of IQOS marketing across Asia, Europe, Middle East, North America, and South America (Jackler et al., 2020; Philip Morris International, 2020b). In Japan, a high-income country (HIC) where IQOS was first introduced, it rapidly gained market share (Adamson et al., 2020; Liu et al., 2018; Sutanto et al., 2020; Tabuchi et al., 2019; Wu et al., 2019). As the industry has expanded HTP sales and marketing into low- and middle-income countries (LMICs) (Ochoa, 2020), no research of which we are aware has evaluated the profiles of LMIC

consumers who have used HTPs. Mexico was one of the first LMICs in Latin America where IQOS was introduced, beginning at the end of 2019.

PMI claims that IQOS, the market leader for HTPs, aims “to replace cigarettes with the smoke-free products” (Philip Morris International, 2020c), but the data suggest this aim is not met. Recent studies in Japan and Korea (Hori et al., 2020; Jun et al., 2021), have found that HTPs are used more frequently by multiple tobacco product users (i.e. poly-tobacco users) than by exclusive smokers (Kang et al., 2020; Kuwabara et al., 2020; Sugiyama and Tabuchi, 2020; Sutanto et al., 2020). Research in the HIC countries of Canada, England, USA and Australia found that 89.8% of current HTP users were also concurrent smoker and vapers (Miller et al., 2020). The low level of exclusive HTP use suggests that complete switching to HTPs is unusual (Hwang et al., 2019; Jackler et al., 2020; Kang et al., 2020; Kim and Cho, 2020; Ratajczak et al., 2020). When smoking cessation attempts have been evaluated, smokers who use HTPs were no more likely than exclusive smokers to attempt to quit smoking (Hwang et al., 2019; Kang et al., 2020; Kim and Cho, 2020).

The introduction of HTPs appears to have increased the likelihood of poly-tobacco use, such that compared to exclusive smokers, dual and triple users of HTPs, e-cigarettes, or combustible cigarettes being more likely to have positive perceptions about HTPs (Fung et al., 2020; Kim and Cho, 2020; Sutanto et al., 2020).

Studies of HTP use among smokers are limited to HICs (Jankowski et al., 2019; Ratajczak et al., 2020). General population studies, most of them conducted in Asia, find that HTPs use is higher among males, young adults and those from higher socioeconomic status groups (Hwang et al., 2019; Kim and Cho, 2020; Marynak et al., 2018; Nyman et al., 2018; Sutanto et al., 2020). Furthermore, those who smoke more frequently appear more likely to use HTPs (Hwang et al., 2019). Also, awareness of tobacco company promotions for IQOS was positively associated with current HTP use in Japan (Tabuchi et al., 2018). It should be noted that while Japan bans e-cigarettes, HTPs can be marketed and sold, and its tobacco control regulations are relatively weak (Tanigaki and Poudyal, 2019), which may help explain the rapid growth in the HTPs market there. Korea has stronger tobacco control regulations than Japan, although HTPs and e-cigarettes can be marketed and sold (Jun et al., 2021).

1.1 Study context

Since 2008, Mexican legislation has prohibited the sale, distribution, and promotion of any product that looks like or mimics a cigarette, which has been interpreted to include e-cigarettes and HTPs (Cámara de Diputados del H. Congreso de la Unión, 2012). Nevertheless, since 2018 PMI has promoted IQOS through social media campaigns to anticipate and accompany the launch of product availability in retail stores at the end of 2019, when it became the first and only brand on the market. In early 2020, the importation of HTPs was banned by presidential decree (Diario Oficial de la Federación, 2020). Despite these regulations, HTP sales have continued as legal appeals have made their way through the court system. HTP advertising and promotions include implicit and explicit claims that IQOS are less risky than cigarettes (Philip Morris International, 2021), similar to those in other countries (i.e., “smoke free,” “doesn’t affect the people around you,” “reduces health risks”). These claims may increase HTP appeal and minimize potential health concerns (Gravelly et al., 2020), including among Mexican smokers, who comprise about 17.5% of the 12- to 65-year old population (Reynales-Shigematsu et al., 2017). Therefore, the present study examines the prevalence, correlates and patterns of HTP use among Mexican smokers, including their reasons for using HTPs.

2. Methods

2.1 Data source

Data come from five surveys of an open cohort of Mexican smokers and e-cigarette users recruited through a non-probability sample of participants from online consumer panel and surveyed between November 2019 to March 2021. Participants had to be adults (≥ 18 years old) and have smoked or used e-cigarettes in the prior 30 days. At each survey, 1500 participants were recruited, with quotas used for last month e-cigarette use ($n > 500$) and educational attainment (approximately 1/3=high school or less; 1/3=technical/trade school or community college/or some college; and 1/3=college degree or higher). We oversampled e-cigarette users to evaluate dual users (33.9%). We excluded from the sample those who did not give information about household income ($n=398$), those who had quit smoking at the time of the survey ($n=436$), and exclusive e-cigarette users ($n=172$), who were excluded

due to their small sample size across HTP use outcomes. Participants were followed to the extent possible, with the sample replenished with new participants to maintain the target sample size at each survey. The final analytic sample included 6,500 observations (Nov. 2019 n=1321, Mar. 2020 n=1282, Jul. 2020 n=1272, Nov. 2020 n=1309 and Mar 2021 n=1316), 3,108 unique participants who at the time of the survey were current cigarette smokers.

Surveys were administered in Spanish using standard questions on tobacco product use (International Agency for Research on Cancer, 2008) and questions on novel tobacco products from the International Tobacco Control (ITC) survey (Thompson et al., 2019). The survey took on average between 20-25 minutes to complete, and the panel provider gave standard compensation for participation (e.g., points-based or monetary rewards, chances to win prizes). All study procedures were approved by the Institutional Review Board and Ethics Committee of the National Institute of Public Health of Mexico (CI 1572).

2.2 Measures

2.2.1 HTP use variables

The survey section on HTPs began with a brief product description and image of IQOS, the only HTP available in Mexico. Product awareness was assessed (i.e., “Have you heard of heated tobacco product (outside of these surveys)?”) with responses dichotomized (yes vs. no or don’t know). Those who reported awareness were asked whether they had ever tried an HTP (yes vs. no or don’t know). Participants who reported HTP trial were queried about the frequency of current use (daily; less than daily, but at least once a week; less than weekly, but at least once a month; less than once a month, but occasionally; not at all). Responses to these questions were used to derive categories of use: never tried (i.e., unaware of HTPs or never tried HTPs); HTP trial (i.e., tried HTPs, but no use in the last month); and current HTP use (i.e., in the last month).

Current HTP users were asked about heatsticks/heets last used, showing images of each variety: Sienna Selection (intense tobacco), Amber Selection (toasted tobacco and nuts), Yellow Selection (smooth tobacco with citrus), Blue Selection (smooth menthol),

Turquoise Selection (deep menthol), Purple Wave (fruit-flavored menthol), other type, and “don't know”. HTP users also reported how they obtained the last IQOS device they used (i.e., bought; gifted; borrowed; free sample). Participants who reported buying it were asked where they bought it (online; vape shop or tobacconist; department store or supermarket; pharmacist; convenience store; temporary or mobile sales location, and gas station), the latter four categories were combined as “other” due to small sample sizes.

2.2.2 Smoking- and e-cigarette related variables

All participants reported smoking frequency and were categorized using cutpoints that generally reflect tertiles of consumption intensity in Mexico (non-daily; daily ≤ 5 cigarettes and daily > 5 cigarettes) (Pan American Health Organization and National Institute of Public Health-Mexico, 2017). Participants also reported recent smoking cessation attempts (i.e., in the last four months; yes vs. no) and intentions to quit smoking (i.e., in the next six months vs. not). Self-reported e-cigarette use frequency in the prior month was used to derive categories of exclusive cigarette smokers, sporadic dual user (i.e., e-cigarette use twice a week or less); and frequent dual user (i.e., e-cigarette use three times a week or more), with cutpoints based on the use median.

2.2.3 Descriptive norms of partner, family and friends

With two separate questions, we asked whether participants had a partner or spouse who smokes and whether a household family member smokes. Responses were combined to indicate smoking by partner or family (i.e., yes vs. no). For two parallel questions on e-cigarette use among partners/family, we used the same coding (yes vs. no). We also asked about HTP use among household family members (yes vs. no). Separate questions asked about current smoking, e-cigarette use, and HTP use among participants' five closest friends with whom they regularly spend time, with responses for each product dichotomized to reflect use by any of these friends (yes vs. no). For all these questions, those who responded “I don't know” or who indicated they were unaware of HTPs were classified as “no”.

2.2.4 Relative risk perceptions and exposure to information about HTPs

Participants who reported awareness of HTPs were asked their perception of HTPs' harmfulness relative to cigarettes ("Compared to smoking cigarettes, how harmful do you think using a heated tobacco product is?"), with responses categorized to reflect lower perceived harm than cigarettes (i.e. much less harmful than smoking cigarettes, somewhat less harmful than smoking cigarettes), equally or more harmful than cigarettes (i.e. equally harmful to smoking cigarettes, somewhat more harmful than smoking cigarettes, much more harmful than smoking cigarettes) and I don't know/lack of awareness about HTPs. Additionally, we assessed potential marketing exposures about HTPs in the last 30 days via: the internet (yes vs. no); inside shops/stores that sell tobacco products (yes vs. no); outside shops/stores that sell tobacco products (yes vs. no); and in newspapers or magazines (yes vs. no). Those who reported being unaware of HTPs were classified as having "no" exposure.

2.2.6 Reasons for HTP and e-cigarette use

Participants who reported current use of HTPs and/or e-cigarettes answered product-specific but parallel questions about their reasons for using the product, with the option to check all that apply: is less harmful to people around me; is more acceptable than smoking ordinary cigarettes to people around me; helps me cut down on the number of ordinary cigarettes I smoke; I can use them in places where I can't smoke; and might help me stay quit from smoking ordinary cigarettes.

2.2.7 Sociodemographic variables

Sociodemographic measures included sex (male and female), age (18–29, 30–39, 40–49, and +50 years), educational attainment (middle school and less; high school, technical or some college; and university and more) and monthly household income in Mexican pesos (1 USD = 21 MXN: less than 8,000 MXN; 8,001 to 15,000 MXN; and 15,001 to >20,000MXN) responses of "refused" and "don't know" were coded as missing.

2.3 Analysis

We evaluated descriptive statistics using unweighted and weighted data, with inverse probability of selection weights based on the sex, age, and educational attainment profiles for exclusive smokers and, separately, for dual users according to nationally-representative data from 2018 (Shamah-Levy et al., 2020). Prevalences for HTP-related variables (awareness, ever tried and current use) were estimated as weighted proportions; and we compared prevalences by survey using weighted chi-square tests. Additionally, we assessed endorsement of different reasons for using HTPs and e-cigarettes among current users of each device, comparing them using weighted chi-square tests. Using mixed effects multinomial logistic regression models to account for repeated measures among those who participated in multiple surveys, we estimated crude and adjusted relative risk ratios (RRRs & ARRRs, respectively) to estimate the likelihood of HTP use (current HTP use, prior HTP use, never tried HTPs=reference) by sociodemographic, smoking-related variables, descriptive social norms, perceived relative harmlessness of HTPs, and exposure to HTP information. As a sensitivity analysis, we re-estimated these models after excluding participants who were unaware of HTPs. Similarly, we re-estimated the models for HTP use without weights. We do not report on the analytic sample (Supplement 1) or unweighted results (available upon request), since results were consistent with those from the full analytic sample using weighted data, and would not have changed our primary interpretations. Because of the purposive nature of our sample, especially the oversample of e-cigarette users, we believe the weighted estimates are of wider interest since they are more suitable for generalizing to the population of smokers in Mexico. Prevalence estimates, confidence intervals, and model inference were based on modified sandwich standard errors that adjusted for any form of within-participant correlation. All analyses were conducted using Stata v.14 (Stata Corp, TX, USA).

3. Results

Table 1 presents weighted characteristics of participants as well as the unweighted sample size for each subgroup of interest (N=6,500). About one-third (31.6%) reported being aware of HTPs, 5% reporting having tried them but not currently and 1.1% using them currently. The weighted prevalence of these variables did not significantly differ across

survey waves, except for HTP trial ($p=0.0001$), this difference appeared driven primarily by the highest estimate from the July 2020 survey (15.6%).

3.1 Factors associated with the use of HTPs among adult smokers

In adjusted models for HTP use (Table 2), likelihood of HTP trial (vs. never tried HTPs) was higher among those with partners/family who use e-cigarettes or HTPs ($ARRR=3.89$, 95% CI 1.79-8.44 and $ARRR=5.32$, 95% CI 1.88-15.08, respectively), as well as those exposed to HTPs information outside shops/stores where tobacco is sold ($ARRR=2.26$, 95% CI 1.10-4.63) or through newspapers or magazines ($ARRR=6.12$, 95% CI 2.92-15.00). Not having an opinion about the harm of HTPs relative to cigarettes, were associated with less likely to current HTPs use ($ARRR_{I\text{ don't know/ unaware vs. equally or more harmful}}=0.16$, 95% CI 0.05-0.54). Other significant correlates of ever tried included: educational attainment ($ARRR_{\text{high school / technical/ some college vs. university}}=2.78$, 95% CI 1.34-5.78) and household income (i.e., 15,001 or more Mexican pesos a month) ($ARRR=2.94$, 95% CI 1.06-8.11).

Likelihood of current HTP use (vs. never tried HTPs) was higher among those who were light daily smokers ($ARRR_{\text{daily } \leq 5 \text{ cigarettes vs. non-daily}}=6.16$, 95% CI 2.52-15.05), intend to quit ($ARRR=2.57$, 95% CI 1.14-4.92), and use e-cigarettes, whether sporadically ($ARRR_{\text{sporadic e-cigarette use vs. no use}}=6.35$, 95% CI 2.46-16.37) or more frequently ($ARRR_{\text{frequent e-cigarette use vs. no use}}=11.20$, 95% CI 3.85-32.90). Current HTP use was also higher among participants: with partners/family who use e-cigarettes ($ARRR=10.04$, 95% CI 3.03-33.22) or HTPs ($ARRR=1.38$, 95% CI 1.46-19.79); and those who reported having seen information about HTPs inside or outside shops where tobacco is sold ($ARRR=2.43$, 95% CI 1.06-5.61 and $ARRR=3.54$, 95% CI 1.67-7.50, respectively). Also, having partners/family members who smoke ($ARRR=0.34$, 95% CI 0.13-0.92) and not having an opinion about the harm of HTPs relative to cigarettes ($ARRR_{I\text{ don't know/ unaware vs. equally or more harmful}}=0.03$, 95% CI 0.01-0.08) were associated with lower likelihood of current HTPs use.

3.2 Use patterns and preferences among current HTP users

Among respondents who reported current HTP use (unweighted $n=665$), 13.5% reported daily use (Table 3). The most popular heatsticks/heets variety they used most

recently were “Blue Selection” (31.8%), followed by “Amber Selection” (28.7%) and “Sienna Selection” (15.1%). Of the approximately 44.2% who bought their HTP device, 68.7% did so online and 18.5% from a vape shop or tobacconist.

3.3 Reasons for use HTPs and e-cigarette

Figure 1 shows the weighted percentages for each of the reasons for using HTPs and e-cigarettes, queried only among current users of HTPs (n=665) and e-cigarettes users (n=2,273). The most prevalent reason for using HTPs was their social acceptability relative to cigarettes (50.2%), followed by because HTPs are less harmful than cigarettes to people around them (40.0%) and because it helps to cut down on the number of combustible cigarettes they smoke (28.6%). Among e-cigarette users, the most frequently endorsed reason for use was lower harm to others (47.7%), followed by their social acceptability (39.7%) and because it helps stay quit from smoking cigarettes (28.1%). Use to cut down on cigarettes was the only reason that was statistically different when we compared HTPs users and e-cigarette users ($p<0.01$, 28.6% vs. 14.9%, respectively).

4. Discussion

This study among Mexican smokers found that the prevalence of current HTP use was approximately 1.1% and remained relatively stable over the year and a half period after their introduction into the market. This is similar to what has been found in European countries (Germany, Greece, Hungary, Poland, Romania and Spain) where 0.8% of adult smokers use HTPs (Lotrean et al., 2020), as well as in Canada, England, USA and Australia where this prevalence was 0.9% (Miller et al., 2020). Our results are also relatively consistent with a recent study across 28 European Union countries, including the United Kingdom, which found 1.3% of the general population used HTPs, although current and former smokers were more likely to have ever or currently used HTPs (Lavery et al., 2021). Longitudinal research is needed to better understand who tries and goes on to use HTPs consistently, as, to our knowledge, all studies in this area are cross-sectional and do not ask about when consumers first tried HTPs.

In line with prior studies (Brose et al., 2018; Hwang et al., 2019; Kang et al., 2020; Kim and Cho, 2020), we found that most smokers who currently use HTPs also use e-cigarettes, making them poly-users of nicotine products. However, we did not find this association with HTP trial, suggesting that e-cigarette use promotes continued use HTPs. Longitudinal studies are needed to better evaluate the trajectories of HTP use, including assessment of quitting combustible cigarettes and the potential for impeding cessation, perhaps by providing a more socially acceptable and less detectable means of nicotine delivery than cigarettes for situations where one cannot smoke, as suggested in other studies (Hair et al., 2018; Tompkins et al., 2020).

Our focus on descriptive norms, or perceptions of actual behavior within a social group (Lapinski and Rimal, 2005), follows from diffusion of innovation theory's emphasis on social networks for explaining why innovations like HTPs are adopted (Rogers, 2003). We found strong associations between use of e-cigarettes and HTPs among partners/family and both trial and current use of HTPs. Furthermore, a substantial proportion of current HTP users obtained their devices as a gift from a relative or friend (34.0%). This follows a prior qualitative study among HTP users and ex-users who reported that family and friends who used HTPs promoted its use as an alternative to smoking and suggested they try it (Tompkins et al., 2020). Indeed, F&M marketing strategies in Mexico include a referral program where current users can get coupons for IQOS products if they promote IQOS use among their family and friends (Philip Morris International, 2020d). Notably, friend use of e-cigs or HTPs was strongly correlated with trial and use, but not in adjusted models, suggesting that familial influence matters more in the Mexican context given that family use was significant in both adjusted and unadjusted models.

Our findings contrast with prior studies that reported a high prevalence of believing HTPs were less harmful than smoking among current users of HTPs (Gravelly et al., 2020; Lavery et al., 2021; Majek et al., 2021). A qualitative study of adult HTP users in the UK perceptions of lower harm were also prominent, although, as our results, there was a great uncertainty about risks from use (East et al., 2021). This could be due to the misperception promoted by the industry and the lack of scientific agreement on the risk of these products. Monitoring HTP marketing with implied and direct reduced risk claims, as well as

evaluations of smokers' responses to this marketing, will be important in evaluating the uptake and consequences of HTP use going forward.

Consistent with the findings from a study of Japanese smokers, we found that the most prevalent exposure to HTP marketing among HTPs users was inside and outside stores (Craig et al., 2020). This engagement may happen after progression from trial to regular IQOS use. This is not entirely surprising, since smokers may be more likely to go to places where marketing activities are more intensive, such as tobacco shops. That exposure to HTP information on the Internet was significant only in bivariate analyses – and only close to statistical significance in multivariate analyses – may be because online ad exposures are more prevalent among young adults (Jackler et al., 2020). Future research should focus on whether the effects of such exposures vary by age group. In Mexico, as in many other countries, the absence of a comprehensive tobacco product advertising ban and challenges around regulating internet content make it difficult to monitor and reduce marketing activities (Jankowski et al., 2019).

In the Metropolitan area of Mexico City, PMI promotes the IQOS through a fifteen day device loan program, with a discount voucher of 400 Mexican pesos (approximately \$19 USD) if the person ultimately purchases the device for around \$36 USD with the discount (Philip Morris International, 2020d). We are not aware of any free IQOS giveaway programs in Mexico, although a small percentage of users in our study reported getting their device as a free sample (12.3%), perhaps because they interpreted the loan program as a free sample. In a global market, making the device more accessible through discount packages gains new consumers who repeatedly buy products, like the tobacco sticks that are needed for HTPs (Jackler et al., 2020).

Consistent with the findings of a study among Japanese smokers (Sutanto et al., 2019), a large percentage of current users of HTPs prefer menthol flavor (41.5%). This is comparable to the prevalence of preference for cigarette varieties with flavor capsules in the filter in Mexico (43%) (Zavala-Arciniega et al., 2020), where mint/menthol flavor is common (Ogunnaike et al., 2020). Tobacco companies have long manipulated menthol content to promote smoking initiation, since menthol can mask the harshness of smoke among those who first experiment with the cigarettes (Kreslake et al., 2008). Indeed, this has driven some concerns about whether the appeal of HTPs for youth has been properly

considered in evaluating their potential public health impact (Lempert and Glantz, 2020). Recent research among Guatemalan adolescents indicates that very few use HTPs (Gottschlich et al., 2020) and most perceive them as less appealing and more harmful than cigarettes (Monzón et al., 2021). Nevertheless, the great popularity of menthol flavor among HTP users, the inclusion of additional flavors like berry, and their potential uptake among youth should be further evaluated to clarify the role of HTPs in tobacco product initiation, relapse, or maintenance of nicotine dependence. Indeed, given the appeal of flavor capsules for youth, (Abad-Vivero et al., 2016; Barrientos-Gutierrez et al., 2021) efforts should be made to impede the introduction of HTP sticks that include capsules, as these have been introduced in other countries (Cho and Thrasher 2015).

Our findings about the main reasons for using HTPs are consistent with previous qualitative and quantitative studies (Adamson et al., 2020; Hair et al., 2018; Queloiz and Etter, 2019; Tompkins et al., 2020). The top reasons for using HTPs were social (can use around others and not harm them), similar to other studies (Adamson et al., 2020; Hair et al., 2018) and for the top reasons smokers in our sample used e-cigarettes. As suggested in other studies, smokers may use HTPs to experiencing relatively less stigma and negative judgments than when they smoke in public or around non-smokers (Hair et al., 2018; Tompkins et al., 2020). The use of HTPs because they are less harmful than cigarettes to people around them may be linked to reduced ash and odor from HTPs (Hair et al., 2018), which is a key message in IQOS marketing materials and which, by extension, gives the appearance of lower toxicity than combustible cigarettes (Queloiz and Etter, 2019). Given the relatively low percentage of smokers who use HTPs to cut down on cigarettes or help them quit, HTPs may mostly be used in social situations where this social use is a substitute for smoking (which may be preferred). Indeed, having recently tried to quit smoking was not associated with either trial or current use of HTPs, although quit intention was positively associated with current use. This finding contrasts with some studies where intention to quit smoking was relatively uncommon among HTP users (Park et al., 2021; Ryu et al., 2020). Given the relatively short period of time that HTPs have been on the Mexican market and the relatively low prevalence of use in Mexico compared to Asian countries, early adopters of novel HTP devices may be more to be interested in trying to quit compared to later adopters.

Our study has some limitations, including the use of a convenience sample from an online panel used for marketing research, where we over-sampled e-cigarette users and under-sampled smokers from lower socioeconomic status groups. Although our analyses integrated weighting to reflect sociodemographic profiles of exclusive smokers and dual users in the general population, our results may still be biased. Also, our analysis of HTP use includes those who were unaware of HTPs, so that the models would consider the characteristics of this subpopulation of smokers. These participants, however, were not queried about perceived harmfulness of HTPs, use of HTPs among family and friends, or exposure to HTP marketing, which is why we grouped them with those who reported not knowing the relative harms, no use of HTPs among network members, and no HTP marketing exposures. However, results from our sensitivity analyses that excluded participants who were unaware of HTPs indicated that the correlates of HTP use were consistent with those we found when analyzing the entire sample. Finally, our study was cross sectional, and so the temporal ordering of relationships is not clear. Longitudinal data are needed to better examine the incidence and correlates of tobacco product transitions related to HTP use.

5. Conclusion

In our sample of Mexican smokers, approximately 1.1% currently use HTPs, similar to findings in a variety of countries. Smokers who use e-cigarettes are particularly likely to use HTPs, leading to poly-product use as has also been found in other countries. Use of HTPs and e-cigarettes among social network members, particularly partners and family, appears to promote use, which industry explicitly encourages through its marketing. While HTP users are more likely to intend to quit smoking than those who have never tried HTPs, other data suggest that HTPs may be mostly used to deliver nicotine in settings where smoking is socially unacceptable, as is also found for e-cigarette use.

References

- Abad-Vivero, E.N., Thrasher, J.F., Arillo-Santillán, E., Pérez-Hernández, R., Barrientos-Gutiérrez, I., Kollath-Cattano, C., Mejía, R., Sargent, J.D., 2016. Recall, appeal and willingness to try cigarettes with flavour capsules: assessing the impact of a tobacco product innovation among early adolescents. *Tob. Control* 25, e113 LP-e119. <https://doi.org/10.1136/tobaccocontrol-2015-052805>
- Adamson, J., Kanitscheider, C., Prasad, K., Camacho, O.M., Beyerlein, F., Bhagavan, Y.K., Proctor, C., Murphy, J., 2020. Results from a 2018 cross-sectional survey in Tokyo, Osaka and Sendai to assess tobacco and nicotine product usage after the introduction of heated tobacco products (HTPs) in Japan. *Harm Reduct. J.* 17, 32. <https://doi.org/10.1186/s12954-020-00374-3>
- Barrientos-Gutiérrez, I., Islam, F., Cho, Y.J., Salloum, R. G., Louviere, J., Arillo-Santillán, E., Reynales-Shigematsu, L.M., Barnoya, J., Saenz de Miera Juárez, B., Hardin, J., Thrasher, J.F., 2021. Assessing cigarette packaging and labelling policy effects on early adolescents: results from a discrete choice experiment. *Tob. Control* 30, 505 LP – 514. <https://doi.org/10.1136/tobaccocontrol-2019-055463>
- Bitzer, Z.T., Goel, R., Trushin, N., Muscat, J., Richie, J.P., 2020. Free Radical Production and Characterization of Heat Not-Burn Cigarettes in Comparison to Conventional and Electronic Cigarettes. *Chem. Res. Toxicol.* <https://doi.org/10.1021/acs.chemrestox.0c00088>
- Brose, L.S., Simonavicius, E., Cheeseman, H., 2018. Awareness and Use of “Heat-not-burn” Tobacco Products in Great Britain. *Tob. Regul. Sci.* 4, 44–50. <https://doi.org/10.18001/trs.4.2.4>
- Cámara de Diputados del H. Congreso de la Unión, 2012. Reglamento de la ley general para el control del tabaco [WWW Document]. URL http://www.diputados.gob.mx/LeyesBiblio/regley/Reg_LGCT.pdf (accessed 5.22.20).
- Cho, Y.J., Thrasher, J.F., 2019. Flavour capsule heat-sticks for heated tobacco products. *Tob. Control* 28, e158–e159. <https://doi.org/10.1136/tobaccocontrol-2018-054472>
- Craig, L. V., Yoshimi, I., Fong, G.T., Meng, G., Yan, M., Mochizuki, Y., Tabuchi, T., Thrasher, J.F., Xu, S.S., Quah, A.C.K., Ouimet, J., Sansone, G., Chung-Hall, J., 2020. Awareness of Marketing of Heated Tobacco Products and Cigarettes and Support for

- Tobacco Marketing Restrictions in Japan: Findings from the 2018 International Tobacco Control (ITC) Japan Survey. *Int. J. Environ. Res. Public Heal.* .
<https://doi.org/10.3390/ijerph17228418>
- Diario Oficial de la Federación, 2020. Decreto por el que se modifica la Tarifa de la Ley de los Impuestos Generales de Importación y de Exportación. [WWW Document]. URL http://www.dof.gob.mx/nota_detalle.php?codigo=5586899&fecha=19/02/2020 (accessed 6.4.20).
- East, K.A., Tompkins, C.N.E., McNeill, A., Hitchman, S.C., 2021. ‘I perceive it to be less harmful, I have no idea if it is or not:’ a qualitative exploration of the harm perceptions of IQOS among adult users. *Harm Reduct. J.* 18, 42. <https://doi.org/10.1186/s12954-021-00490-8>
- Foundation for a Smoke-Free World, 2019. Global trends in nicotine [WWW Document]. URL <https://www.smokefreeworld.org/wp-content/uploads/2019/08/fsfw-report-trends-in-nicotine-1005201811.pdf> (accessed 5.16.20).
- Fung, M.D.T., Diemert, L.M., Zhang, P., O'Connor, S., Schwartz, R., 2020. Awareness and Perceived Risk of Heated Tobacco Products. *Tob. Regul. Sci.* 6, 15–19.
<https://doi.org/10.18001/TRS.v6i1.2>
- Gottschlich, A., Mus, S., Monzón, J.C., Thrasher, J.F., Barnoya, J., 2020. Cross-sectional study on the awareness, susceptibility and use of heated tobacco products among adolescents in Guatemala City, Guatemala. *BMJ Open* 10, e039792.
<https://doi.org/10.1136/bmjopen-2020-039792>
- Gravelly, S., Fong, G.T., Sutanto, E., Loewen, R., Ouimet, J., Xu, S.S., Quah, A.C.K., Thompson, M.E., Boudreau, C., Li, G., Goniewicz, M.L., Yoshimi, I., Mochizuki, Y., Elton-Marshall, T., Thrasher, J.F., Tabuchi, T., 2020. Perceptions of Harmfulness of Heated Tobacco Products Compared to Combustible Cigarettes among Adult Smokers in Japan: Findings from the 2018 ITC Japan Survey. *Int. J. Environ. Res. Public Health.* <https://doi.org/10.3390/ijerph17072394>
- Hair, E.C., Bennett, M., Sheen, E., Cantrell, J., Briggs, J., Fenn, Z., Willett, J.G., Vallone, D., 2018. Examining perceptions about IQOS heated tobacco product: Consumer studies in Japan and Switzerland. *Tob. Control* 27, s70–s73.
<https://doi.org/10.1136/tobaccocontrol-2018-054322>

- Hori, A., Tabuchi, T., Kunugita, N., 2020. Rapid increase in heated tobacco product (HTP) use from 2015 to 2019: from the Japan Society and New Tobacco Internet Survey (JASTIS). *Tob. Control*. <https://doi.org/10.1136/tobaccocontrol-2020-055652>
- Hwang, J.H., Ryu, D.H., Park, S.W., 2019. Heated tobacco products: Cigarette complements, not substitutes. *Drug Alcohol Depend.* 204. <https://doi.org/10.1016/j.drugalcdep.2019.107576>
- International Agency for Research on Cancer, 2008. Methods for evaluating tobacco control policies, in: *Methods for Evaluating Tobacco Control Policies*. Lyon, France, p. 459.
- Jackler, R., Ramamurthi, D., Axelrod, A., Jung, J., Louis-Ferdinand, N., Reidel, J., Yu, A., Jackler, L., Chau, C., 2020. Global Marketing of IQOS The Philip Morris Campaign to Popularize “Heat Not Burn” Tobacco [WWW Document]. URL <http://tobacco.stanford.edu/iqosanalysis> (accessed 1.22.20).
- Jankowski, M., Brożek, G.M., Lawson, J., Stoczek, S., Majek, P., Zejda, J.E., 2019. New ideas, old problems? Heated tobacco products – a systematic review. *Int. J. Occup. Med. Environ. Health* 92, 595–634. <https://doi.org/10.13075/ij.oreh.1896.01433>
- Jun, J., Kim, S.-H., Thrasher, J., Cho, Y.J., Heo, Y.-J., 2021. Heated debates on regulations of heated tobacco products in South Korea: the news valence, source and framing of relative risk/benefit. *Tob. Control*. <https://doi.org/10.1136/tobaccocontrol-2020-056131>
- Kang, S.Y., Lee, S., Cho, H.-J., 2020. Prevalence and predictors of heated tobacco product use and its relationship with attempts to quit cigarette smoking among Korean adolescents. *Tob. Control* tobaccocontrol-2019-055114. <https://doi.org/10.1136/tobaccocontrol-2019-055114>
- Kim, Cho, 2020. Prevalence and correlates of current use of heated tobacco products among a nationally representative sample of Korean adults: Results from a cross-sectional study. *Tob. Induc. Dis.* 18. <https://doi.org/10.18332/tid/125232>
- Kreslake, J.M., Wayne, G.F., Alpert, H.R., Koh, H.K., Connolly, G.N., 2008. Tobacco Industry Control of Menthol in Cigarettes and Targeting of Adolescents and Young

- Adults. *Am. J. Public Health* 98, 1685–1692.
<https://doi.org/10.2105/AJPH.2007.125542>
- Kuwabara, Y., Kinjo, A., Fujii, M., Imamoto, A., Osaki, Y., McNeill, A., Beckley-Hoelscher, N., 2020. Comparing Factors Related to Any Conventional Cigarette Smokers, Exclusive New Alternative Product Users, and Non-Users among Japanese Youth: A Nationwide Survey. *Int. J. Environ. Res. Public Health*.
<https://doi.org/10.3390/ijerph17093128>
- Lapinski, M.K., Rimal, R.N., 2005. An explication of social norms. *Commun. Theory* 127–147.
- Laverty, A.A., Vardavas, C.I., Filippidis, F.T., 2021. Prevalence and reasons for use of Heated Tobacco Products (HTP) in Europe: an analysis of Eurobarometer data in 28 countries. *Lancet Reg. Heal. - Eur.* 8, 100159.
<https://doi.org/https://doi.org/10.1016/j.lanepe.2021.100159>
- Lempert, L.K., Glantz, S., 2020. Analysis of FDA’s ITC OS marketing authorisation and its policy impacts. *Tob. Control tobaccocontrol*-2019-055585.
<https://doi.org/10.1136/tobaccocontrol-2019-055585>
- Liu, X., Lugo, A., Spizzichino, L., Febuchi, T., Gorini, G., Gallus, S., 2018. Heat-Not-Burn Tobacco Products Are Getting Hot in Italy. *J. Epidemiol.* 28, 274–275.
<https://doi.org/10.2183/jea.1E20180040>
- Lotrean, L., Trofor, A., Radu Loghin, C., Eremia, M., Mihaltan, F., Driezen, P., Kyriakos, C.N., Mons, U., Domján, T., Fernández, E., Katsaounou, P.A., Przewoźniak, K., Filippidis, F.T., Gravelly, S., Fong, G.T., Vardavas, C.I., 2020. Awareness and use of heated tobacco products among adult smokers in six European countries: findings from the EUREST-PLUS ITC Europe Surveys. *Eur. J. Public Health* 30, iii78–iii83.
<https://doi.org/10.1093/eurpub/ckz228>
- Majek, P., Jankowski, M., Nowak, B., Macherski, M., Nowak, M., Gil, A., Nakiela, P., Lewicka, B., Lawson, J.A., Zejda, J.E., Brożek, G.M., 2021. The Frequency of Use and Harm Perception of Heated Tobacco Products (HTPs): The 2019 Cross-Sectional Survey among Medical Students from Poland. *Int. J. Environ. Res. Public Heal.* .
<https://doi.org/10.3390/ijerph18073381>
- Marynak, K.L., Wang, T.W., King, B.A., Agaku, I.T., Reimels, E.A., Graffunder, C.M.,

2018. Awareness and Ever Use of “Heat-Not-Burn” Tobacco Products Among U.S. Adults, 2017. *Am. J. Prev. Med.* 55, 551–554.
<https://doi.org/10.1016/J.AMEPRE.2018.04.031>
- Miller, C.R., Sutanto, E., Smith, D.M., Hitchman, S.C., Gravely, S., Yong, H.H., Borland, R., O’Connor, R.J., Cummings, K.M., Fong, G.T., Hyland, A., Quah, A.C.K., Goniewicz, M.L., 2020. Awareness, trial and use of heated tobacco products among adult cigarette smokers and e-cigarette users: findings from the 2018 ITC Four Country Smoking and Vaping Survey. *Tob. Control* tobaccocontrol-2020-055985.
<https://doi.org/10.1136/tobaccocontrol-2020-055985>
- Monzón, J., Islam, F., Mus, S., Thrasher, J.F., Barnoya, J., 2021. Effects of tobacco product type and characteristics on appeal and perceived harm: Results from a discrete choice experiment among Guatemalan adolescents. *Prev. Med. (Baltim)*. 148, 106590.
<https://doi.org/https://doi.org/10.1016/j.ypmed.2021.106590>
- Nyman, A.L., Weaver, S.R., Popova, L., Pechacek, T.J., Huang, J., Ashley, D.L., Eriksen, M.P., 2018. Awareness and use of heated tobacco products among US adults, 2016–2017. *Tob. Control* 27, s55 LP-s61. <https://doi.org/10.1136/tobaccocontrol-2018-054323>
- Ochoa, E., 2020. Desenmascarando a la industria tabacalera en América Latina [WWW Document]. URL <https://saludjusta.mx/wp-content/uploads/Reporte-Regional-Desenmascarando-a-la-IT-en-AL.pdf> (accessed 3.4.21).
- Ogunnaike, A., Barrientos-Gutierrez, I., Arillo-Santillán, E., Gallegos, K., Tharsher, J., 2020. Why smoke flavor capsule cigarettes? preferences and perceptions among adult smokers in Mexico. [WWW Document]. URL https://cdn.ymaws.com/www.srnt.org/resource/resmgr/conferences/2020_%0Dannual_meeting/SRNT20_Rapid_Abstracts_02272.pdf (accessed 6.1.21).
- Pan American Health Organization, National Institute of Public Health-Mexico, 2017. Global Adult Tobacco Survey (GATS) Mexico 2015 [WWW Document]. URL https://www.gob.mx/cms/uploads/attachment/file/239099/GATS_2009.pdf (accessed 3.21.21).
- Park, J., Kim, H.J., Shin, S.H., Park, E., Oh, J.-K., Park, E.Y., Lim, M.K., 2021. Perceptions of heated tobacco products (HTPs) and intention to quit among adult

- tobacco users in Korea. *J. Epidemiol. advpub.* <https://doi.org/10.2188/jea.JE20200213>
- Philip Morris International, 2021. Cambia a una alternativa de tabaco sin humo [WWW Document]. URL <https://www.iqos.com/mx/es/Noticias/cambia-alternativa-tabaco-sin-humo.html> (accessed 2.2.21).
- Philip Morris International, 2020a. Tobacco meets technology [WWW Document]. URL <https://www.pmi.com/smoke-free-products/iqos-our-tobacco-heating-system> (accessed 6.16.20).
- Philip Morris International, 2020b. Building Leading Brands [WWW Document]. URL <https://www.pmi.com/investor-relations/overview/building-leading-brands> (accessed 6.4.20).
- Philip Morris International, 2020c. Delivering a smoke-free future [WWW Document]. URL <https://www.pmi.com/our-transformation/delivering-a-smoke-free-future> (accessed 6.4.20).
- Philip Morris International, 2020d. Programa de Prestamo IQOS [WWW Document]. URL https://www.probar-iqos.com/?utm_source=iqosreferral&utm_medium=topnav&utm_campaign=menu.html (accessed 10.2.20).
- Queloz, S., Etter, J.F., 2019. A nationwide survey of users of tobacco vaporizers, reasons and modes of utilization, perceived advantages and perceived risks. *BMC Public Health* 19. <https://doi.org/10.1186/s12889-019-6957-0>
- Ratajczak, A., Janikowski, P., Strus, P., Feleszko, W., 2020. Heat Not Burn Tobacco Products – A New Global Trend: Impact of Heat-Not-Burn Tobacco Products on Public Health, a Systematic Review. *Int. J. Environ. Res. Public Heal.* . <https://doi.org/10.3390/ijerph17020409>
- Reynales-Shigematsu, L., Zavala-Arciniega, L., Paz-Ballesteros, WC. Gutiérrez-Torres, D., García-Buendía, J., Rodríguez-Andrade, MA. Gutiérrez-Reyes, J., Franco-Núñez, A., Romero-Martínez, M., Mendoza-Alvarado, L., 2017. Encuesta Nacional de Consumo de Drogas, Alcohol y Tabaco 2016-2017: Reporte de Tabaco. [WWW Document]. INPRFM. URL http://inprf.gob.mx/psicosociales/archivos/ena/ENCODAT_DROGAS_2016-2017.pdf (accessed 1.21.02).

- Rogers, E.M., 2003. Diffusion of Innovations, 5th Editio. ed. The Free Press, New York, United States.
- Ryu, D.-H., Park, S.-W., Hwang, J.H., 2020. Association between Intention to Quit Cigarette Smoking and Use of Heated Tobacco Products: Application of Smoking Intensity Perspective on Heated Tobacco Product Users. *Int. J. Environ. Res. Public Health* 17, 8471. <https://doi.org/10.3390/ijerph17228471>
- Shamah-Levy, T., Vielma-Orozco, E., Heredia-Hernández, O., Romero-Martínez, M., Mojica-Cuevas, J., Cuevas-Nasu, L., Santaella-Castell, J., Rivera-Dommarco, J., 2020. Encuesta Nacional de Salud y Nutrición 2018-19: Resultados Nacionales. Instituto Nacional de Salud Pública, Cuernavaca, México.
- Sugiyama, T., Tabuchi, T., 2020. Use of Multiple Tobacco and Tobacco-Like Products Including Heated Tobacco and E-Cigarettes in Japan: A Cross-Sectional Assessment of the 2017 JASTIS Study. *Int. J. Environ. Res. Public Health* 17, 2161. <https://doi.org/10.3390/ijerph17062161>
- Sutanto, E., Miller, C., Smith, D.M., Brennan, K., Hyland, A., Cummings, K.M., Quah, A.C.K., Xu, S.S., Fong, G.T., Ouimet, J., Yoshimi, I., Mochizuki, Y., Tabuchi, T., O'Connor, R.J., Goniewicz, M.L., 2020. Concurrent Daily and Non-Daily Use of Heated Tobacco Products with Combustible Cigarettes: Findings from the 2018 ITC Japan Survey. *Int. J. Environ. Res. Public Health* 17, 2098. <https://doi.org/10.3390/ijerph17062098>
- Sutanto, E., Miller, C., Smith, D.M., O'connor, R.J., Quah, A.C.K., Cummings, K.M., Xu, S., Fong, G.T., Hyland, A., Ouimet, J., Yoshimi, I., Mochizuki, Y., Tabuchi, T., Goniewicz, M.L., 2019. Prevalence, use behaviors, and preferences among users of heated tobacco products: Findings from the 2018 itc Japan survey. *Int. J. Environ. Res. Public Health* 16. <https://doi.org/10.3390/ijerph16234630>
- Tabuchi, T., Gallus, S., Shinozaki, T., Nakaya, T., Kunugita, N., Colwell, B., 2018. Heat-not-burn tobacco product use in Japan: its prevalence, predictors and perceived symptoms from exposure to secondhand heat-not-burn tobacco aerosol. *Tob. Control* 27, e25 LP-e33. <https://doi.org/10.1136/tobaccocontrol-2017-053947>
- Tabuchi, T., Shinozaki, T., Kunugita, N., Nakamura, M., Tsuji, I., 2019. Study Profile: The Japan “Society and New Tobacco” Internet Survey (JASTIS): A Longitudinal Internet

- Cohort Study of Heat-Not-Burn Tobacco Products, Electronic Cigarettes, and Conventional Tobacco Products in Japan. *J. Epidemiol.* 29, 444–450.
<https://doi.org/10.2188/jea.JE20180116>
- Tanigaki, J., Poudyal, H., 2019. Challenges and opportunities for greater tobacco control in Japan. *Int. J. Drug Policy* 70, 78–86.
<https://doi.org/https://doi.org/10.1016/j.drugpo.2019.05.008>
- Thompson, M., Fong, G., Boudreau, C., Driezen, P., Li, G., Gravely, S., Cummings, K., Heckman, B., O'Connor, R., Thrasher, J.F., Nahhas, G., Borland, R., Yong, H., McNeill, A., Hitchman, S., Quah, A.C., 2019. Methods of the IT4 Four Country Smoking and Vaping Survey, wave 1 (2016). *Addiction* 114 Suppl 1, 6–14.
<https://doi.org/10.1111/add.14528>
- Tompkins, C.N.E., Burnley, A., McNeill, A., Hitchman, A.C., 2020. Factors that influence smokers' and ex-smokers' use of IQOS: a qualitative study of IQOS users and ex-users in the UK. *Tob. Control* tobaccocontrol-2019-055306.
<https://doi.org/10.1136/tobaccocontrol-2019-055306>
- Wu, Y.S., Wang, M.P., Ho, S.Y., Li, H.C.W., Cheung, Y.T.D., Tabuchi, T., Kwong, A.C.S., Lai, V., Lam, T.H., 2019. Heated tobacco products use in Chinese adults in Hong Kong: a population-based cross-sectional study. *Tob. Control* 29, 277 LP – 281.
<https://doi.org/10.1135/tobaccocontrol-2018-054719>
- Zavala-Arciniega, L., Gutiérrez-Torres, D.S., Reynales-Shigematsu, L.M., Barrientos-Gutiérrez, I., Fleischer, N.L., Meza, R., Thrasher, J.F., 2020. Cigarros con cápsulas de sabor en México: prevalencia, proporción de uso entre fumadores y predictores de consumo. *Ensanut 2018-19. Salud Publica Mex.* 62, 820–828.
<https://doi.org/10.21149/11566>




Figure 1. Weighted percentages about the reasons for use HTPs and e-cigarettes among adult smokers. Histograms show percentage and 95% CIs *p-value: p<0.01

Table 1. Analytic sample characteristics (n=6,500)

	n [#]	Unweighted % (95% CI)						Weighted % (95% CI)		
Survey										
November 2019	1,321	20.3	(19.3	-	21.3)	23.5	(## - ##
March 2020	1,282	19.7	(18.8	-	20.7)	23.2	(## - ##
July 2020	1,272	19.6	(18.6	-	20.5)	18.3	(## - ##
November 2020	1,309	20.1	(19.2	-	21.1)	16.3	(## - ##
March 2021	1,316	20.2	(19.3	-	21.2)	18.6	(## - ##
Sex										
Male	3,411	52.5	(51.3	-	53.7)	71.7	(## - ##
Female	3,089	47.5	(46.3	-	48.7)	28.3	(## - ##
Age										
18-29	1,759	27.1	(26.0	-	28.1)	34.7	(## - ##
30-39	1,973	30.4	(29.2	-	31.5)	29.1	(## - ##
40-49	1,277	19.6	(18.7	-	20.6)	13.3	(8.9 - ##
>50	1,491	22.9	(21.9	-	24.0)	22.8	(## - ##
Education										
Middle school and less	622	9.6	(8.9	-	10.3)	75.1	(71 - ##
High school / technical/ some college	3,622	55.7	(54.5	-	56.9)	17.8	(15 - ##

University and more	2,256	34. 7	(33. 6	- 35. 9)	7.1	(5.9	- 8.5)
Household income									
Less than 8,000 MXN	1,552	23. 9	(22. 8	- 24. 9)	42.4	(## #	- ## #)
8,001 to 15,000 MXN	2,070	31. 8	(30. 7	- 33. 0)	41.8	(## #	- ## #)
15,001 to >20,000 MXN	2,878	44. 3	(43. 1	- 45. 5)	15.7	(## #	- ## #)
Smoking frequency									
Non-daily	3,309	50. 9	(49. 7	- 52. 1)	52.9	(## #	- ## #)
Daily <=5 cigarettes	1,473	22. 7	(21. 6	- 23. 7)	16.5	(## #	- ## #)
Daily >5 cigarettes	1,718	26. 4	(25. 4	- 27. 5)	30.6	(## #	- ## #)
Recent quit attempt									
No	3,879	59. 7	(58. 5	- 60. 9)	58.9	(## #	- ## #)
Yes	2,621	40. 3	(39. 1	- 41. 5)	41.1	(## #	- ## #)
Intention to quit (next 6 months)									
No	4,155	63. 9	(62. 8	- 65. 1)	67.3	(## #	- ## #)
Yes	2,345	36. 1	(34. 9	- 37. 2)	32.7	(## #	- ## #)
E-cigarette use frequency									
Exclusive smoker	4,227	65. 0	(63. 9	- 66. 2)	96.5	(## #	- ## #)
Sporadic dual user	1,365	21. 0	(20. 0	- 22. 0)	2.1	(1.5	- 2.9)
Frequent dual user	908	14. 0	(13. 1	- 14. 8)	1.5	(1.1	- 2.0)
Partners/family smoke									
No	2,248	34. 6	(33. 4	- 35. 7)	29.1	(## #	- ## #)
Yes	4,252	65. 4	(64. 3	- 66. 6)	70.9	(## #	- ## #)
Partners/family use e-cigarette									
No	4,909	75. 5	(74. 5	- 76. 6)	91.6	(## #	- ## #)
Yes	1,591	24. 5	(23. 4	- 25. 5)	8.4	(5.6	- #)
Family use HTPs									
No	5,629	86. 6	(85. 8	- 87. 4)	97.4	(## #	- ## #)
Yes	871	13. 4	(12. 6	- 14. 2)	2.6	(1.9	- 3.5)
Friends smoke									
No	1,143	17. 6	(16. 7	- 18. 5)	28.9	(## #	- ## #)
Yes	5,357	82. 4	(81. 5	- 83. 3)	71.1	(## #	- ## #)
Friends use e-cigarette									
No	4,113	63. 3	(62. 1	- 64. 4)	87.0	(## #	- ## #)
Yes	2,387	36. 7	(35. 6	- 37. 9)	13.0	(9.4	- #)
Friends use HTPs									
No	5,452	83. 9	(83. 0	- 84. 8)	97.1	(## #	- ## #)
Yes	1,048	16. 1	(15. 2	- 17. 0)	2.9	(2.2	- 3.9)
Perceived harmfulness of HTPs relative to cigarettes									
HTPs are equally or more harmful than cigarettes	1,397	21. 5	(20. 5	- 22. 5)	18.8	(## #	- ## #)

HTPs are less harmful than cigarettes	1,121	17.2	(16.3 - 18.2)	8.1	(5.5 - ##)
I don't know	309	4.8	(4.2 - 5.3)	4.8	(# - ##)
Unaware	3,673	56.5	(55.3 - 57.7)	68.4	(## - ##)
Information sources related to HTPS (in the past 30 days)					
On the Internet					
No	4,567	70.3	(69.2 - 71.4)	83.9	(## - ##)
Yes	1,933	29.7	(28.6 - 30.8)	16.1	(## - ##)
Inside shops/ stores					
No	5,638	86.7	(85.9 - 87.6)	94.9	(## - ##)
Yes	862	13.3	(12.4 - 14.1)	5.1	(2.8 - 9.1)
Outside shops/ stores					
No	5,333	82.0	(81.1 - 83.0)	93.1	(## - ##)
Yes	1,167	18.0	(17.0 - 18.9)	6.9	(4.3 - ##)
In newspapers or magazines					
No	5,423	83.4	(82.5 - 84.3)	92.5	(## - ##)
Yes	1,077	16.6	(15.7 - 17.5)	7.5	(4.9 - ##)
Aware of HTPs	2,827	43.5	(42.3 - 44.7)	31.6	(## - ##)
Ever tried HTPs	1,217	18.7	(17.8 - 19.7)	5.0	(3.6 - ##)
Current use of HTPs	665	10.2	(9.5 - 11.0)	1.1	(0.8 - 1.5)

§Unweighted data

Table 2. Weighted multinomial logistic regression of factors associated with the use of HTPs in adult smokers and vapers, keep those who never tried HTPs as the reference (N=6,500)

Sex	N e v e r t r i e d H T P s	Ever tried HTPs					Current HTP use				
		%	R R (95% CI)	P- v a l u e	AR RR § (95% CI)	p- val ue	%	RR (95% CI)	P- v a l u e	A R R § (95% CI)	P- v a l u e

Male	9 3 7	5 . 3	R e f		Re f		1.0	Re f		Re f					
Female	9 4 6	4 . 2	0 7 9	(0 - 2)	0 . 6 1 3	1.2 8 4	(0 - 2)	0.4 78	1.2 0	(0.5 - 2)	0 . 5 3	1 . 6 3	(# - 3)	0 . 5 2	3 8 2
Age															
18-29	9 3 8	4 . 4	R e f		Re f		1.8	Re f		Re f					
30-39	9 0 4	8 . 7	2 0 7	(0 - 8)	0 . 3 2 9	2.2 3 5	(0 - 5)	0.0 66	0.9 8	(0.2 - 0)	0 . 9 5	1 . 0 3	(# - 2)	0 . 2 4	6 2 4
40-49	9 6 6	2 . 6	0 5 8	(0 - 1)	0 . 2 3 4	1.0 7 3	(0 - 2)	0.8 91	0.9 6	(0.1 - 1)	0 . 4 5 1	1 . 2 0	(# - 6)	0 . 7 4	5 0 4
>50	9 7 2	2 . 4	0 5 2	(0 - 1)	0 . 2 5 7	1.1 2 8	(0 - 3)	0.8 36	0.4 2	(0.0 - 0)	0 . 6 8 1	1 . 0 2	(# - 5)	0 . 5 1	3 8 5
Edu cati on															
Mid dle scho ol and less	9 4 6	4 . 8	0 9 3	(0 - 2)	0 . 5 3 4	2.4 3 0	(0 - 6)	0.0 81	0.6 1	(0.0 - 0)	< 0 . 2 3 0 1	0 . 3 5 0 1	(# - 1)	0 . 2 6 4	0 8
High scho ol/ tech nical / some colle ge	9 2 9	5 . 7	1 1 3	(0 - 1)	0 . 8 8 0	2.7 8 0	(1 - 5)	0.0 06	1.4 6	(0.1 - 0)	< 0 . 4 2 0 1	0 . 6 5 0 1	(# - 1)	0 . 5 4	0 2
Univ ersit y and more	8 9 9	4 . 8	R e f		Re f			5.3	Re f			Re f			
Hou seho ld inco me															
Less than 8,00 0M XN	9 3 1	5 . 9	R e f		Re f			1.1	Re f			Re f			
8,00 1 to 15,0 00M XN	9 5 6	3 . 7	0 6 2	(0 - 2)	0 . 4 6	1.2 6 4	(0 - 2)	0.5 87	0.7 9	(0.2 - 1)	0 . 4 9	1 . 2 6 7	(# - 2)	0 . 6 4	9 8

15,001 to >20,000 MXN	9 5 0 (0 - 3) 0	2.9 4 (1 - 8) 0.0 38	2.3 2.1 (1.0 - 4) 0 1 (# - 5) 0	
Smoking frequency	2
Non-daily	8 9 2 6 9	0 1	4 7 # 6 3	
Daily <=5 cigarettes	0 9 7 4 9 3	6 1	2 3 0 7 8 9	
Daily >5 cigarettes				
Recent quit attempt				
No	9 5 3 (0 - 5) 0	1.7 6 (0 - 4) 0.1 80	2.9 4.6 (2.1 - 9) < 6 (# - 1) <	
Yes	1	0	
Intention to quit	2 9 3 9 4 0	7 7 2 1	7 0 6 5 0 1	
No	9 3 5 (0 - 2) 0	0.9 7 (0 - 2) 0.9 33	1.4 0 (0.6 - 3) 0 1 (# - 3) 0	
In the next six months	4 5 1 1 4 2	4 0 6 3	2 4 3 8 9 8 7	
E-cigarette use frequency				
Exclusive smoker	9 4 5 (2 - 2) 0	2.1 9 (0 - 5) 0.0 70	## 49. (## - #) < 6 (# - 1) <	
Sporadic dual	6 2 7 (2 - 2) 0	5 4 . . 8 . 9	
	. . 9 1 . 0	9 1	

user	1	1	0	7	7	0	4	3	#	0	5	3	0																		
					1	2			#	0		7	0																		
										1			1																		
Frequent	5	1	4	(2	-	8)	<	0.6	(0	-	2)	0.5	##	20	(##	-	#)	<	1	(#	-	3)	<
dual	2	0	3	13	#	8.7	##	##	.	.	.	0	1	(#	-	2)	0
user	.	.	2	1	4	1	5	6						
	3	5	7	7	2	0	6	3	#	0	2	9	0	0	0	#	0	6						1				9	0	0	
						1																							0	0	1
Partners/family smokers																															
No	9	3	R						Re		0.7	Re		R																	
	6	.	e						f			f		f																	
	.	0	f																												
	3																														
Yes	9	5	2	(0	-	6)	0	0.8	(0	-	1)	0.5	1.2	1.8	(0.8	-	3)	0	0	(#	-	0)	0
	3	2	67		2		5
	.	8	0	6	0	2	4	6	.	.	4	6
	0	0	6	8	2	0	1	3	0	0	1	3	1	2	4																
Partners/family use e-cigarette																															
No	9	2	R						Re		0.3	Re		R																	
	7	.	e						f			f		f																	
	.	6	f																												
	1																														
Yes	6	3	1	(5	-	7)	<	3.8	(1	-	8)	0.0	9.3	44.	(##	-	8)	<	1	(#	-	3)	<
	0	0	9	.	2	0	9	.	.	9	01		00		##	6	0	0	(#	-	3)	0		
	.	.	.	0	.	.	.	7	4	.	7	4
	1	7	1	6	1	0	9	4	0	0	9	4	0	0	0																
Family use HT Ps																															
No	9	4	R						Re		0.4	Re		R																	
	5	.	e						f			f		f																	
	.	0	f																												
	6																														
Yes	3	4	3	(#	-	8)	<	5.3	(1	-	1)	0.0	##	17	(##	-	#)	<	5	(#	-	1)	0
	1	2	2	#	6	0	2	.	.	2	.	5	.	.	.	02	#	6.7	##	##	#	0	.	(#	-	9)	.	.	
	.	.	.	#	8
	8	7	5	#	1	0	8	0	0	0	8	0	0	0	0																
Friends smoke																															
No	9	3	R						Re		0.5	Re		R																	
	6	.	e						f			f		f																	
	.	3	f																												
	2																														
Yes	9	5	1	(0	-	5)	0	0.7	(0	-	1)	0.5	1.3	2.4	(0.8	-	7)	0	1	(#	-	3)	0
	3	6	47		8		0
	.	6	7	5	7	3	3	8	.	.	3	8
	1	7	5	1	4	2	3	3	0	0	2	3	7	1	4																

[illegible]

No	9 5 .5	4 .1	R e f	Re f	0.4	Re f	R e f
Yes	4 2 .3	3 4 .5	1 9 .1	(. 0 2	7 0 .0	- 0 .8	5 0 .4
				<	1.4	(0 5 8	- 3 6 4
					0.4	## 24	12 4.6 8
					(## ##	- # #	# # #
					<	2	(# #
						-	6
)	0
							0
							7
							8
							0
							9

	HTP s are equa lly or mor e har mful than cigar ettes	7 6 .	1 6 .	1 .	(.	0 2	- 4	4 8) .	0 8	0.8 3	(.	0 4	- 6	1 2) .	0.5 86	7.4 7	3.0 7	(.	1.1 7	- .	8 0) .	0 0	1 3	(.	# #	- 7	2 8) .	0 5	3 5
I don't kno w/ 0	9 9 .	1 .	0 0	(.	0 0	- 2	0 5) .	< .	0.1 6	(.	0 5	- 4	0 4) .	0.0 03	0.0 04	0.0 0	(.	0.0 0	- .	0 0) .	< .	0 0	(.	# #	- 0	0 8) .	< .	0 0	

by all variables in
the table and date
of survey

Table 3. Variables related to HTPs among current users (n=665)

	n [§]	Weighted % (95% CI)				
Frequency of use						
Daily	99	13.5	(8.1	-	###)
Not daily but at least once a week	334	46.9	(33.6	-	###)
Less than once a week, but at least once a month	232	39.6	(27.5	-	###)
Most recent variety of heatsticks/heets used						
Sienna Selection (intense tobacco)	139	15.1	(9.3	-	###)
Amber Selection (toasted tobacco and nuts)	117	28.7	(16.4	-	###)
Yellow Selection (smooth tobacco with citrus)	84	8.5	(4.9	-	###)
Blue Selection (smooth menthol)	161	31.8	(20.5	-	###)
Turquoise Selection (deep menthol)	70	5.0	(3.4	-	7.4)
Purple Wave (menthol with fruits)	84	9.9	(4.3	-	###)
Other and I don't know	10	1.1	(0.5	-	2.5)
How did you obtain their HTPs device?						
I bought it	376	44.2	(32.1	-	###)
It was a gift from a relative or friend	211	34.0	(21.6	-	###)
It was a free sample	40	12.3	(5.4	-	###)
It was borrowed	38	9.4	(3.0	-	###)
Where did you buy their HTPs device?						
Online	213	68.7	(57.5	-	###)
Vape shop or tobacconist	83	18.5	(11.7	-	###)
Department store or supermarket	62	10.4	(6.8	-	###)
Other	18	2.5	(1.4	-	4.5)

[§]Unweighted data

Funding

This research was funded by of the Fogarty International Center of the National Institutes of Health under award number R01 TW010652. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health

Contributors

Lizeth Cruz-Jimenez, and James F. Thrasher have contributed to all sections of manuscript, conceptualization, preparation, writing, review, and editing. Inti Barrientos-Gutiérrez, Edna Arillo-Santillán, Katia Gallegos-Carrillo, and Rosibel Rodríguez-Bolaños contributed to conceptualization, supervision, review, and editing. Luis Zavala-Arciniega and Shannon Gravely contributed to methodology, supervision, and writing – review. All authors have approved the final article for submission.

Conflicts of Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Conflicts of Interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Contributors: Lizeth Cruz-Jimenez, and James F. Thrasher have contributed to all sections of manuscript, conceptualization, preparation, writing, review, and editing. Inti Barrientos-Gutiérrez, Edna Arillo-Santillán, Katia Gallegos-Carrillo, and Rosibel Rodríguez-Bolaños contributed to conceptualization, supervision, review, and editing. Luis Zavala-Arciniega and Shannon Gravely contributed to methodology, supervision, and writing – review. All authors have approved the final article for submission.

Highlights

- Uptake of HTPs is low among Mexican smokers.
- Strong correlates of current HTPs use include frequent e-cigarette use and use of e-cigarettes and HTPs among close network members.
- Social acceptability and lower harm compared to cigarettes were the top reasons for use.

Journal Pre-proof