

Surveillance of moist snuff: total nicotine, moisture, pH, un-ionized nicotine, and tobacco-specific nitrosamines

Patricia Richter, Knachelle Hodge, Stephen Stanfill, Liqin Zhang, Clifford Watson

Received 6 September 2007; accepted 11 March 2008

In 2005, approximately 2.3% of U.S. adults used smokeless tobacco. Moist snuff leads all types of smokeless tobacco in revenues and marketing expenditures. The U.S. Surgeon General has concluded that smokeless tobacco use can lead to nicotine addiction. The National Toxicology Program of the National Institutes of Health has classified smokeless tobacco as a human carcinogen. Tobacco-specific nitrosamines (TSNAs) are potent carcinogens in smokeless tobacco products, and the pH of the product influences the content of un-ionized nicotine which is the form of nicotine most rapidly absorbed in the mouth. The Centers for Disease Control and Prevention analyzed 40 top-selling brands of moist snuff to measure nicotine, moisture, pH, un-ionized nicotine, and TSNAs, including 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL). The study findings indicate that moist snuff brands varied widely in content of rapidly absorbed, addictive un-ionized nicotine (500-fold range) and of carcinogenic TSNAs (18-fold range). Product characteristics such as packaging and moisture content appeared to be correlated with concentrations of un-ionized nicotine, and flavor characteristics of low-priced brands may correlate with TSNA concentrations. These findings warrant further study in light of (a) the marketing of smokeless tobacco for use in places where smoking is prohibited, (b) the promotion of smokeless tobacco as a harm-reduction product, and (c) the ever-expanding number of highly flavored smokeless varieties brought to the market.

Introduction

In 2005, approximately 2.3% of U.S. adults used smokeless tobacco (Centers for Disease Control and Prevention [CDC], 2006). Smokeless tobacco use is much higher among U.S. adults with low levels of education and for some racial and ethnic groups such as American Indians and Alaska Natives (Nelson, Mowery, Tomar, Marcus, Giovino, & Zhao, 2006). Among the smokeless tobacco products, moist snuff

receives the greatest advertising and marketing support and leads in revenues and marketing expenditures. In 2005, expenditures on advertising and promoting moist snuff totaled \$210.43 million compared with \$16.75 million for loose leaf tobacco and chewing tobacco, \$72,000 for plug tobacco and twist tobacco, and \$103,000 for Scotch snuff and dry snuff (Federal Trade Commission [FTC], 2007). Dollar sales for moist snuff (\$2.23 billion) represented 85.2% of total sales of smokeless tobacco in 2005, and more pounds of moist snuff were sold in 2005 (75.7 million) than all other smokeless tobacco products combined (FTC, 2007). From 1986 to 2003, use of moist snuff increased more than 80-fold while large decreases in sales of chewing tobacco and dry snuff products fueled an overall decline in sales volumes for smokeless tobacco (Nelson et al., 2006).

In 1986, the U.S. Surgeon General concluded that smokeless (“spit”) tobacco is not a safe substitute for smoking cigarettes or cigars and that smokeless

Patricia Richter, Ph.D., Office on Smoking and Health, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, Atlanta, GA; Knachelle Hodge, M.P.H., University of Maryland School of Medicine, Baltimore, MD; Stephen Stanfill, M.S., Liqin Zhang, M.S., Clifford Watson, Ph.D., Division of Laboratory Sciences, National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, GA.

Correspondence: Dr. Patricia Richter, Centers for Disease Control and Prevention, Office on Smoking and Health, 4770 Buford Highway, N.E., MS K-50, Atlanta, GA 30341, USA. Tel: +1 (770) 488-5703; Fax: +1 (770) 488-5848; E-mail: pir1@cdc.gov

tobacco products cause various cancers, noncancerous oral conditions, and nicotine addiction (U.S. Department of Health and Human Services [USDHHS], 1986). The National Toxicology Program of the National Institutes of Health determined that smokeless tobacco is a human carcinogen (National Toxicology Program [NTP], 2005). Smokeless tobacco causes pancreatic cancer (International Agency for Research on Cancer [IARC], 2007) and has also been linked to cardiovascular disease, dental diseases, and adverse pregnancy outcomes (Critchley & Unal, 2003). *N'*-Nitrosonornicotine (NNN) and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) are quantitatively the most prevalent strong carcinogens in smokeless tobacco. 4-[Methylnitrosamino]-1-[3-pyridyl]-1-butanol (NNAL) is also carcinogenic while NAB (*N'*-nitrosoanabasine) is a moderately potent carcinogen and *N'*-nitrosoanatabine (NAT) is generally considered inactive as a rodent carcinogen (Hecht & Hoffmann 1988). Both NNK and NNN have recently been classified as reasonably anticipated by the IARC as carcinogenic to humans (Group 1) (IARC, 2007). Nicotine is the substance in smokeless tobacco that causes addiction; un-ionized (unprotonated or free) nicotine is the form of nicotine most rapidly and easily absorbed in the mouth (Armitage & Turner, 1970; USDHHS, 1986, 1988). The pH of a smokeless tobacco product is a major determinant of the amount of un-ionized nicotine delivered to the user (Tomar & Henningfield, 1997).

The purpose of this study was to independently measure nicotine, moisture, pH, and un-ionized nicotine, and TSNA in 40 top-selling brands of moist snuff. In addition, we examined the relation between select product characteristics (e.g., packaging, cut, flavor) and un-ionized nicotine levels and TSNA.

Materials and methods

Materials

Forty brand varieties of U.S. moist snuff smokeless tobacco products manufactured by five companies were selected for analysis on the basis of market share. Unit-based market share measured in percent (%) was determined from proprietary sales data (Scan-Trac data, Grocery Channel, 50 Retail Markets; ACNielsen, Schaumburg, Illinois). Collectively, these brand varieties held more than a 97% share of the U.S. market in the year they were purchased. CDC personnel purchased five packs of each of the 40 moist snuff products from Internet retailers or in the metropolitan Atlanta area in 2004. Samples were stored in the original packaging at

–70°C until testing. The five packs of each product were pooled, and the pooled samples were used for all analyses.

Methods

Samples were analyzed for moisture, pH, and nicotine content as previously described (Richter & Spierto, 2003) except for the use of mass selective detection instead of flame ionization detection. This method has been shown to obtain equivalent results for nicotine over the range investigated. TSNA were measured by CDC in the following manner. Samples were ground and approximately 0.25 g was weighed into a stainless steel extraction cell. 100 ng each of ¹³C₆-labeled *N'*-nitrosonornicotine (NNN), 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK), *N'*-nitrosoanabasine (NAB), *N'*-nitrosoanatabine (NAT) and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) were added to the vial. Samples were then extracted into ethylacetate at 100°C and 1500 psi for 5 min using a Dionex ASE® 200 Accelerated Solvent Extractor (Dionex, Sunnyvale, California). The extracts were dried under nitrogen and reconstituted in 10 mL of methanol. Then samples were diluted 10-fold in 20 mM ammonium acetate aqueous solution for analysis by high-performance liquid chromatography and electrospray ionization tandem mass spectrometry (LC/MS/MS). Details of the mass spectral detection method were published previously (Wu, Ashley, & Watson, 2003).

Statistical analysis

Statistical analyses were performed by using the Statistical Analysis System software (SAS for Windows, Version 9.1; SAS Institute, Cary, NC). Statistical comparisons were performed by using analysis of variance and Student *t*-test; *p* < .05 was considered statistically significant.

Results

Total nicotine, moisture, pH, and un-ionized nicotine

Levels of nicotine and TSNA measured in the 40 brands of moist snuff products are expressed as the amount detected per gram of wet tobacco weight. Results are shown with brands ordered by total nicotine content (Table 1). One brand, Oregon Mint Snuff, had no detectable nicotine. Excluding Oregon Mint Snuff, total nicotine concentrations ranged from 4.4 milligrams of nicotine per gram of tobacco (mg/g) (Hawken Rough Wintergreen) to 25.0 mg/g (W.B. Cut Regular) (mean, 11.9 mg/g; median, 12.6 mg/g). Total un-ionized nicotine ranged from 0.01 mg/g (Hawken Rough Wintergreen) to 7.8 mg/g

(Kodiak Ice Long Cut Regular) (mean, 3.8 mg/g; median 4.4 mg/g). Un-ionized nicotine, expressed as a percentage of total nicotine, ranged from 0.3% (Hawken Rough Wintergreen) to 79.9% (Kodiak Ice Long Cut Regular). Moisture content for all 40 brands ranged from 3.2% (Oregon Mint Snuff) to 54.5% (Rooster Long Cut Bold Wintergreen) (mean, 50.4%; median 53.1%).

Total nicotine was not correlated with un-ionized nicotine ($r=.01$) or pH ($r=-.11$) while pH demonstrated a highly significant ($p<.0001$) positive correlation with both total ($r=.92$) and percentage of un-ionized nicotine ($r=.90$). The results demonstrated positive correlation between pH and total un-ionized nicotine ($p<.0001$) and between pH and percentage of un-ionized nicotine ($p<.0001$), and these correlations showed strong statistical significance. There was also significant positive correlation between moisture and total un-ionized nicotine ($p<.0002$), percentage of un-ionized nicotine ($p<.0016$), and pH ($p<.0001$). Moist snuff is sold in a variety of tobacco cuts, including fine cut, long cut, and rough cut. In this study there was a significant difference in the pH ($p=.0057$) between the groups.

Five of the varieties of moist snuff in our sample consisted of tobacco in a portion pouch. The other varieties were packaged as loose tobacco. For loose tobacco varieties, but not portion pouches, we observed statistically significant correlations of moisture content with total un-ionized nicotine ($p=.0003$), percentage of un-ionized nicotine ($p=.0034$), and pH ($p<.0001$). In addition, mean pH ($p=.01$), mean concentrations of moisture ($p=.009$), total nicotine ($p=.01$), total un-ionized nicotine ($p=.001$), as well as mean percentage of un-ionized nicotine ($p=.005$), were significantly higher in loose smokeless tobacco varieties than in pouch varieties.

In general, the correlations between pH and un-ionized nicotine and moisture existed regardless of retail price or market share. In general, for brands that sold at more or less than the mean retail price of \$2.95 per unit or the median market share of 0.72%, the positive correlations between pH and un-ionized nicotine (total or percentage) and between moisture and un-ionized nicotine and between moisture and pH were statistically significant. Brands that sold for less than the mean retail price were exceptions. The results for these brands demonstrated a significant positive correlation of moisture with total un-ionized nicotine ($p=.008$) but not with percentage of un-ionized nicotine. There was also a significant positive correlation between moisture and total nicotine ($p=.0001$) and cut and total nicotine ($p=.044$) for brands with a high market share.

One brand of moist snuff had a market share approximately 3 times higher than that of the brand

with the next highest market share and more than 100 times higher than the brand with the lowest market share. When this brand was excluded from the analysis, market share was positively correlated with the percentage of un-ionized nicotine ($p=.04$) and pH ($p=.05$) in the high market share group, and this correlation was statistically significant.

Tobacco-specific nitrosamines

Because Oregon Mint Snuff had no detectable TSNAs, this brand was excluded from further statistical analysis of TSNAs. Ranges and differences between the lowest and highest concentrations, means, and medians for concentrations measured in nanograms of each TSNA per gram of tobacco (wet weight) were as follows (Table 1):

- NAB—range, 123–4,242 ng/g (34-fold difference); mean, 616 ng/g; median, 395 ng/g;
- NAT—range, 938–31,866 ng/g (34-fold); mean, 6,452 ng/g; median, 4,369 ng/g;
- NNN—range, 2,204–42,554 ng/g (19-fold); mean, 6,883 ng/g; median, 5,271 ng/g;
- NNK—range, 382–9,950 ng/g (26-fold); mean, 1,811 ng/g; median, 1,261 ng/g;
- NNAL—range, 21–1,412 ng/g (67-fold); mean, 169 ng/g; median, 120 ng/g.

The sums of concentrations of all TSNAs ranged from 4.87 $\mu\text{g/g}$ to 90.02 $\mu\text{g/g}$ (4,874–90,024 ng/g) (18-fold difference) (Table 1). The sums of concentrations of carcinogenic TSNAs (NNN+NNK+NNAL) ranged from 2.99 $\text{m}\mu\text{g/g}$ to 53.92 $\text{m}\mu\text{g/g}$ (2,986–53,916 ng/g) (18-fold difference). The brand variety with the highest concentrations of TSNAs had total TSNA and carcinogenic TSNA concentrations 3 times higher than the brand variety with the next highest concentration. The ratios of the most potent carcinogenic TSNAs to total TSNAs ($[\text{NNN}+\text{NNK}+\text{NNAL}]/[\text{NAB}+\text{NAT}+\text{NNN}+\text{NNK}+\text{NNAL}]$) ranged from .47 (Timber Wolf Fine Cut Regular) to .78 (Hawken Rough Wintergreen). The average ratio of carcinogenic TSNAs to total TSNAs was .55; the median ratio was .56.

We observed no correlations between concentrations of carcinogenic TSNAs or total TSNAs with total nicotine, un-ionized nicotine, moisture, or retail price or market share in the overall sample, and none were observed in evaluations of low versus high retail price or market share groups. There was a statistically significant positive correlation between flavor in moist snuff and total TSNAs ($p=.02$) and carcinogenic TSNAs ($p=.02$) among brands in the low retail price group (below mean retail price). Flavored varieties are those that have a flavor descriptor in their name (e.g., Skoal Long Cut Berry Blend). For the high market share varieties (above the median

Table 1. Total nicotine, moisture, pH, un-ionized nicotine, and tobacco-specific nitrosamines in top-selling brands of moist snuff^a.

| Brand variety | Manufacturer | Total nicotine (mean mg/g ± SD) | Total moisture (mean % ± SD) | pH (mean ± SD) | Un- ionized nicotine (%) | Un-ionized nicotine (mg/g) | NAB (mean ng/ g ± SD) | NAT (mean ng/g ± SD) | NNN (mean ng/ g ± SD) | NNK (mean ng/ g ± SD) | NNAL (mean ng/ g ± SD) | Sum of total TSNAs (ng/g) | Sum of NNN, total NNK, NNAL (ng/g) |
|--------------------------------------|--------------------------|---------------------------------------|---------------------------------|-------------------|-----------------------------------|----------------------------------|-----------------------------|-------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------------|---|
| W.B. Cut Regular | United States Tobacco | 25.03 ± 0.10 | 33.0 ± 0.30 | 5.71 ± 0.01 | 0.5 | 0.12 | 142 ± 9.0 | 3652 ± 220 | 4184 ± 200 | 446 ± 62.0 | 23 ± 12.0 | 8447 | 4653 |
| Red Seal Fine Cut Wintergreen | United States Tobacco | 14.19 ± 0.08 | 53.3 ± 0.20 | 7.55 ± 0.00 | 25.3 | 3.59 | 123 ± 12.0 | 1907 ± 141 | 2599 ± 217 | 583 ± 33.0 | 60 ± 22.0 | 5272 | 3242 |
| Timber Wolf Long Cut, Wintergreen | Swedish Match | 14.01 ± 0.03 | 54.3 ± 0.40 | 7.62 ± 0.02 | 28.7 | 4.01 | 169 ± 9.5 | 2543 ± 346 | 2432 ± 263 | 902 ± 183 | 50 ± 13.6 | 6096 | 3384 |
| Timber Wolf Fine Cut Regular | Swedish Match | 13.91 ± 0.02 | 50.2 ± 0.30 | 7.41 ± 0.01 | 19.8 | 2.75 | 1196 ± 270 | 12056 ± 1100 | 9556 ± 1571 | 2287 ± 642 | 123 ± 17.3 | 25218 | 11966 |
| Timber Wolf Straight Regular | Swedish Match | 13.88 ± 0.08 | 54.0 ± 0.40 | 7.75 ± 0.01 | 34.9 | 4.86 | 193 ± 54.1 | 2874 ± 682 | 2877 ± 1127 | 919 ± 179 | 37 ± 24.8 | 6901 | 3833 |
| Skoal Long Cut Berry Blend | United States Tobacco | 13.56 ± 0.04 | 53.5 ± 0.20 | 7.52 ± 0.02 | 24.0 | 3.25 | 649 ± 142 | 6964 ± 346 | 7630 ± 1576 | 1739 ± 600 | 207 ± 26.8 | 17190 | 9576 |
| Skoal Straight Cut Regular | United States Tobacco | 13.35 ± 0.13 | 54.2 ± 0.10 | 7.72 ± 0.01 | 33.9 | 4.47 | 836 ± 83.2 | 9252 ± 275 | 8903 ± 1285 | 3383 ± 949 | 151 ± 27.6 | 22527 | 12438 |
| Skoal Key | United States Tobacco | 13.19 ± 0.10 | 53.3 ± 0.10 | 7.90 ± 0.01 | 43.1 | 5.68 | 4242 ± 1174 | 31866 ± 1288 | 42554 ± 16049 | 9950 ± 392 | 1412 ± 222 | 90024 | 53916 |
| Grizzly Fine Cut Regular | Conwood | 13.17 ± 0.07 | 53.8 ± 0.20 | 7.81 ± 0.00 | 38.1 | 5.03 | 862 ± 111 | 10908 ± 2968 | 8690 ± 878 | 1692 ± 149.2 | 281 ± 9.0 | 22434 | 10664 |
| Skoal Fine Cut Wintergreen | United States Tobacco | 13.13 ± 0.17 | 53.4 ± 0.20 | 7.66 ± 0.01 | 30.9 | 3.99 | 650 ± 180 | 7642 ± 1894 | 8395 ± 1765 | 2031 ± 575 | 103 ± 22.5 | 18821 | 10529 |
| Red Seal Fine Cut Regular | United States Tobacco | 13.09 ± 0.03 | 54.3 ± 0.60 | 8.00 ± 0.02 | 49.7 | 6.37 | 1676 ± 531 | 14544 ± 2552 | 14080 ± 4472 | 3878 ± 910 | 247 ± 35.7 | 34424 | 18205 |
| Copenhagen Long Cut Regular | United States Tobacco | 12.99 ± 0.02 | 53.7 ± 0.30 | 7.91 ± 0.01 | 44.0 | 5.67 | 181 ± 48.2 | 1793 ± 313 | 2204 ± 577 | 911 ± 880 | 21 ± 14.2 | 5110 | 3136 |
| Timberwolf Fine Cut Wintergreen | Swedish Match | 12.98 ± 0.50 | 52.8 ± 0.50 | 7.22 ± 0.01 | 13.7 | 1.76 | 205 ± 25.0 | 2738 ± 174 | 2874 ± 489 | 1061 ± 37.3 | 43 ± 13.5 | 6921 | 3978 |
| Red Seal Long Cut Wintergreen | United States Tobacco | 12.97 ± 0.03 | 53.6 ± 0.70 | 7.67 ± 0.01 | 31.0 | 4.03 | 139 ± 9.6 | 1749 ± 268 | 2570 ± 132 | 382 ± 89.6 | 35 ± 3.3 | 4874 | 2986 |
| Copenhagen Regular | United States Tobacco | 12.94 ± 0.29 | 52.6 ± 0.10 | 7.73 ± 0.01 | 34.2 | 4.36 | 365 ± 42.4 | 3879 ± 738 | 3987 ± 302 | 960 ± 218 | 62 ± 1.7 | 9253 | 5009 |
| Skoal Long Cut Regular | United States Tobacco | 12.83 ± 0.15 | 53.3 ± 0.50 | 7.77 ± 0.00 | 36.0 | 4.59 | 1046 ± 15.0 | 16054 ± 1228 | 14031 ± 614 | 3428 ± 38.0 | 269 ± 8.0 | 34828 | 17728 |
| Timber Wolf Cool Wintergreen | Swedish Match | 12.77 ± 0.04 | 54.4 ± 0.60 | 7.84 ± 0.00 | 39.5 | 5.04 | 208 ± 119 | 2684 ± 1063 | 2557 ± 905 | 653 ± 400 | 26 ± 10.5 | 6128 | 3237 |
| Skoal Long Cut Wintergreen | United States Tobacco | 12.71 ± 0.29 | 53.5 ± 0.30 | 7.75 ± 0.01 | 35.5 | 4.41 | 501 ± 116 | 5101 ± 823 | 6455 ± 535 | 1309 ± 786 | 85 ± 13.8 | 13452 | 7849 |
| Skoal Long Cut Classic | United States Tobacco | 12.70 ± 0.21 | 54.0 ± 0.40 | 7.83 ± 0.02 | 39.3 | 4.99 | 730 ± 96.5 | 8305 ± 1179 | 8186 ± 411 | 2217 ± 411 | 178 ± 24.9 | 19616 | 10581 |
| Skoal Long Cut Cherry | United States Tobacco | 12.64 ± 0.22 | 52.6 ± 0.20 | 7.49 ± 0.02 | 23.4 | 2.89 | 530 ± 107 | 6271 ± 795 | 6237 ± 1368 | 1579 ± 957 | 65 ± 8.6 | 14682 | 7881 |
| Renegades Wintergreen | Swedish Match | 12.41 ± 0.08 | 53.0 ± 0.60 | 6.70 ± 0.01 | 4.5 | 0.56 | 177 ± 17.4 | 3092 ± 485 | 2651 ± 245 | 761 ± 86.9 | 23 ± 5.7 | 6705 | 3436 |
| Copenhagen Black Bourbon | United States Tobacco | 12.32 ± 0.05 | 54.4 ± 0.20 | 7.82 ± 0.01 | 38.9 | 4.79 | 406 ± 54.5 | 3366 ± 361 | 4407 ± 468 | 2264 ± 2171 | 56 ± 4.9 | 10498 | 6726 |

Table 1. Continued.

| Brand variety | Manufacturer | Total nicotine (mean mg/g ± SD) | Total moisture (mean % ± SD) | pH (mean ± SD) | Un- ionized nicotine (%) | Un-ionized nicotine (mg/g) | NAB (mean ng/ g ± SD) | NAT (mean ng/g ± SD) | NNN (mean ng/ g ± SD) | NNK (mean ng/ g ± SD) | NNAL (mean ng/ g ± SD) | Sum of NNN, NNK, total NNAL | |
|--------------------|---------------|---------------------------------------|---------------------------------|-------------------|-----------------------------------|----------------------------------|-----------------------------|-------------------------|-----------------------------|-----------------------------|------------------------------|-----------------------------------|----------------|
| | | | | | | | | | | | | TSNAs (ng/g) | NNK, (ng/g) |
| Redwood Regular | Swisher | 12.17 ± 0.07 | 52.2 ± 0.50 | 7.44 ± 0.01 | 21.0 | 2.55 | 1348 ± 61.2 | 7787 ± 547 | 7918 ± 1347 | 2308 ± 275 | 255 ± 28.7 | 19616 | 10481 |
| Skoal Long Cut | United States | 12.08 ± 0.08 | 52.2 ± 0.40 | 7.85 ± 0.01 | 40.3 | 4.87 | 395 ± 137 | 5418 ± 1365 | 5541 ± 1072 | 1451 ± 468 | 120 ± 41.5 | 12925 | 7112 |
| Mint | Tobacco | | | | | | | | | | | | |
| Silver Creek Fine | Swisher | 11.96 ± 0.04 | 52.6 ± 1.20 | 7.08 ± 0.00 | 10.3 | 1.24 | 637 ± 62.0 | 6091 ± 289 | 5946 ± 192 | 2925 ± 95.0 | 268 ± 14.0 | 15867 | 9139 |
| Cut Wintergreen | | | | | | | | | | | | | |
| Rooster Long Cut | United States | 11.46 ± 0.13 | 54.4 ± 0.20 | 7.85 ± 0.01 | 40.3 | 4.61 | 523 ± 73.2 | 4369 ± 594 | 5271 ± 583 | 1092 ± 925 | 70 ± 18.2 | 11326 | 6433 |
| Icy Mint | Tobacco | | | | | | | | | | | | |
| Rooster Long Cut | United States | 11.06 ± 0.07 | 54.5 ± 0.10 | 7.86 ± 0.01 | 40.6 | 4.49 | 1432 ± 110 | 11713 ± 386 | 11081 ± 575 | 2731 ± 143 | 150 ± 2.7 | 27107 | 13962 |
| Bold Wintergreen | Tobacco | | | | | | | | | | | | |
| Silver Creek Long | Swisher | 10.26 ± 0.14 | 50.7 ± 0.10 | 6.64 ± 0.01 | 4.0 | 0.41 | 1120 ± 108 | 11711 ± 2919 | 10540 ± 1264 | 4321 ± 559 | 302 ± 48.1 | 27994 | 15162 |
| Cut Straight | | | | | | | | | | | | | |
| Regular | | | | | | | | | | | | | |
| Silver Creek Long | Swisher | 10.21 ± 0.09 | 50.6 ± 0.60 | 7.08 ± 0.01 | 10.2 | 1.04 | 212 ± 20.6 | 2297 ± 60 | 2722 ± 249 | 944 ± 31.3 | 97 ± 2.7 | 6272 | 3763 |
| Cut Wintergreen | | | | | | | | | | | | | |
| Kodiak Straight | Conwood | 9.85 ± 0.07 | 52.4 ± 0.30 | 8.29 ± 0.01 | 65.1 | 6.39 | 335 ± 50.4 | 5082 ± 1212 | 4476 ± 822 | 1147 ± 563 | 217 ± 36.5 | 11257 | 5840 |
| Cut Regular | | | | | | | | | | | | | |
| Kodiak Ice Long | Conwood | 9.80 ± 0.08 | 53.2 ± 0.10 | 8.62 ± 0.06 | 79.9 | 7.81 | 314 ± 23.0 | 4255 ± 149 | 4550 ± 181 | 1233 ± 97.0 | 184 ± 11.0 | 10536 | 5967 |
| Cut Regular | | | | | | | | | | | | | |
| Kodiak Wintergreen | Conwood | 9.56 ± 0.07 | 52.5 ± 0.40 | 8.49 ± 0.02 | 75.6 | 7.14 | 492 ± 88.2 | 7380 ± 1474 | 6923 ± 1674 | 1744 ± 503 | 418 ± 100.6 | 16958 | 9085 |
| Grizzly Long Cut | Conwood | 9.40 ± 0.08 | 51.9 ± 0.10 | 8.42 ± 0.03 | 71.5 | 6.71 | 215 ± 6.1 | 3100 ± 20 | 3633 ± 250 | 975 ± 36.9 | 168 ± 16.7 | 8091 | 4776 |
| Wintergreen | | | | | | | | | | | | | |
| Cougar Long Cut | Conwood | 9.14 ± 0.06 | 53.0 ± 0.10 | 8.16 ± 0.03 | 58.0 | 5.27 | 213 ± 21.7 | 2813 ± 348 | 4039 ± 375 | 1261 ± 206 | 153 ± 10.0 | 8478 | 5453 |
| Natural | | | | | | | | | | | | | |
| Cougar Long Cut | Conwood | 8.72 ± 0.07 | 52.2 ± 0.80 | 7.44 ± 0.02 | 20.8 | 1.83 | 708 ± 45.0 | 9465 ± 518 | 9352 ± 1072 | 1660 ± 58.0 | 306 ± 32.0 | 21491 | 11318 |
| Wintergreen | | | | | | | | | | | | | |
| Cougar Regular | Conwood | 8.48 ± 0.10 | 53.9 ± 0.20 | 8.07 ± 0.01 | 52.9 | 4.48 | 238 ± 15.6 | 2907 ± 99 | 4567 ± 155 | 1053 ± 41 | 132 ± 10.9 | 8897 | 5752 |
| Skoal Bandits Mint | United States | 7.64 ± 0.10 | 44.5 ± 0.70 | 6.72 ± 0.00 | 4.8 | 0.37 | 231 ± 24.6 | 4257 ± 427 | 6069 ± 764 | 931 ± 286 | 52 ± 18.5 | 11540 | 7052 |
| | Tobacco | | | | | | | | | | | | |
| Skoal Bandits | United States | 6.23 ± 0.09 | 48.0 ± 0.30 | 6.84 ± 0.00 | 6.3 | 0.39 | 181 ± 10.6 | 2789 ± 168 | 4552 ± 109 | 751 ± 202 | 31 ± 4.7 | 8303 | 5334 |
| Wintergreen | Tobacco | | | | | | | | | | | | |
| Hawken Rough | Conwood | 4.42 ± 0.02 | 27.4 ± 0.20 | 5.54 ± 0.01 | 0.3 | 0.01 | 202 ± 22.1 | 938 ± 238 | 3203 ± 190 | 757 ± 241 | 117 ± 21.3 | 5218 | 4077 |
| Wintergreen | | | | | | | | | | | | | |
| Oregon Mint Snuff | Oregon Mint | ND | 3.23 ± 0.34 | 6.51 ± 0.03 | ND | ND | ND | ND | ND | ND | ND | ND | ND |
| | Snuff | | | | | | | | | | | | |

Note. SD, standard deviation; NAB, N'-nitrosoanabasine; NAT, N'-nitrosoanatabine; NNN, N'-nitrosoanornicotine; NNK, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNAL, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAs, tobacco-specific nitrosamines; ND, not detected.^a Concentrations of nicotine and TSNAs are expressed as amount detected per gram of wet tobacco.

market share), the correlation between cut and total TSNA ($p=.03$) and carcinogenic TSNA ($p=.05$) was significant. No correlation was observed between packaging (loose or in portion pouches) and either total TSNA or carcinogenic TSNA.

Discussion

Under the Comprehensive Smokeless Tobacco Health Education Act of 1986 (15 U.S.C. 4401 et seq., Pub. L. 99-252), smokeless tobacco manufacturers report annually to the Centers for Disease Control and Prevention (CDC) on the total nicotine content, moisture content, pH, and estimated un-ionized nicotine in smokeless tobacco products. This law does not require information on concentrations of other chemicals such as tobacco-specific nitrosamines (TSNAs) in smokeless tobacco products. The data that are reported to CDC under this act are considered "trade secret" and cannot be released to the public in accordance with 5 U.S.C. 552(b)(4) and 18 U.S.C. 1905. The purpose of this study was to provide consumers, researchers, and public health officials with information on pH, levels of total nicotine, un-ionized nicotine, moisture and TSNA in popular smokeless tobacco brands.

Our previous findings demonstrated that brands of moist snuff with the largest amount of un-ionized nicotine are the most frequently sold brands (Richter & Spierto, 2003). In that study, we used data on market share in an aggregated brand family level (e.g., Skoal). In the present study, this limitation was addressed by using data on market share and retail price reported on a brand variety level (e.g., Skoal Long Cut Berry Blend). The investigation was also expanded to include TSNA concentrations. As in the earlier study, pH proved to be a good predictor of un-ionized nicotine content in the moist snuff products. Market share and retail price were not factors in the relationships between pH and un-ionized nicotine among the brand varieties tested. Exclusion of one very high market share brand resulted in a significant positive correlation between market share and pH and market share and the percentage of un-ionized nicotine among brands in the high market share group (above median market share). This finding suggests that the most popular brands of moist snuff continue to deliver the highest amounts of the rapidly absorbed un-ionized nicotine to consumers.

The observation that moisture concentrations in moist snuff positively correlate with both pH and un-ionized nicotine content is intriguing and, to our knowledge, has not been previously reported. Moisture level is a product characteristic controlled by the manufacturer. Among the brand varieties tested in this study, tobacco cut, another product

characteristic, was also positively correlated with pH, both overall and in the high retail price varieties. Others have reported that tobacco cut influences oral absorption of nicotine from smokeless tobacco and that binding agents used with larger cuts of tobacco may allow the smokeless tobacco to be formed in a compacted mass that slows nicotine release (Connolly, 1995; Djordjevic, Hoffmann, Glynn, & Connolly, 1995). Although our analysis was limited by the small number of varieties sold in portion pouches, the results suggest that packaging is a factor in product moisture and un-ionized nicotine content, as well as pH level, but not TSNA content.

The nicotine dose received by a smokeless tobacco user is influenced by the amount of nicotine in the product, the tobacco cut size, pH, and to a much lesser extent, behavioral factors such as the rate of expectoration (Tomar and Henningfield, 1997; USDHHS, 1986). An additional factor that influences the oral absorption of nicotine is the buffering capacity of saliva. The buffering capacity of moist snuff products is higher than the buffering capacity of saliva and the saliva pH for moist snuff products is determined by the acid-base buffering capacities of both the saliva and the moist snuff (Ciolino, McCauley, Fraser, & Wolnik, 2001). Consuming acidic foods such as coffee or cola reduces saliva pH and oral nicotine absorption (Henningfield, Radzius, Cooper, & Clayton, 1990). Among the top-selling U.S. smokeless tobacco products in our current study, there was more than a 5-fold difference in the range of total nicotine concentrations and more than a 500-fold difference in the range of total un-ionized nicotine concentrations. Five brands of moist snuff reported on in 2003 were included in this study (Richter & Spierto, 2003). Compared with the earlier samples, two had lower concentrations of total nicotine and lower (more acidic) pH; one had a higher total nicotine and higher pH; and two were mixed: one had lower total nicotine but higher pH; and one had higher total nicotine but lower pH. Increasing the alkalinity of smokeless tobacco promotes the absorption of nicotine and increases its physiological effects (Tomar & Henningfield, 1997). It is not known whether these differences in nicotine and pH were related to changes in product formulation or whether they reflect variations in an agricultural-based product.

One goal of this study was to survey concentrations of TSNA in popular brands of moist snuff and look for correlations between concentrations of TSNA and other chemical and physical properties. We found an almost 20-fold difference in the range of sums for total TSNA and for carcinogenic TSNA. The difference in the range of NNAL concentrations was almost 70-fold (Table 1), a much wider range than that for other carcinogenic TSNA (NNN,

19-fold; NNK, 26-fold). NNAL has been detected at low concentrations in toombak, a Sudanese fermented snuff (Prokopczyk et al., 1995), but no data were available on NNAL in commercial smokeless tobacco products. Our study found no significant correlation between moisture and TSNAs. However, others have shown that TSNA concentrations in moist snuff products stored at room temperature increase more in high-moisture products than in low-moisture products (Andersen, Fleming, Hamilton-Kemp, & Hildebrand, 1993).

Concentrations of NAB, NAT, NNN, and NNK measured by gas chromatography with a thermal-energy analyzer were recently reported for a variety of tobacco products including traditional moist snuff and new oral tobacco products (Stepanov, Jensen, Hatsukami, & Hecht, 2006). Comparison of the results from that study with those from our study showed that although there is good agreement in total concentrations of these TSNAs for some brands (e.g., Copenhagen long cut); differences are notable for other brands. For example, in our study using high-performance liquid chromatography and electrospray ionization tandem mass spectrometry (LC/MS/MS), the sum of these four TSNAs for Skoal Long Cut Straight Regular was 35 µg/g. In the study by Stepanov and associates, it was 9.2 µg/g. The reproducibility and accuracy of the method we used was reported previously (Wu et al., 2003). This difference may be related to differences in extraction efficiency or year-to-year variations in TSNA concentrations. Further investigation is warranted. In the work by Stepanov and associates, TSNA concentrations in new oral tobacco products were approximately threefold lower than those in traditional smokeless tobacco brands, and the overall ratio of carcinogenic TSNAs to total TSNAs ((NNN+NNK)/(NAB+NAT+NNN+NNK)) was .67, which falls within the range of ratios for the same four TSNAs (.47–.78) in our study. (NNAL concentrations were not reported in the study by Stepanov and associates.) Those results suggest that it is possible to produce commercial smokeless tobacco products with lower TSNA concentrations. However, the biological relevance of small reductions in TSNA concentrations in smokeless tobacco, a human carcinogen, cannot be determined from these data.

The results of our study are subject to two limitations. First, the study samples were the 40 top-selling brands of domestic moist snuff, which were selected by using sales-based data on market share. This sampling captured brands ranging in average retail price from \$1.36 to \$4.13 per unit. Despite the large number of brands, there may be regional, niche, or deeply discounted varieties that are not represented by the brands included in this

study. The study design focused on the most widely available popular brands of moist snuff and did not include new or modified smokeless tobacco products that are being sold or are in test market in some regions of the country. The products were purchased from Internet retailers who ship tobacco products to consumers throughout the United States. Thus they are typical of products available to consumers nationwide.

Second, results are based on five samples from the same lot of a brand that were pooled before analyses. Although this approach ensured an adequate quantity of the moist snuff product to complete all analyses with a single pooled sample, it did not allow evaluation of variations within a product or variations between lots or locations. Consequently, we did not draw direct between-brand conclusions from our comparisons.

To conclude, results from this study indicate that top-selling brands of moist snuff varied widely in content of addictive un-ionized nicotine (500-fold) and carcinogenic TSNAs (18-fold). Product characteristics such as packaging and moisture content appeared to be correlated with concentrations of un-ionized nicotine. In addition, the findings indicate that flavor characteristics of low-priced brands may correlate with TSNA concentrations. Such findings warrant further study in light of (a) the marketing of smokeless tobacco for use in places where smoking is prohibited; (b) the promotion of smokeless tobacco as a harm-reduction product (Henningfield, Rose, & Giovino, 2002; Nelson et al., 2006); and (c) the ever-expanding number of highly flavored smokeless varieties brought to the market (e.g., Vanilla Blend Skoal and Kayak Peach Long Cut).

Acknowledgments

The authors would like to acknowledge Lily T. Jia for assistance with the analysis of the nicotine content of the smokeless tobacco samples, Dr. Michelle O'Hegarty and Bill Marx for assistance in obtaining the smokeless tobacco brands, and Dr. Brian Armour for calculating the market share data.

All research was supported by internal funds of the Centers for Disease Control and Prevention (CDC). Use of trade names is for informational purposes only and in no way implies endorsement by the U.S. Government, the U.S. Department of Health and Human Services, or CDC. The findings and conclusions in this report are those of the authors and do not necessarily represent the views of CDC.

The authors report no conflicts of interest.

References

- Andersen, R. A., Fleming, P. D., Hamilton-Kemp, T. R., & Hildebrand, D. F. (1993). pH changes in smokeless tobaccos undergoing nitrosation during prolonged storage: Effects of moisture, temperature, and duration. *Journal of Agricultural and Food Chemistry*, 41, 968–972.
- Armitage, A. K., & Turner, D. M. (1970). Absorption of nicotine in cigarette and cigar smoke through the oral mucosa. *Nature*, 226, 1231–1232.

- Centers for Disease Control and Prevention. (2006). Tobacco use among adults—United States, 2005. *Morbidity and Mortality Weekly*, 55, 1145–1148.
- Ciolino, L. A., McCauley, H. A., Fraser, D. B., & Wolnik, K. A. (2001). The relative buffering capacities of saliva and moist snuff: implications for nicotine absorption. *Journal of Analytical Toxicology*, 25, 15–25.
- Connolly, G. N. (1995). The marketing of nicotine addiction by one oral snuff manufacturer. *Tobacco Control*, 4, 73–79.
- Critchley, J. A., & Unal, B. (2003). Health effects associated with smokeless tobacco: A systematic review. *Thorax*, 58, 435–443.
- Djordjevic, M. V., Hoffmann, D., Glynn, D. T., & Connolly, G. N. (1995). U.S. commercial brands of moist snuff, 1994—Assessment of nicotine, moisture, and pH. *Tobacco Control*, 4, 62–66.
- Hecht, S. S., & Hoffmann, D. (1988). Tobacco-specific nitrosamines, an important group of carcinogens in tobacco and tobacco smoke. *Carcinogenesis*, 9, 875–884.
- Henningfield, J. E., Radzius, A., Cooper, T. M., & Clayton, R. R. (1990). Drinking coffee and carbonated beverages blocks absorption of nicotine from nicotine polacrilex gum. *Journal of the American Medical Association*, 264, 1560–1564.
- Henningfield, J. E., Rose, C., & Giovino, G. (2002). Brave new world of tobacco disease prevention. Promoting dual tobacco-product use? *American Journal of Preventive Medicine*, 23, 226–228.
- International Agency for Research on Cancer. (2007). IARC Monographs on the evaluation of carcinogenic risk to humans. Volume 89: Smokeless tobacco and some tobacco-specific nitrosamines. Lyon: World Health Organization.
- National Toxicology Program. (2005). Report on Carcinogens, Eleventh Edition; U.S. Department of Health and Human Services, Public Health Service, National Toxicology Program.
- Nelson, D. E., Mowery, P., Tomar, S., Marcus, S., Giovino, G., & Zhao, L. (2006). Trends in smokeless tobacco use among adults and adolescents in the United States. *American Journal of Public Health*, 96, 897–905.
- Prokopczyk, B., Wu, M., Cox, J. E., Amin, S., Desai, D., Idris, A. M., & Hoffmann, D. (1995). Improved methodology for the quantitative assessment of tobacco-specific N-nitrosamines in tobacco by supercritical fluid extraction. *Journal of Agricultural and Food Chemistry*, 43, 916–922.
- Richter, P., & Spierto, F. W. (2003). Surveillance of smokeless tobacco nicotine, pH, moisture, and unprotonated nicotine content. *Nicotine & Tobacco Research*, 5, 885–889.
- Stepanov, I., Jensen, J., Hatsukami, D., & Hecht, S. S. (2006). Tobacco-specific nitrosamines in new tobacco products. *Nicotine & Tobacco Research*, 8, 309–313.
- Tomar, S. L., & Henningfield, J. E. (1997). Review of the evidence that pH is a determinant of nicotine dosage from oral use of smokeless tobacco. *Tobacco Control*, 6, 219–225.
- U.S. Department of Health and Human Services. (1986). *The Health Consequences of Using Smokeless Tobacco: A Report of the Surgeon General*. NIH Publication No. 86-2874. Bethesda, MD: Department of Health and Human Services, Public Health Service, Office of the Surgeon General.
- U.S. Department of Health and Human Services. (1988). *The Health Consequences of Smoking: Nicotine Addiction: A Report of the Surgeon General*. DHHS publication No. (CDC) 88-8406. Rockville, MD: Department of Health and Human Services, Public Health Service, Office of the Surgeon General.
- U.S. Federal Trade Commission. (2007). *Smokeless Tobacco Report for the Years 2002–2005*. Washington: Federal Trade Commission.
- Wu, W., Ashley, D. L., & Watson, C. H. (2003). Simultaneous determination of five tobacco-specific nitrosamines in mainstream cigarette smoke by isotope dilution liquid chromatography/electrospray ionization tandem mass spectrometry. *Analytical Chemistry*, 75, 4827–4832.