

5"

5"

1"

3"

Prazi-Med™ (praziquantel powder for immersion)

For use in ornamental finfish,
except *Callichthyidae* and
Clariidae catfish

Net Contents:
One 10 gram mylar packet

For aquariums and ponds equal
to or less than 250 gallons
250 mg Measuring Scoop included

LEGAL STATUS—In order to be legally marketed, a new animal drug intended for a minor species must be Approved, Conditionally Approved, or Indexed by the Food and Drug Administration. **THIS PRODUCT IS INDEXED**— MIF 900-054. Extra-label use is prohibited.

THIS PRODUCT IS NOT TO BE USED IN ANIMALS INTENDED FOR USE AS FOOD FOR HUMANS OR FOOD-PRODUCING ANIMALS.

It is a violation of federal law to use this product in a manner other than directed in the labeling.



Before using this drug, read package insert for full product information. For accurate dosing, use the enclosed 250 mg scoop and refer to the dosing chart for use with the 10 gram mylar packet included in the package insert.

USES: PRAZI-MED™ praziquantel powder for immersion is used for the treatment of internal parasites (flukes and tapeworms) and external parasites (flukes) susceptible to praziquantel in ornamental finfish, except *Callichthyidae* and *Clariidae* catfish.

DIRECTIONS: PRAZI-MED™ is intended for use only in personal home aquariums, personal home ponds, or public aquariums and zoos.

WARNINGS: USER SAFETY WARNINGS: Not for use in humans. Keep out of reach of children. Use in well-ventilated area and avoid inhalation. Avoid dust formation. Wear eye protection, dust mask, and chemical-resistant gloves when handling the drug. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco products, or using the toilet. **INHALATION:** If inhaled and breathing becomes difficult, move victim to fresh air. Seek medical attention. **INGESTION:** If ingested: If the person is conscious, rinse mouth out with water and then drink plenty of water. Seek

medical attention.

EYE CONTACT: If in eyes: Rinse eye immediately with water for at least 15 minutes. If symptoms occur, seek medical attention.

SKIN CONTACT and/or CLOTHING CONTAMINATION: Wash skin with soap and water, and launder clothing with detergent. If symptoms occur, seek medical attention.

STORAGE, HANDLING AND DISPOSAL: Store at or below 25°C (77°F) and protect from light. Keep product and empty container away from heat and sources of ignition. See package insert for further instructions for disposing of treated water. Do not conduct water changes during the treatment period. Prior to conducting a water change, subject treated water to chemical filtration for at least 24 hours prior to discharge to remove residual drug. Dispose of unused product, spills, and treated water in a manner that will not lead to contamination of surface water. Contact your State Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance pertaining to disposal of unused product.

QUESTIONS/COMMENTS? To report side effects, for technical assistance or to request an SDS, contact AQION at 1-719-256-0044. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

Manufactured for: AQION LLC
539 W. Commerce Street, Suite 6825, Dallas, TX 75208
PRAZI-MED™ is a trademark of KDG Impresa LLC.
© 2025 KDG Impresa LLC. All rights reserved. ~ Rev. 01/2026

Lot #:
Expiration date:
NDC# 43457-534-10



BLEED AREA CUTTING AREA
 SAFE AREA PASTING AREA