



LUCEMYRA® IS THE ONLY FDA-APPROVED, NON-OPIOID MEDICINE
PROVEN TO HELP WITH SYMPTOMS OF OPIOID WITHDRAWAL.

[Read More](#)



PATIENT PRESCRIBING INFO

[Click to see full Important Safety Information & Prescribing Information](#)



FIND A PROVIDER

[Expand](#)





PATIENT PRESCRIBING INFO



FIND A PROVIDER



PATIENT RESOURCES

See a collection of interactive and downloadable content, and links to external organizations.

[SEE RESOURCES](#)



LUCEMYRA can help relieve withdrawal symptoms



[Click to see full Important Safety Information & Prescribing Information](#)

Expand ^





LUCEMYRA can help relieve withdrawal symptoms



Significantly improves the percentage of patients who successfully complete withdrawal treatment



Demonstrated safety profile in studies



Provides the biggest improvement in symptom relief when symptoms were often most severe



2020 ASAM National Practice Guideline: Recommendations for Treating Opioid Withdrawal

The American Society of Addiction Medicine (ASAM) recognizes two main strategies for managing opioid withdrawal: gradually tapering the dose of the current opioid(s), or using a non-opioid medication, like LUCEMYRA (lofexidine).



[Click to see full Important Safety Information & Prescribing Information](#)

[Expand](#) ^





2020 ASAM National Practice Guideline: Recommendations for Treating Opioid Withdrawal

The American Society of Addiction Medicine (ASAM) recognizes two main strategies for managing opioid withdrawal: gradually tapering the dose of the current opioid(s), or using a non-opioid medication, like LUCEMYRA (lofexidine).

Regarding the use of alpha-2 adrenergic agonists, the Guideline states:

“Lofexidine should therefore be the preferred choice for withdrawal management in an outpatient setting, where monitoring of blood pressure and management of hypotension is more difficult.”

-American Society of Addiction Medicine, 2020





Regarding the use of alpha-2 adrenergic agonists, the Guideline states:

“Lofexidine should therefore be the preferred choice for withdrawal management in an outpatient setting, where monitoring of blood pressure and management of hypotension is more difficult.”



-American Society of Addiction Medicine, 2020



[Click to see full Important Safety Information & Prescribing Information](#)

[Expand](#) ^



[Privacy Policy](#) | [Terms & Conditions](#)

This product information is intended for US patients only. Any products discussed herein may have different product labeling in different countries. Under license from Britannia Pharmaceuticals LTD to BioCorRx Pharmaceuticals Inc. USWM LLC is the distributor of LUCEMYRA® in the United States and its territories. LUCEMYRA® is a registered trademark of BioCorRx Pharmaceuticals Inc. Luminat® is a registered trademark of BioCorRx Pharmaceuticals Inc.





withdrawal, gradually tapering the dose of the current opioid(s), or using a non-opioid medication, like LUCEMYRA (lofexidine).

Regarding the use of alpha-2 adrenergic agonists, the Guideline states:

[Collapse](#)

What is LUCEMYRA?

LUCEMYRA is a non-opioid prescription medicine used in adults to help with the symptoms of opioid withdrawal that may happen when you stop taking an opioid suddenly. LUCEMYRA will not completely prevent the symptoms of opioid withdrawal and is not a treatment for opioid use disorder.

Important Safety Information

LUCEMYRA can cause serious side effects, including low blood pressure, slow heart rate, and fainting. Watch for symptoms of low blood pressure or heart rate, including dizziness, lightheadedness, or feeling faint at rest or when quickly standing up; if you experience these symptoms, call your healthcare provider right away and do not take your next dose of LUCEMYRA until you have talked to your healthcare provider. Avoid becoming dehydrated or overheated and be careful not to stand up too suddenly from lying or sitting, as these may increase your risk of low blood pressure and fainting. When your treatment is complete, you will need to stop taking LUCEMYRA gradually, or your blood pressure could increase. After a period of not using opioid drugs, you can become more sensitive to the effects of opioids if you start using them again. This may increase your risk of overdose and death. Before taking LUCEMYRA, tell your healthcare provider about all your medical conditions, including if you have low blood pressure, slow heart rate, any heart problems including history of heart attack or a condition called long QT syndrome, liver or kidney problems, or if you drink alcohol. Tell your healthcare provider if you are pregnant, plan on becoming pregnant, or are breastfeeding; it is not known if LUCEMYRA can harm your unborn baby or whether LUCEMYRA passes into your breast milk. Especially tell your healthcare provider if you take benzodiazepines, barbiturates, tranquilizers, or sleeping pills, as taking these with LUCEMYRA can cause serious side effects. The most common side effects of LUCEMYRA include low blood pressure or symptoms of low blood pressure such as lightheadedness, slow heart rate, dizziness, sleepiness, and dry mouth. **To report SUSPECTED ADVERSE REACTIONS or product complaints, contact BioCorRx Pharmaceuticals Inc at 1-833-LUCEMYRA. You may also report SUSPECTED ADVERSE REACTIONS to the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.** Click here to see full [Prescribing Information](#).

