


[Is It Time for Niktimvo?](#)
[About Niktimvo](#)
[How Niktimvo May Help](#)
[Taking Niktimvo](#)
[Possible Side Effects](#)
[Support and Resources](#)

Is your chronic graft-versus-host disease (GVHD) holding you back?

A CHANCE TO STAY ONE STEP AHEAD WITH NIKTIMVO

Niktimvo is the **FIRST** and **ONLY** treatment that takes a different approach to treat chronic GVHD in people who weigh at least 88.2 pounds (40 kg) and received at least 2 prior treatments that did not work.



See how Niktimvo may help

Niktimvo may offer fast and lasting responses

[Explore Study Results](#)


Is Niktimvo right for me?

Find out if Niktimvo is the right choice for you

[Learn When to Start](#)


Discover Niktimvo support and resources

Get more information about what you can expect with Niktimvo treatment

[Access Resources](#)

If you continue to experience chronic GVHD symptoms, ask your doctor if it's time for Niktimvo



IncyteCARES is here to support you through every step of your journey with Niktimvo. [See support](#)

INDICATIONS AND USAGE

Niktimvo™ (axatilimab-csfr) is a prescription medicine used to treat adults and children who weigh at least 88.2 pounds (40 kg) with chronic graft-versus-host disease (cGVHD) after receiving at least 2 prior treatments (systemic therapy) and they did not work.

It is not known if Niktimvo is safe and effective in adults and children weighing less than 88.2 pounds (40 kg).

IMPORTANT SAFETY INFORMATION

What are the possible side effects of Niktimvo?

Niktimvo may cause serious side effects, including:

Infusion-related reactions. Infusion-related reactions are common with Niktimvo and can be serious. Your healthcare provider will monitor you for infusion-related reactions during your treatment. If you have a reaction, your healthcare provider may temporarily or completely stop your treatment with Niktimvo. Tell your healthcare provider right away if you have fever, chills, rash, flushing, shortness of breath, trouble breathing, nausea, vomiting, or symptoms of high blood pressure such as chest pain, headaches, or blurred vision during an infusion of Niktimvo.

The most common side effects of Niktimvo include:

- infections
- increased blood level of liver enzymes
- decreased blood level of phosphate
- low red blood cell count (anemia)
- muscle, bone, or joint pain
- increased blood level of pancreatic enzymes
- low energy
- increased blood level of calcium
- increased blood level of a muscle enzyme
- increased blood level of a bone enzyme
- nausea
- headache
- diarrhea
- cough
- fever
- shortness of breath

These are not all the possible side effects of Niktimvo. Call your doctor for medical advice about side effects.

Before receiving Niktimvo, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had liver problems.
- are pregnant or plan to become pregnant. Niktimvo may harm your unborn baby.

Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with Niktimvo.
- You should use an effective method of birth control during your treatment and for 30 days after your last dose of Niktimvo. Talk to your healthcare provider about birth control methods that you can use during this time.
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with Niktimvo.

- are breastfeeding or plan to breastfeed. It is not known if Niktimvo passes into your breast milk. Do not breastfeed during treatment and for 30 days after your last dose of Niktimvo.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see the Full Prescribing Information, which includes a more complete discussion of the risks associated with Niktimvo.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

You may also report side effects to Incyte Medical Information at 1-855-463-3463.




[Is It Time for Niktimvo?](#)
[About Niktimvo](#)
[How Niktimvo May Help](#)
[Taking Niktimvo](#)
[Possible Side Effects](#)
[Support and Resources](#)

In a study of people experiencing chronic GVHD symptoms after taking at least 2 therapies,

Niktimvo™ (axatilimab-csfr) delivered fast and lasting responses

Study overview

Niktimvo was studied in 79 people who had received a stem cell transplant, received at least 2 prior treatments for chronic graft-versus-host disease (GVHD), and weighed at least 88.2 pounds (40 kg).

[How Niktimvo Was Studied](#)
[Possible Side Effects](#)

Most people achieved a response with Niktimvo

A response to treatment with Niktimvo was defined as:

- A resolution of all signs and symptoms of chronic GVHD (complete response), or
- Improvement in at least one affected area without worsening signs or symptoms in other areas (partial response)

75% of people (59 out of 79) experienced a response



Some people saw a fast response



Of the people who responded to treatment, half saw their first response in **6 weeks or less**

The time to first response ranged from **4 weeks to 22 weeks**.

Niktimvo provided lasting responses



More than half (60%) of the people who showed a response to Niktimvo continued to respond **for at least 1 year**

People achieved responses in multiple affected organs

A complete or partial response in each organ was evaluated by recording changes in chronic GVHD severity from the beginning of the study to the end.

[Click on the tiles below for more information about organ\(s\).](#)



Niktimvo offered symptom improvement

44 out of 79 people who received Niktimvo (56%) reported improvement in the specific chronic GVHD symptoms assessed in the study.

Symptom improvement was defined as a reduction in chronic GVHD symptoms compared to the start of the study.



Niktimvo may provide fast and lasting responses across a range of affected organs

Every person is unique. How your chronic GVHD progresses and how you respond to Niktimvo will depend on your individual circumstances.

Talk to your doctor about how people responded to Niktimvo in the study and the potential safety concerns. It may help you and your doctor decide if Niktimvo is right for you.

[See how IncyteCARES may help](#)
[Get Support](#)
[Explore possible side effects](#)
[Learn About Safety](#)

INDICATIONS AND USAGE

Niktimvo™ (axatilimab-csfr) is a prescription medicine used to treat adults and children who weigh at least 88.2 pounds (40 kg) with chronic graft-versus-host disease (cGVHD) after receiving at least 2 prior treatments (systemic therapy) and they did not work.

It is not known if Niktimvo is safe and effective in adults and children weighing less than 88.2 pounds (40 kg).

IMPORTANT SAFETY INFORMATION

What are the possible side effects of Niktimvo?

Niktimvo may cause serious side effects, including:

Infusion-related reactions. Infusion-related reactions are common with Niktimvo and can be serious. Your healthcare provider will monitor you for infusion-related reactions during your treatment. If you have a reaction, your healthcare provider may temporarily or completely stop your treatment with Niktimvo. Tell your healthcare provider right away if you have fever, chills, rash, flushing, shortness of breath, trouble breathing, nausea, vomiting, or symptoms of high blood pressure such as chest pain, headaches, or blurred vision during an infusion of Niktimvo.

The most common side effects of Niktimvo include:

- infections
- increased blood level of liver enzymes
- decreased blood level of phosphate
- low red blood cell count (anemia)
- muscle, bone, or joint pain
- increased blood level of pancreatic enzymes
- low energy
- increased blood level of calcium
- increased blood level of a muscle enzyme
- increased blood level of a bone enzyme
- nausea
- headache
- diarrhea
- cough
- fever
- shortness of breath

These are not all the possible side effects of Niktimvo. Call your doctor for medical advice about side effects.

Before receiving Niktimvo, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had liver problems.
- are pregnant or plan to become pregnant. Niktimvo may harm your unborn baby.

Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with Niktimvo.
- You should use an effective method of birth control during your treatment and for 30 days after your last dose of Niktimvo. Talk to your healthcare provider about birth control methods that you can use during this time.
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with Niktimvo.
- are breastfeeding or plan to breastfeed. It is not known if Niktimvo passes into your breast milk. Do not breastfeed during treatment and for 30 days after your last dose of Niktimvo.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see the Full Prescribing Information, which includes a more complete discussion of the risks associated with Niktimvo.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

You may also report side effects to Incyte Medical Information at 1-855-463-3463.

