

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use OCTAGAM 10% safely and effectively. See full prescribing information for OCTAGAM 10%.

OCTAGAM 10% [Immune Globulin Intravenous (Human)]

liquid solution for intravenous administration

Initial U.S. Approval: 2014

WARNING

THROMBOSIS, RENAL DYSFUNCTION AND ACUTE RENAL FAILURE

See full prescribing information for complete boxed warning

- Thrombosis may occur with immune globulin intravenous (IGIV) products, including OCTAGAM 10%. Risk factors may include: advanced age, prolonged immobilization, hypercoagulable conditions, history of venous or arterial thrombosis, use of estrogens, indwelling vascular catheters, hyperviscosity, and cardiovascular risk factors.
- Renal dysfunction, acute renal failure, osmotic nephropathy, and death may occur with the administration of Immune Globulin Intravenous (Human) (IGIV) products in predisposed patients. Renal dysfunction and acute renal failure occur more commonly in patients receiving IGIV products containing sucrose. OCTAGAM 10% does not contain sucrose.
- For patients at risk of thrombosis, renal dysfunction or renal failure, administer OCTAGAM 10% at the minimum infusion rate practicable. Ensure adequate hydration in patients before administration. Monitor for signs and symptoms of thrombosis and assess blood viscosity in patients at risk for hyperviscosity.

INDICATIONS AND USAGE

- OCTAGAM 10% is an immune globulin intravenous (human) liquid preparation indicated for the treatment of chronic immune thrombocytopenic purpura (ITP) in adults (1).

DOSAGE AND ADMINISTRATION

For intravenous use only.

Indication	Dose	Initial Infusion rate	Maintenance Infusion Rate (if tolerated)
Chronic ITP	1 g/kg daily for 2 consecutive days	1.0 mg/kg/min (0.01 mL/kg/min)	Up to 12.0 mg/kg/min (Up to 0.12 mL/kg/min)

- Ensure that patients with pre-existing renal insufficiency are not volume depleted; discontinue OCTAGAM 10% if renal function deteriorates (2, 5.2).
- For patients at risk of renal dysfunction or thrombotic events, administer OCTAGAM 10% at the minimum infusion rate practicable (5.2, 5.5).

DOSAGE FORMS AND STRENGTHS

Solution containing 10% IgG (0.1 g/L) (3, 16)

CONTRAINDICATIONS

- History of anaphylactic or severe systemic reactions to human immunoglobulin (4)
- IgA deficient patients with antibodies against IgA and a history of hypersensitivity (4)

WARNINGS AND PRECAUTIONS

- IgA-deficient patients with antibodies against IgA are at greater risk of developing severe hypersensitivity and anaphylactic reactions to OCTAGAM 10%. Epinephrine should be available immediately to treat any severe acute hypersensitivity reactions (5.1).
- Monitor renal function, including blood urea nitrogen and serum creatinine, and urine output in patients at risk of developing acute renal failure (5.2).
- Falsely elevated blood glucose readings may occur during and after the infusion of OCTAGAM 10% with testing by some glucometers and test strip systems (5.3).
- Hyperproteinemia, increased serum osmolality and hyponatremia may occur in patients receiving OCTAGAM 10% (5.4).
- Hemolysis that is either intravascular or due to enhanced red blood cell sequestration can develop subsequent to OCTAGAM 10% treatments. Risk factors for hemolysis include high doses and non-O-blood group. Closely monitor patients for hemolysis and hemolytic anemia (5.6).
- Aseptic Meningitis Syndrome may occur in patients receiving OCTAGAM 10%, especially with high doses or rapid infusion (5.7).
- Monitor patients for pulmonary adverse reactions (transfusion-related acute lung injury (TRALI)) (5.8).
- OCTAGAM 10% is made from human plasma and may contain infectious agents, e.g. viruses and, theoretically, the Creutzfeldt-Jakob disease agent (5.9).

ADVERSE REACTIONS

The most common adverse reactions reported in greater than 5% of subjects during a clinical trial were headache, fever and increased heart rate (6).

To report SUSPECTED ADVERSE REACTIONS, contact Octapharma at 1-866-766-4860 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

The passive transfer of antibodies may:

Confound the results of serological testing (7).

Interfere with the immune response to live viral vaccines, such as measles, mumps, and rubella (7).

USE IN SPECIFIC POPULATIONS

- Pregnancy: no human or animal data. Use only if clearly needed (8.1).
- Geriatric Use: In patients over age 65 or in any person at risk of developing renal insufficiency, do not exceed the recommended dose, and infuse OCTAGAM 10% at the minimum infusion rate practicable (8.5).

See 17 for PATIENT COUNSELING INFORMATION.

Revised: July 2014

FULL PRESCRIBING INFORMATION: CONTENTS*

BOXED WARNING: THROMBOSIS, RENAL DYSFUNCTION AND ACUTE RENAL FAILURE

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosage
- 2.2 Preparation and handling
- 2.3 Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Hypersensitivity
- 5.2 Renal Dysfunction/Failure
- 5.3 Blood Glucose Monitoring
- 5.4 Hyperproteinemia, Increased Serum Viscosity and Hyponatremia
- 5.5 Thrombosis
- 5.6 Hemolysis
- 5.7 Aseptic meningitis syndrome
- 5.8 Transfusion-Related Acute Lung Injury (TRALI)
- 5.9 Transmission of Infectious Agents
- 5.10 Monitoring Laboratory Tests

- 6 ADVERSE REACTIONS**
 - 6.1 Clinical Trials Experience
 - 6.2 Postmarketing Experience
 - 7 DRUG INTERACTIONS**
 - 8 USE IN SPECIFIC POPULATIONS**
 - 8.1 Pregnancy
 - 8.3 Nursing Mothers
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use
 - 10 OVERDOSAGE**
 - 11 DESCRIPTION**
 - 12 CLINICAL PHARMACOLOGY**
 - 12.1 Mechanism of Action
 - 12.3 Pharmacokinetics
 - 13 NON-CLINICAL TOXICOLOGY**
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
 - 13.2 Animal Toxicology and/or Pharmacology
 - 14 CLINICAL STUDIES**
 - 15 REFERENCES**
 - 16 HOW SUPPLIED/STORAGE AND HANDLING**
 - 17 PATIENT COUNSELING INFORMATION**
- *Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

OCTAGAM 10% [Immune Globulin Intravenous (Human)] Liquid Preparation

WARNING: THROMBOSIS, RENAL DYSFUNCTION AND ACUTE RENAL FAILURE

- **Thrombosis may occur with immune globulin intravenous (IGIV) products, including OCTAGAM 10%. Risk factors may include: advanced age, prolonged immobilization, hypercoagulable conditions, history of venous or arterial thrombosis, use of estrogens, indwelling central vascular catheters, hyperviscosity, and cardiovascular risk factors. Thrombosis may occur in the absence of known risk factors. (See WARNING and PRECAUTIONS [5.5], PATIENT COUNSELING INFORMATION [17])**
- **Renal dysfunction, acute renal failure, osmotic nephrosis, and death may occur in predisposed patients who receive IGIV products, including OCTAGAM 10%. Patients predisposed to renal dysfunction include those with a degree of pre-existing renal insufficiency, diabetes mellitus, age greater than 65, volume depletion, sepsis, paraproteinemia, or patients receiving known nephrotoxic drugs. Renal dysfunction and acute renal failure occur more commonly in patients receiving IGIV product containing sucrose. OCTAGAM 10% does not contain sucrose.**
- **For patients at risk of thrombosis, renal dysfunction or acute renal failure, administer OCTAGAM 10% at the minimum dose and infusion rate practicable. Ensure adequate hydration in patients before administration. Monitor for signs and symptoms of thrombosis and assess blood viscosity in patients at risk for hyperviscosity. (See DOSAGE and ADMINISTRATION [2.3], WARNINGS and PRECAUTIONS [5.5])**

1 INDICATIONS AND USAGE

OCTAGAM 10% is indicated in Chronic Immune Thrombocytopenic Purpura to rapidly raise platelet counts to control or prevent bleeding in adults.

2 DOSAGE AND ADMINISTRATION

For intravenous use only.

2.1 Dosage

Administer OCTAGAM 10% at a total dose of 2 g/kg, divided into two doses of 1 g/kg (10mL/kg) given on two consecutive days.

2.2 Preparation and handling

- Parenteral products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Do not use if OCTAGAM 10% is turbid and/or if discoloration is observed.
- Do not mix OCTAGAM 10% with other medicinal products or administer simultaneously with other intravenous preparation in the same infusion set. Do not mix OCTAGAM 10% with immune globulin intravenous (IGIV) products from other manufacturers.
- Do not freeze OCTAGAM 10%. Do not use solutions that have been frozen.
- OCTAGAM 10% contains no preservative. OCTAGAM 10% bottle is for single use only. Use promptly any bottle that has been entered or opened, and discard partially used bottles.
- The content of OCTAGAM 10% bottles may be pooled under aseptic conditions into sterile infusion bags and infused within 8 hours after pooling.
- Do not use after expiration date.
- Do not dilute OCTAGAM 10%.
- The infusion line may be flushed before and after administration of OCTAGAM 10% with either normal saline or 5% dextrose in water.

2.3 Administration

- Administer OCTAGAM 10%, which is to be at room temperature, only by the intravenous route.
- OCTAGAM 10% is not supplied with an infusion set. If an in-line filter is used the pore size should be 0.2 – 200 microns.
- Do not use a needle of larger than 16 gauge to prevent the possibility of coring. Insert needle only once, within the stopper area delineated (by the raised ring for penetration). Penetrate the stopper perpendicular to its plane and within the ring.

For rate of administration, see Table 1.

Table 1

Rate of Administration	mg/kg/min (mg/kg/hour)	mL/kg/min
first 30 min	1 (60)	0.01
next 30 min, if above is tolerated	2 (120)	0.02
next 30 min, if above is tolerated	4 (240)	0.04
next 30 min, if above is tolerated	8 (480)	0.08
Maximum	≤ 12 (≤720)	≤0.12

- Monitor the patient carefully throughout the infusion. Certain adverse drug reactions may be related to the rate of infusion. Slowing or stopping the infusion usually allows the symptoms to disappear promptly. Once the symptoms subside, the infusion may then be resumed at a lower rate.
- Ensure that patients with pre-existing renal insufficiency are not volume depleted. For patients at risk of renal dysfunction or thromboembolic events, administer OCTAGAM 10% at the minimum infusion rate practicable, not to exceed 3.3 mg/kg/min (0.03 mL/kg/min). Discontinue OCTAGAM 10% if renal function deteriorates.

3 DOSAGE FORMS AND STRENGTHS

Solution containing 10% IgG (0.1 g/L) (See How Supplied/Storage and Handling (16)).

4 CONTRAINDICATIONS

- OCTAGAM 10% is contraindicated in patients who have a history of severe systemic hypersensitivity reactions, such as anaphylaxis, to human immunoglobulin.
- OCTAGAM 10% contains trace amounts of IgA (average 106 µg/mL in a 10% solution). It is contraindicated in IgA-deficient patients with antibodies against IgA and history of hypersensitivity.

5 WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity

Severe hypersensitivity reactions may occur (*See Contraindications (4)*).^[1] In case of hypersensitivity, discontinue OCTAGAM 10% infusion immediately and institute appropriate treatment. Epinephrine should be immediately available for treatment of severe acute hypersensitivity reactions.

OCTAGAM 10% contains trace amounts of IgA (average 106 µg/mL in a 10% solution). IgA-deficient patients with antibodies against IgA are at greater risk of developing severe hypersensitivity and anaphylactoid reactions when administered OCTAGAM 10% (*See Contraindications (4)*).

OCTAGAM 10% may cause hypersensitivity reactions in patients with corn allergy. OCTAGAM 10% contains maltose, a disaccharide sugar which is derived from corn.

5.2 Renal Dysfunction/Failure

Renal dysfunction, acute renal failure, osmotic nephropathy, and death may occur upon use of OCTAGAM 10% in predisposed patients.

Ensure that patients are not volume depleted prior to the initiation of the infusion of OCTAGAM 10%.

For patients at risk of renal dysfunction because of pre-existing renal insufficiency or predisposition to acute renal failure (such as individuals with diabetes mellitus, age greater than 65, volume depletion, sepsis, paraproteinemia, or those receiving known nephrotoxic drugs), administer OCTAGAM 10% at the minimum infusion rate practicable, not to exceed 3.3mg/kg/min (0.03mL/kg/min) (*See Boxed Warning, and Dosage and Administration (2)*).

Periodic monitoring of renal function tests and urine output is particularly important in patients judged to be at risk of developing acute renal failure. Assess renal function, including a measurement of blood urea nitrogen (BUN)/serum creatinine, prior to the initial infusion of OCTAGAM 10% and again at appropriate intervals thereafter. If renal function deteriorates, consider discontinuation of the product.

5.3 Blood Glucose Monitoring

Some glucose monitoring systems give falsely elevated glucose readings because they interpret the maltose in OCTAGAM 10% as glucose.^[2] This occurs in systems that use glucose dehydrogenase pyrroloquinoline quinone (GDH-PQQ) or glucose-dye-oxidoreductase methods for detecting blood glucose. Inaccurate glucose readings can cause inappropriate administration of insulin and resultant potentially life-threatening hypoglycemia. Also, true hypoglycemia may go undetected when masked by a falsely elevated glucose reading.

Monitor glucose levels in diabetic patients with a glucose-specific method only. Review the product information of blood glucose testing systems, including the test strips, to determine if the system is appropriate to use with maltose-containing parenteral products. Contact the manufacturer if there is any uncertainty about the system.

5.4 Hyperproteinemia, Increased Serum Viscosity and Hyponatremia

Hyperproteinemia, increased serum viscosity and hyponatremia may occur in patients receiving OCTAGAM 10% therapy. It is clinically critical to distinguish true hyponatremia from pseudohyponatremia related to hyperproteinemia with concomitant decreased calculated serum osmolality or elevated osmolar gap, because treatment aimed at decreasing serum free water in patients with pseudohyponatremia may lead to volume depletion, a further increase in serum viscosity and a higher risk of thromboembolic events.^[3]

5.5 Thrombosis

Thrombosis may occur following treatment with immune globulin products, including OCTAGAM 10%. Risk factors may include: advanced age, prolonged immobilization, hypercoagulable conditions, history of venous or arterial thrombosis, use of estrogens, indwelling central vascular catheters, hyperviscosity, and cardiovascular risk factors. Thrombosis may occur in the absence of known risk factors.

Consider baseline assessment of blood viscosity in patients at risk for hyperviscosity, including those with cryoglobulins, fasting chylomicronemia / markedly high triacylglycerols (triglycerides), or monoclonal gammopathies. For patients at risk of thrombotic events, administer OCTAGAM 10% at the minimum dose and infusion rate practicable. Ensure adequate hydration in patients before administration. Monitor for signs and symptoms of thrombosis and assess blood viscosity in patients at risk for hyperviscosity. ^[4-6] (*See BOXED WARNING, DOSAGE and ADMINISTRATION [2.3], PATIENT COUNSELING INFORMATION [17]*)

5.6 Hemolysis

OCTAGAM 10% may contain blood group antibodies that can act as hemolysins and induce *in vivo* coating of red blood cells (RBCs) with immunoglobulin, causing a positive direct antiglobulin test (DAT) (Coombs' test) result and hemolysis. Delayed hemolytic anemia can develop subsequent to IGIV therapy due to enhanced RBC sequestration and acute hemolysis, consistent with intravascular hemolysis, has been reported. Cases of severe hemolysis-related renal dysfunction/failure or disseminated intravascular coagulation have occurred following infusion of IGIV.

The following risk factors may be associated with the development of hemolysis following IGIV administration: high doses (e.g. ≥ 2 g/kg), given either as a single administration or divided over several days, and non-O blood group.^[7] Other individual patient factors, such as an underlying inflammatory state (as may be reflected by, for example, elevated C-reactive protein or erythrocyte sedimentation rate), have been hypothesized to increase the risk of hemolysis following administration of IGIV.^[8], but their role is uncertain. Hemolysis has been reported following administration of IGIV for a variety of indications, including ITP.

Closely monitor patients for clinical signs and symptoms of hemolysis, particularly patients with risk factors noted above. Consider appropriate laboratory testing in higher risk patients, including measurement of hemoglobin or hematocrit prior to infusion and within approximately 36 to 96 hours post infusion. If clinical signs and symptoms of hemolysis or a significant drop in hemoglobin or hematocrit have been observed, perform confirmatory laboratory testing. If transfusion is indicated for patients who develop hemolysis with clinically compromising anemia after receiving IGIV, perform adequate cross-matching to avoid exacerbating on-going hemolysis.

5.7 Aseptic meningitis syndrome

Aseptic meningitis syndrome (AMS) may occur with OCTAGAM treatment. Discontinuation of treatment has resulted in remission of AMS within several days without sequelae. The syndrome usually begins within several hours to two days following rapid infusion with OCTAGAM treatment. It is characterized by symptoms and signs including severe headache, nuchal rigidity, drowsiness, fever, photophobia, painful eye movements, nausea and vomiting. Cerebrospinal fluid (CSF) studies frequently reveal pleocytosis up to several thousand cells per cubic millimeter, predominantly from the granulocytic series, and elevated protein levels up to several hundred mg/dl, but negative culture results. Conduct a thorough neurological examination in patients exhibiting such symptoms and signs, including CSF studies, to rule out other causes of meningitis. It appears that patients with a history of migraine may be more susceptible.^[9]

AMS may occur more frequently following high doses (≥ 2 g/kg) and/or rapid infusion of IGIV.

5.8 Transfusion-Related Acute Lung Injury (TRALI)

Noncardiogenic pulmonary edema [Transfusion-Related Acute Lung Injury (TRALI)] may occur in patients administered IGIV. [10] TRALI is characterized by severe respiratory distress, pulmonary edema, hypoxemia, normal left ventricular function, and fever. Signs and symptoms typically appear within 1 to 6 hours after transfusion. Patients with TRALI may be managed using oxygen therapy with adequate ventilatory support.

Monitor recipients of OCTAGAM 10% for pulmonary adverse reactions. If TRALI is suspected, perform appropriate tests for the presence of anti-HLA and anti-neutrophil antibodies in the product.

5.9 Transmission of Infectious Agents

Because OCTAGAM 10% is made from human blood, it may carry a risk of transmitting infectious agents, e.g., viruses and theoretically, the Creutzfeldt-Jakob disease (CJD) agent. Report all infections thought by a physician or other healthcare provider to have been possibly transmitted by this product to Octapharma at 1-866-766-486. Discuss the risks and benefits of OCTAGAM 10% with the patient before prescribing or administering this product.

5.10 Monitoring Laboratory Tests

- After infusion of immunoglobulin, the transitory rise of the various passively transferred antibodies in the patient's blood may yield positive serological testing results, with the potential for misleading interpretation.
- Passive transmission of antibodies to erythrocyte antigens (e.g. A, B, and D) may cause a positive direct or indirect antiglobulin (Coombs') test. Clinically assess patients with known renal dysfunction, diabetes mellitus, age greater than 65, volume depletion, sepsis, paraproteinemia, or those receiving nephrotic agents, and monitor as appropriate (BUN; serum creatinine, urine output) during therapy with OCTAGAM 10%.
- Consider baseline assessment of blood viscosity in patients at risk for hyperviscosity, including those with polycythemia, cryoglobulin, fasting chylomicronemia/markedly high triglycerides, or monoclonal gammopathies.
- Consider measuring hemoglobin or hematocrit at baseline and approximately 36 to 96 hours post infusion in patients at higher risk of hemolysis. If signs and/or symptoms of hemolysis are present after an infusion of OCTAGAM 10%, perform appropriate laboratory testing for confirmation.
- If TRALI is suspected, perform appropriate tests for the presence of anti-neutrophil antibodies in both the product and patient's serum.

6 ADVERSE REACTIONS

The most common adverse reactions observed with OCTAGAM 10% treatment during clinical trial were headache, fever, and increased heart rate.

The most serious adverse reaction observed with OCTAGAM 10% treatment during clinical trial was headache.

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a product cannot be directly compared to rates in the clinical trials of another product and may not reflect the rates observed in clinical practice.

In a prospective, open-label, single-arm, multicenter study 116 subjects with newly diagnosed or chronic ITP received OCTAGAM 10% at a dose of 2 g/kg, administered daily as 1 g/kg intravenous infusions on 2 consecutive days. In the initial study phase, the infusion rate allowed for OCTAGAM 10% administration was up to 6 mg/kg/min [0.06 mL/kg/min], which was achieved in 24 of 26 subjects (92%). Subsequently, the maximum infusion rate allowed was 12 mg/kg/min [0.12mL/kg/min], and 54 (60%) of 90 subjects received OCTAGAM 10% at this rate in the second phase of the trial.

Pre-medication to alleviate potential adverse drug reactions was allowed, and was only administered to 1 (0.9%) of the 116 patients.

The investigators judged 129 (54%) of the temporally associated ("infusional") adverse reactions (AR) in 62 patients to be at least possibly related to the infusion of OCTAGAM 10%. The most common causally related ARs observed in >5% of subjects are shown in Table 2.

Table 2: Drug-related Adverse Reactions Experienced by >5% of Subjects

Reactions	No. of Subjects (% of Subjects [n=116])
Headache	29 (25)
Pyrexia	17 (15)
Heart Rate Increased	13 (11)

Table 3 lists the ARs that occurred in more than 5% of subjects during and within 72 hours after the end of a treatment cycle (two consecutive infusions) with OCTAGAM 10%, irrespective of causality.

Table 3: ARs in >5% Subjects During and Within 72 Hours After the End of 2-Days Treatment Cycle, Irrespective of Causality

Reactions	No. of Subjects (% of Subjects [n=116])
Headache	35 (30)
Heart Rate Increased	25 (22)
Pyrexia	20 (17)
Heart Rate Decreased	18 (16)
Hypertension	7 (6)

One subject experienced a serious adverse reaction (headache).

6.2 Postmarketing Experience

Because postmarketing reporting of adverse reactions is voluntary and from a population of uncertain size, it is not always possible to reliably estimate the frequency of these reactions or establish a causal relationship to product exposure.

OCTAGAM 5% Postmarketing Experience

The following adverse reactions have been identified during post-approval use of OCTAGAM 5%. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or to establish a causal relationship to OCTAGAM 5%.

Blood and lymphatic system disorders Leucopenia, hemolytic anemia
Immune system disorders Hypersensitivity, anaphylactic shock, anaphylactic reaction, anaphylactoid reaction, angioedema, face edema
Metabolic and nutritional disorders Fluid overload
Psychiatric disorders Agitation
Nervous system disorders Headache, cerebrovascular accident, aseptic meningitis, migraine, dizziness, paraesthesia
Cardiac disorders Myocardial infarction, tachycardia, palpitations, cyanosis
Vascular disorders Hypotension, thrombosis, peripheral circulatory failure, hypertension
Respiratory, thoracic and mediastinal disorders Respiratory failure, pulmonary embolism, pulmonary edema, bronchospasm, dyspnea, cough
Gastrointestinal disorders Nausea, vomiting, diarrhea, abdominal pain
Skin and subcutaneous tissue disorders Eczema, urticaria, rash, dermatitis, pruritus, alopecia
Musculoskeletal and connective tissue disorders Back pain, arthralgia, myalgia, pain in extremity
Renal and urinary disorders Acute renal failure
General disorders and administration site conditions Fatigue, injection site reaction, pyrexia, chills, chest pain, hot flush, flushing, hyperhidrosis, malaise
Investigations Hepatic enzymes increased, falsely elevated blood glucose level

Postmarketing Experience with IGIV Products

The following adverse reactions have been identified during post-approval use of IGIV products. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or to establish a causal relationship to IGIV products:

Infusion reactions: Hypersensitivity (e.g., anaphylaxis), headache, diarrhea, tachycardia, fever, fatigue, dizziness, malaise, chills, flushing, urticaria or other skin reactions, wheezing or other chest discomfort, nausea, vomiting, rigors, back pain, myalgia, arthralgia, and changes in blood pressure
Renal Acute renal dysfunction/failure, osmotic nephropathy
Respiratory Apnea, Acute Respiratory Distress Syndrome (ARDS), TRALI, cyanosis, hypoxemia, pulmonary edema, dyspnea, bronchospasm
Cardiovascular Cardiac arrest, thromboembolism, vascular collapse, hypotension
Neurological Coma, loss of consciousness, seizures, tremor, aseptic meningitis syndrome
Integumentary Stevens-Johnson syndrome, epidermolysis, erythema multiforme, dermatitis (e.g., bullous dermatitis)
Hematologic Pancytopenia, leukopenia, hemolysis, positive direct antiglobulin (Coombs') test
Gastrointestinal Hepatic dysfunction, abdominal pain
General/Body as a Whole Pyrexia, rigors

7 DRUG INTERACTIONS

Admixtures of OCTAGAM 10% with other drugs and intravenous solutions have not been evaluated. It is recommended that OCTAGAM 10% be administered separately from other drugs or medications which the patient may be receiving. Do not mix the product.

Do not mix OCTAGAM 10% with IGIVs from other manufacturers.

Passively transferred antibodies in immunoglobulin preparations can confound the results of serological testing.

Antibodies in OCTAGAM 10% may interfere with the response to live viral vaccines, such as measles, mumps, and rubella. Inform physicians of recent therapy with OCTAGAM 10%, so that administration of live viral vaccines, if indicated, can be appropriately delayed for 3 or more months from the time of OCTAGAM 10% administration.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C. Animal reproduction studies have not been conducted with OCTAGAM 10%. It is also not known whether OCTAGAM 10% can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. OCTAGAM 10% should be given to a pregnant woman only if clearly needed.

8.3 Nursing Mothers

It is not known whether OCTAGAM 10% is excreted in human milk. Because many drugs are excreted in human milk caution should be exercised when OCTAGAM 10% is administered to a nursing woman.

8.4 Pediatric Use

The safety and effectiveness of OCTAGAM 10% has not been established in pediatric patients with ITP.

8.5 Geriatric Use

Patients > 65 years of age may be at increased risk for developing certain adverse reactions such as thromboembolic events and acute renal failure (*See Boxed Warnings and Thrombotic Events (5.5) and Renal Failure (5.2)*). Do not exceed recommended doses in this population, and the applied infusion rate should be the minimum practicable. Clinical studies of OCTAGAM 10% did not include sufficient numbers of subjects > 65 years to determine whether they respond differently from younger subjects.

10 OVERDOSAGE

With intravenous administration, overdose of OCTAGAM 10% may lead to fluid overload and hyperviscosity. Patients at risk of complications of fluid overload and hyperviscosity include elderly patients and those with cardiac or renal impairment.

11 DESCRIPTION

Immune Globulin Intravenous (Human), OCTAGAM 10%, is a solvent/detergent (S/D)-treated, sterile preparation of highly purified immunoglobulin G (IgG) derived from large pools of human plasma. OCTAGAM 10% is a solution for infusion to be administered intravenously.

This preparation contains approximately 100 mg of protein per mL (10%) of which not less than 96% is normal human immunoglobulin G. OCTAGAM 10% contains not more than 3% aggregates, not less than 94% monomers and dimers and not more than 3% fragments. On average, the product contains 106 µg/mL of IgA and even lower amounts of IgM.

The sodium content of the final solution is not more than 30 mmol/L and the pH is between 4.5 and 5.0. The osmolality is 310 - 380 mosmol/kg.

The manufacturing process for OCTAGAM 10% isolates IgG without additional chemical or enzymatic modification, and the Fc portion is maintained intact. OCTAGAM 10% contains the IgG antibody activities present in the donor population. IgG subclasses are fully represented with the following approximate percents of total IgG: IgG₁ is 65%, IgG₂ is 30%, IgG₃ is 3% and IgG₄ is 2%.

OCTAGAM 10% contains a broad spectrum of IgG antibodies against bacterial and viral agents that are capable of opsonization and neutralization of microbes and toxins. It contains maltose (90 mg/mL), but no preservatives or sucrose.

All units of human plasma used in the manufacture of OCTAGAM 10% are provided by FDA-approved blood establishments, and are tested by FDA-licensed serological tests for HBsAg, antibodies to HCV and HIV and Nucleic Acid Test (NAT) for HCV and HIV-1 and found to be non-reactive (negative).

The product is manufactured by the cold ethanol fractionation process followed by ultrafiltration and chromatography. The manufacturing process includes treatment with an organic S/D mixture composed of tri-n-butyl phosphate (TNBP) and Triton X-100 (Octoxynol). The OCTAGAM 10% manufacturing process shows significant viral reduction in in vitro studies (Table 4). These reductions are achieved through a combination of process steps including cold ethanol fractionation, S/D treatment and pH 4 treatment.

Table 4: In vitro reduction factor during OCTAGAM 10% manufacturing

Production step	In vitro reduction factor [\log_{10}]				
	Enveloped viruses			Non-enveloped viruses	
	HIV-1	PRV	SBV	MEV	PPV
Cold ethanol fractionation	≥ 4.81	≥ 6.28	≥ 7.13	≥ 7.13	≥ 6.53
S/D treatment	≥ 4.93	5.23	≥ 6.77	Not applicable	
pH4 treatment	≥ 4.33	≥ 6.71	6.71	5.07	< 1*
Global reduction factor	≥ 14.07	≥ 18.22	≥ 20.61	≥ 12.20	≥ 6.53

*not calculated for global LRF

HIV-1: Human Immunodeficiency Virus - 1

PRV: Pseudorabies Virus

SBV: Sindbis Virus

MEV: Mouse Encephalomyelitis Virus

PPV: Porcine Parvovirus

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The mechanism of action of immunoglobulins in the treatment of *chronic* ITP has not been fully elucidated.

12.3 Pharmacokinetics

Pharmacokinetic studies with OCTAGAM 10% have not been performed in patients with *chronic* ITP.

13 NON-CLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No studies were conducted on carcinogenesis, mutagenesis, or impairment of fertility with OCTAGAM 10%.

Repeated-dose toxicity studies and genotoxic studies gave no evidence of carcinogenic properties of TNBP and Octoxynol.[11,12].

The results of *in vitro* and *in vivo* genotoxicity studies for TNBP and Octoxynol were negative. The results of studies on the embryotoxic and teratogenic properties of TNBP and Octoxynol in rats and rabbits at a wide range of i.v. doses were also negative.

13.2 Animal Toxicology and/or Pharmacology

No studies were conducted on pharmacology and pharmacokinetics with OCTAGAM 10% in animals.

A variety of single-dose toxicity studies were performed for TNBP and Octoxynol alone or in combination: the lowest toxic dose of TNBP + Octoxynol (in a ratio of 1:5) was 10,000 mcg/kg BW in rats after intravenous administration. Studies on 13-week toxicity were performed for intravenous combinations of TNBP + Octoxynol in a broad dose range in dogs and rats: the lowest toxic dose for rats was local 60 mcg TNBP/kg + 300 mcg Octoxynol/kg BW (concentration: 0.0006% and 0.003%, respectively) and systemic 300 mcg TNBP/kg + 1,500 mcg Octoxynol/kg BW (concentration: 0.003% and 0.015%, respectively); the lowest toxic dose for dogs was local 50 mcg TNBP/kg + 250 mcg Octoxynol/kg BW (concentration: 0.005% and 0.025%, respectively) and systemic 500 mcg TNBP/kg + 2,500 mcg Octoxynol/kg BW (concentration: 0.05% and 0.25%, respectively).

A pharmacokinetic study was carried out in rats given 300 mcg of TNBP/kg and 1,500 mcg Octoxynol/kg BW i.v. The plasma half-life for TNBP was approximately 20 minutes. Octoxynol was not detected.

14 CLINICAL STUDIES

A prospective, open-label, single-arm, multicenter study assessed the efficacy, safety, and tolerability of OCTAGAM 10% in 116 subjects with tentative newly diagnosed or *chronic* ITP and a platelet count of $20 \times 10^9/L$ or less. Subjects ranged in age from 17 to 88; 64% were female and 36% were male.

Subjects received a 2 g/kg dose of OCTAGAM 10% administered as two daily 1 g/kg intravenous doses, given on 2 consecutive days. Pre-medication to alleviate potential adverse drug reactions was allowed but only given to 1 (0.9%) of 116 patients. In the initial study phase, the infusion rate allowed for OCTAGAM 10% administration was up to 6 mg/kg/min [0.06 mL/kg/min], which was achieved in 24 of 26 subjects (92%). Subsequently, the maximum infusion rate allowed was 12 mg/kg/min [0.12mL/kg/min], which was achieved in 54 of 90 subjects (60%). Platelet counts were measured on Days 1 to 7, 14, 21, and 63.

The study was designed to determine the response rate defined as the percentage of subjects with an increase in platelet count to at least $50 \times 10^9/L$ within 7 days after the first infusion (responders). Additionally, maximum platelet count, the time to reach a platelet count of at least $50 \times 10^9/L$ within the first 7 days, the duration of that response (i.e. the number of days the platelet count remained in excess of $50 \times 10^9/L$), and the regression of hemorrhages in subjects who had bleeding at baseline were observed.

Of the 66 subjects with chronic ITP in the efficacy analysis, 54 (82%) responded to OCTAGAM 10% with a rise in platelet counts to at least $50 \times 10^9/L$ within 7 days after the first infusion. The lower bound of the overall 95% confidence interval for the response rate (73%) is above the predefined response rate of 70%.

The mean maximum platelet count achieved in the 66 subjects with chronic ITP was $227 \times 10^9/L$.

Table 5. Maximum Platelet Count ($\times 10^9/L$)

	Chronic ITP subjects (n=66)
Median and range	212.0 (18 to 686)
Mean \pm s.d.	227.0 \pm 155.63

The median time to reach a platelet response of at least $50 \times 10^9/L$ was 2 days after the first infusion. The duration of platelet response was analyzed for the 54 subjects with chronic ITP who achieved a response within 7 days after the first infusion: the median duration of platelet response in these subjects was 12 days (range: 1 to 79 days).

Table 6. Time to and Duration of Platelet Response (Responders Only).

	Time to platelet response (days)	Duration of platelet response (days)
	Chronic ITP subjects (n=54)	Chronic ITP subjects (n=54)
Median and range	2 (1 to 5)	12 (1 to 79)
Mean \pm s.d.	1.8 \pm 0.91	20.9 \pm 21.76

In 35 of the 45 subjects with chronic ITP (78%) who had bleeding at baseline, the hemorrhages had completely resolved by Day 7. A decrease in the severity of hemorrhage from baseline to Day 7 was observed in 38 of 45 subjects (84%) with chronic ITP.

Table 7. Assessment of Bleeding at Day 7 in Subjects with Baseline Bleeding

Number of Subjects with Baseline Bleeding	Severity at Day 7				
	None	Minor	Mild	Moderate	Missing
Chronic ITP (N=45)	35 (78%)	8 (18%)	0	0	2 (4%)

15 REFERENCES

- Duhem C, Dicato MA, Ries F: Side-effects of intravenous immune globulins. Clin.Exp.Immunol. 1994;97 Suppl 1:79-83.
- Kannan S, Rowland CH, Hockings GI, et al: Intragam can interfere with blood glucose monitoring. Med J Aust. 2004;180:251-252.
- Steinberger BA, Ford SM, Coleman TA: Intravenous immunoglobulin therapy results in post-infusional hyperproteinemia, increased serum viscosity, and pseudohyponatremia. Am.J Hematol. 2003;73:97-100.
- Dalakas MC: High-dose intravenous immunoglobulin and serum viscosity: risk of precipitating thromboembolic events. Neurology 1994;44:223-226.
- Go RS, Call TG: Deep venous thrombosis of the arm after intravenous immunoglobulin infusion: case report and literature review of intravenous immunoglobulin-related thrombotic complications. Mayo Clin Proc 2000;75:83-85.
- Wolberg AS, Kon RH, Monroe DM, et al: Coagulation factor XI is a contaminant in intravenous immunoglobulin preparations. Am.J.Hematol. 2000;65:30-34.
- Kahwaji J, Barker E, Pepkowitz S, et al: Acute Hemolysis After High-Dose Intravenous Immunoglobulin Therapy in Highly HLA Sensitized Patients. Clinical journal of the American Society of Nephrology : CJASN 2009;DOI:10.2215/CJN.04540709.
- Daw Z, Padmore R, Neurath D, et al: Hemolytic transfusion reactions after administration of intravenous immune (gamma) globulin: a case series analysis. Transfusion 2008;48:1598-1601.
- Sekul EA, Cupler EJ, Dalakas MC: Aseptic meningitis associated with high-dose intravenous immunoglobulin therapy: frequency and risk factors. Ann Intern.Med 1994;121:259-262.
- Rizk A, Gorson KC, Kenney L, et al: Transfusion-related acute lung injury after the infusion of IVIG. Transfusion 2001;41:264-268.
- Auletta CS, Kotkoskie LA, Saulog T, et al: A dietary oncogenicity study of tributyl phosphate in the CD-1 mouse. Toxicology 1998;128:135-141.
- Auletta CS, Weiner ML, Richter WR: A dietary toxicity/oncogenicity study of tributyl phosphate in the rat. Toxicology 1998;128:125-134

16 HOW SUPPLIED/STORAGE AND HANDLING

OCTAGAM 10% is supplied in 2 g, 5 g, 10 g or 20 g single use bottles.

NDC Number	Size	Grams Protein
Octapharma AB		
68982 – 850 – 01	20 ml	2.0
68982 – 850 – 02	50 ml	5.0
68982 – 850 – 03	100 ml	10.0
68982 – 850 – 04	200 ml	20.0

OCTAGAM 10% is not supplied with an infusion set. If a filtered infusion set is used (not mandatory), choose a filter size of 0.2 – 200 microns.

Components used in the packaging of OCTAGAM 10% are latex-free.

Store OCTAGAM 10% for 24 months at +2°C to +8°C (36°F to 46°F) from the date of manufacture. Within the first 12 months of this shelf-life, the product may be stored up to 6 months at ≤ +25°C (77°F). After storage at ≤ +25°C (77°F) the product must be used or discarded.

Do not use after expiration date.

Do not freeze. Do not use frozen product.

Dispose of any unused product or waste material in accordance with local requirements.

17 PATIENT COUNSELING INFORMATION

Inform patients of the signs and symptoms of hypersensitivity reactions including urticaria, generalized urticaria, tightness of the chest, wheezing, hypotension, and anaphylaxis, and to contact their physicians immediately if allergic symptoms occur,

Inform patients to immediately report the signs and symptoms of the following conditions to their physician:

- renal failure, such as decreased urine output, sudden weight gain, fluid retention/edema, and/or shortness of breath
- aseptic meningitis, such as headache, neck stiffness, drowsiness, fever, sensitivity to light, painful eye movements, nausea, and vomiting
- hemolysis, such as fatigue, increased heart rate, yellowing of the skin or eyes, and dark-colored urine
- TRALI, such as troubled breathing, chest pain, blue lips or extremities, and fever. TRALI typically occurs within 1 to 6 hours following transfusion

Thrombosis

Instruct patients to immediately report symptoms of thrombosis. These symptoms may include: pain and/or swelling of an arm or leg with warmth over the affected area, discoloration of an arm or leg, unexplained shortness of breath, chest pain or discomfort that worsens on deep breathing, unexplained rapid pulse, numbness or weakness on one side of the body.

Inform patients that OCTAGAM 10% is made from human plasma and may contain infectious agents that can cause disease (e.g., viruses, and, theoretically, the CJD agent), and that the risk of infectious agent transmission has been reduced by (a) screening plasma donors for prior exposure to certain viruses, (b) testing the donated plasma for certain viral infections and (c) inactivating and/or removing certain viruses during manufacture.

Inform patients that administration of OCTAGAM 10% may interfere with the response to live viral vaccines such as measles, mumps and rubella, and to notify their immunizing physician of their therapy with OCTAGAM 10%.

Octapharma USA Inc.
Hoboken, NJ 07030