

N-095735-31-2090310
Monensin Med. Feedlot Cattle Feed - Cocci/FE
Dry Type B Med Feed
Page 1 of 2

Monensin Medicated Feedlot Cattle Feed

Type B Medicated Feed

For Cattle Fed in Confinement for Slaughter

For prevention and control of coccidiosis due to *Eimeria bovis* and *Eimeria zuernii*.

For improved feed efficiency.

Active Drug Ingredient

Monensin, USP41 to 80,000 g/ton*

Guaranteed Analysis

Crude Protein, not less than.....%
Non-Protein Nitrogen (NPN)¹, not more than.....%
Crude Fat, not less than.....%
Crude Fiber, not more than.....%
Calcium, not less than.....%
Calcium, not more than.....%
Phosphorus, not less than.....%
Salt², not less than.....%
Salt², not more than.....%
Sodium³, not less than.....%
Sodium³, not more than.....%
Potassium, not less than.....%
Vitamin A^{2,4}, not less than..... I.U./lb

¹When added.

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

Ingredients

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

*Final printed label on formulated Type B medicated feed must bear a single drug concentration.

N-095 735-4-0309 0310
Monensin Med. Feedlot Cattle Feed - Cocci
Dry Type B Med. Feed.
Page 2 of 2

Mixing Directions

Thoroughly mix monensin Type B Medicated Feed into the appropriate amount of non-medicated feed to obtain the correct concentration in the Type C Medicated Feed, 5 to 40 g/ton monensin for feed efficiency, or 10 to 40 g/ton monensin for prevention and control of coccidiosis. No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 g/ton (360 mg monensin per head per day).

Use only the portion of the table below that is applicable to the concentration of monensin in the Type C Medicated Feed you manufacture.

(Example: 600 g/ton Type B Medicated Feed)

Monensin Level Desired in Type C Medicated Feed	Amount of Type B Medicated Feed to make a ton of Type C Medicated Feed	Amount of Non- Medicated Feed
g/ton	lb/ton	lb/ton
5	16.7	1983.3
10	33	1967
20	67	1933
30 [†]	100	1900
40 [†]	133	1867

[†] No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 g/ton (360 mg monensin per head per day).

Caution

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Manufactured By
Blue Bird Feed Mill
Any town, USA 12345

Net Weight lb on Bag or Bulk