

N-095735-U-~~0309~~⁰³¹⁰
 Monensin Med. Feedlot Cattle Feed - cocci
 Type C Med. Feed
 Page 1 of 2

Monensin Medicated Feedlot Cattle Feed

Type C Medicated Feed

**For Cattle Fed in Confinement for Slaughter
 (For prevention and control of coccidiosis)**

For prevention and control of coccidiosis due to *Eimeria bovis* and *Eimeria zuernii*.

Active Drug Ingredient

Monensin, USP10 to 40 g/ton*

Guaranteed Analysis

Crude Protein, not less than	_____	%
Non-Protein Nitrogen (NPN) ¹ , not more than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than	_____	%
Salt ² , not less than	_____	%
Salt ² , not more than	_____	%
Sodium ³ , not less than	_____	%
Sodium ³ , not more than	_____	%
Potassium, not less than	_____	%
Vitamin A ^{2,4} , not less than	_____	I.U./lb

¹When added

²If added

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

Ingredients

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

Feeding Directions

Feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending upon the severity of challenge, up to maximum of 480 milligrams per head per day.

* Final printed label on formulated Type C medicated feed must bear a single drug concentration

N-095735-u ~~0310~~

monensin Med. Feedlot Cattle Feed - cocci

Type C Med. Feed

page 2 of 2

Caution

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning

A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

Manufactured By
Blue Bird Feed Mill
Anytown, USA 12345

Net Weight lb on bag or bulk