

**Oxytetracycline Chicken Ration for Control of Infectious Synovitis and Control of Fowl Cholera  
Type C Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

For control of infectious synovitis caused by Mycoplasma synoviae and control of fowl cholera caused by Pasteurella multocida susceptible to oxytetracycline.

**ACTIVE DRUG INGREDIENT**

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to \*100-200 grams oxytetracycline hydrochloride/ton.

**GUARANTEED ANALYSIS**

|                                      |         |
|--------------------------------------|---------|
| Crude Protein, (Min) . . . . .       | XX.XX % |
| Lysine, (Min) . . . . .              | X.XX%   |
| Methionine, (Min) . . . . .          | X.XX%   |
| Crude Fat, (Min) . . . . .           | X.XX%   |
| Crude Fiber, (Max) . . . . .         | X.XX%   |
| Calcium, (Min) . . . . .             | X.XX%   |
| Calcium, (Max) . . . . .             | X.XX%   |
| Phosphorus, (Min) . . . . .          | X.XX%   |
| Salt, (Min) <sup>1</sup> . . . . .   | X.XX%   |
| Salt, (Max) <sup>1</sup> . . . . .   | X.XX%   |
| Sodium, (Min) <sup>2</sup> . . . . . | X.XX%   |
| Sodium, (Max) <sup>2</sup> . . . . . | X.XX%   |

<sup>1</sup>If added

<sup>2</sup>Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

**INGREDIENTS**

Ingredients as defined by AAFCO

**DIRECTIONS FOR USE**

Feed continuously for 7-14 days as the sole ration.

**RESIDUE WARNING:** Do not feed to chickens producing eggs for human consumption. Zero-day withdrawal period. Do not use in low-calcium feed containing less than 0.55% dietary calcium. Use in such feeds may result in violative residues.

\*The medicated feed label must state a single drug concentration.

Net Weight \_\_\_\_\_ lb (\_\_\_\_\_ kg)

Manufactured by  
Blue Bird Feed Company

Blue Bird, MD 00000