

**Oxytetracycline Turkey Supplement
Type B Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated turkey feed.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to *2-20 grams oxytetracycline hydrochloride/lb.

GUARANTEED ANALYSIS

Crude Protein, (Min).....	XX.XX %
Lysine, (Min).....	X.XX%
Methionine, (Min).....	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min).....	X.XX%
Calcium, (Max).....	X.XX%
Phosphorus, (Min).....	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below. To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing as a complete feed.

Indications for Use

Oxytetracycline in
Type C medicated feed

TURKEYS

For Turkeys: Mix 50 pounds of Type B (2 g/lb) or 5 pounds of Type B (20 g/lb) with non-medicated feed ingredients to make a ton of turkey feed for control of hexamitiasis caused by Hexamita meleagridis susceptible to oxytetracycline containing 100 g/ton to be fed continuously for 7 to 14 days.

100 g/ton

RESIDUE WARNING: Do not feed to turkeys producing eggs for human consumption.

For Turkeys: Mix 100 pounds of Type B (2 g/lb) or 10 pounds of Type B (20 g/lb) with non-medicated feed ingredients to make a ton of turkey feed for control of infectious synovitis caused by Mycoplasma synoviae susceptible to oxytetracycline containing 200 g/ton to be fed continuously for 7 to 14 days. 200 g/ton

RESIDUE WARNING: Do not feed to turkeys producing eggs for human consumption.

Mix an amount of Type B feed to provide approximately 25 mg/lb body weight/day varying with body weight and feed consumption with non-medicated ingredients to make a ton of turkey feed for control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline to be fed continuously for 7 to 14 days. 25 mg/lb body weight/day

RESIDUE WARNING: Zero-day withdrawal period. Do not feed to turkeys producing eggs for human consumption.

*The medicated feed label must state a single drug concentration.

Net Weight _____ lb (_____ kg)
Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000