

**Oxytetracycline Turkey Ration for Control of Hexamitiasis  
Type C Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

For control of hexamitiasis caused by Hexamita meleagridis susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to 100 grams oxytetracycline hydrochloride/ton.

GUARANTEED ANALYSIS

Crude Protein, (Min).....	XX.XX %
Lysine, (Min).....	X.XX%
Methionine, (Min).....	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min).....	X.XX%
Calcium, (Max).....	X.XX%
Phosphorus, (Min).....	X.XX%
Salt, (Min) <sup>1</sup> .....	X.XX%
Salt, (Max) <sup>1</sup> .....	X.XX%
Sodium, (Min) <sup>2</sup> .....	X.XX%
Sodium, (Max) <sup>2</sup> .....	X.XX%

<sup>1</sup>If added

<sup>2</sup>Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO

DIRECTIONS FOR USE

Feed continuously for 7-14 days as the sole ration.

**RESIDUE WARNING: Do not feed to turkeys producing eggs for human consumption.  
Zero-day withdrawal period.**

Net Weight \_\_\_\_\_ lb (\_\_\_\_\_ kg)

Manufactured by  
Blue Bird Feed Company  
Blue Bird, MD 00000