

**Oxytetracycline Turkey Ration for Control of Infectious Synovitis
Type C Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of infectious synovitis caused by Mycoplasma synoviae susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to 200 grams oxytetracycline hydrochloride/ton.

GUARANTEED ANALYSIS

Crude Protein, (Min)	XX.XX %
Lysine, (Min)	X.XX%
Methionine, (Min)	X.XX%
Crude Fat, (Min)	X.XX%
Crude Fiber, (Max)	X.XX%
Calcium, (Min)	X.XX%
Calcium, (Max)	X.XX%
Phosphorus, (Min)	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO

DIRECTIONS FOR USE

Feed continuously for 7-14 days as the sole ration.

**RESIDUE WARNING: Do not feed to turkeys producing eggs for human consumption.
Zero-day withdrawal period.**

Net Weight _____ lb (_____ kg)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000