

**Oxytetracycline Turkey Ration for Control of Complicating Bacterial Organisms Associated with
Bluecomb
Type C Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to _____ grams* oxytetracycline hydrochloride/ton to provide approximately 25 mg/lb body weight/day varying with body weight and feed consumption.

GUARANTEED ANALYSIS

Crude Protein, (Min).	XX.XX %
Lysine, (Min).	X.XX%
Methionine, (Min).	X.XX%
Crude Fat, (Min).	X.XX%
Crude Fiber, (Max).	X.XX%
Calcium, (Min).	X.XX%
Calcium, (Max).	X.XX%
Phosphorus, (Min).	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO

DIRECTIONS FOR USE

Feed continuously for 7-14 days as the sole ration.

**RESIDUE WARNING: Do not feed to turkeys producing eggs for human consumption.
Zero-day withdrawal period.**

*The medicated feed label must state a single drug concentration.

Net Weight _____ lb (_____ kg)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000