

**Oxytetracycline Cattle Supplement
Type B Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated cattle feed.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to *2-20 grams oxytetracycline hydrochloride/lb.

GUARANTEED ANALYSIS

Crude Protein, (Min).....	XX.XX %
NPN, (Max) ¹	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min) ¹	X.XX%
Calcium, (Max) ¹	X.XX%
Phosphorus, (Min) ¹	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%
Potassium, (Min) ¹	X.XX%
Vitamin A, (Min) ³	___IU/Lb

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

³Other than precursors of Vitamin A, if added

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below. To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing as a complete feed.

Indications for Use

Oxytetracycline in
Type C medicated feed

For Growing Cattle (over 400 lbs): Mix an amount of Type B feed to provide approximately 75 mg/head/day varying with body weight and feed consumption with non-medicated feed ingredients to make the daily complete feed for the reduction of the incidence of liver abscesses to be fed continuously.

75 mg/head/day

For Cattle: Mix an amount of Type B feed to provide approximately 0.5-2.0 g/head/day varying with body weight and feed consumption with non-medicated feed ingredients to make the daily complete feed for the prevention and treatment of the early stages of the shipping fever complex. Feed 3-5 days before and after animal arrival in feedlots.

0.5-2.0 g/head/day

For Calves, Beef and Nonlactating Dairy Cattle: Mix an amount of Type B feed to provide approximately 10 mg/lb body weight/day varying with body weight and feed consumption with non-medicated feed ingredients to make the daily complete feed for treatment of bacterial enteritis caused by Escherichia coli and bacterial pneumonia (shipping fever complex) caused by Pasteurella multocida susceptible to oxytetracycline to be fed continuously for 7-14 days.

10 mg/lb body weight/day

RESIDUE WARNING: 5-day withdrawal period at 10 mg/lb dosage.

*The medicated feed label must state a single drug concentration.

Net Weight _____ lb (_____ kg)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000