

**Oxytetracycline Cattle Ration for Treatment of Bacterial Enteritis and Bacterial Pneumonia
Type C Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For treatment of bacterial enteritis caused by Escherichia coli and bacterial pneumonia (shipping fever complex) caused by Pasteurella multocida organisms susceptible to oxytetracycline in calves, beef and non-lactating dairy cattle.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to _____ grams* oxytetracycline hydrochloride/ton to provide approximately 10 mg/lb body weight/day varying with body weight and feed consumption.

GUARANTEED ANALYSIS

Crude Protein, (Min).....	XX.XX %
NPN, (Max) ¹	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min) ¹	X.XX%
Calcium, (Max) ¹	X.XX%
Phosphorus, (Min) ¹	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%
Potassium, (Min) ¹	X.XX%
Vitamin A, (Min) ³	____IU/Lb

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

³Other than precursors of Vitamin A, if added

INGREDIENTS

Ingredients as defined by AAFCO

DIRECTIONS FOR USE

Feed continuously for 7-14 days as the sole ration.

RESIDUE WARNING: 5-day withdrawal period at 10 mg/lb dosage.

*The medicated feed label must state a single drug concentration.

Net Weight _____ lb (_____ kg)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000