

Bloat Guard Type C Medicated Feed ~~Admixture~~
N-032704-5-0091

POLOXALENE
BLUE BIRD

TYPE C MEDICATED CATTLE FEED

For prevention of legume (alfalfa, clover) and wheat pasture bloat in cattle

ACTIVE DRUG INGREDIENT

Poloxalene 1-100 g/lb*

GUARANTEED ANALYSIS

Crude Protein (Min)	_____	%
(This includes not more than 2.9% equivalent crude protein from non-protein nitrogen)		
Crude Fat (Min)	_____	%
Crude Fiber (Max)	_____	%
Calcium (Min)	_____	%
Calcium (Max)	_____	%
Phosphorus (Min)	_____	%
Salt (Min)	_____	%
Salt (Max)	_____	%
Potassium (Min)	_____	%
Vitamin A (Min)	_____	%

Minimum Selenium in parts per million (ppm)

Minimum Vitamin A, other than precursors of Vitamin A, in International Units per pound (if added)

INGREDIENTS

Ingredients as defined by AAFCO

FEEDING DIRECTIONS

Feed to beef cattle on pasture. Feed the required amount (depending on concentration in the finished feed) to deliver 1 - 2 g of poloxalene per 100 lb of body weight. For example:

If the feed contains 1 g poloxalene/lb, feed 1 - 2 lb per 100 lb of body weight.

If the feed contains 10 g poloxalene/lb, feed 0.1 - 0.2 lb per 100 lb of body weight.

Provide plenty of fresh, clean water at all times.

The dosage of Bloat Guard is proportional to body weight and also depends upon the severity of the bloat-producing conditions. The normal dosage of Bloat Guard (1 g of poloxalene per 100 lb of body weight) is recommended for cattle under moderate bloat-producing conditions. For cattle under severe bloat-producing conditions, the normal dosage should be doubled (2 g of poloxalene per 100 lb of body weight).

Repeat the feeding of Bloat Guard when animals are exposed to bloat-producing conditions more than 12 hours from the last feeding of Bloat Guard, but do not exceed the double dosage of Bloat Guard in any 24-hour period. Animals should be fed the recommended amounts of Bloat Guard starting 2 or 3 days before they are exposed to bloat-producing conditions.

CAUTION: It is essential that each animal consumes the total recommended dosage of Bloat Guard daily for adequate protection.

WARNING: The normal life of this product is at least 24 months. However, when the product is subjected to extreme temperatures (38°C/100°F) for long periods of time (6 months), spontaneous combustion may occur. The product is not combustible unless it develops a strong, irritating odor; if this occurs, flush with water and discard immediately.

MANUFACTURED BY
BLUE BIRD FEED MILL
ANYTOWN, USA 12345

Net Weight on Bag or Bulk

* The finished feed should contain a specific concentration of poloxalene, and feeding directions for that concentration. A feed concentration up to 10 g poloxalene per lb will suffice under typical feeding conditions.