

N-095735-U-~~0209~~⁰²¹⁰

Monensin Medicated Growing Cattle (Pasture) Feed - WG/Cocci

Type B Med. Feed.

page 1 of 2

MONENSIN MEDICATED GROWING CATTLE FEED

TYPE B MEDICATED FEED MONENSIN

For Growing Cattle on Pasture or in Dry Lot

For increased rate of weight gain; For prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).

Active Drug Ingredient

Monensin, USP 401 to 80,000 g/ton*

Guaranteed Analysis

Crude Protein, not less than.....	_____	%
Non-Protein Nitrogen (NPN) ¹ , not more than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ² , not less than.....	_____	%
Salt ² , not more than.....	_____	%
Sodium ³ , not less than.....	_____	%
Sodium ³ , not more than.....	_____	%
Potassium, not less than.....	_____	%
Vitamin A ^{2,4} , not less than.....	_____	I.U./lb

¹When added.

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

Ingredients

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

*Final printed label on formulated Type B medicated feed must bear a single drug concentration

Monensin Med. Growing Cattle (Pasture) Feed - wG/cocci

Type B Med. Feed

page 2 of 2

Mixing Directions

Thoroughly mix monensin Type B medicated feed into the appropriate amount of non-medicated feed to obtain the correct concentration in the Type C medicated feed (25 to 400 g/ton). Use only the portion of the table below that is applicable to the concentration of monensin in the Type C medicated feed you manufacture.

Monensin Level Desired in Type C Medicated Feed	Monensin Level in Type B Medicated Feed	Amount of Type B Medicated Feed to make a ton of Type C Medicated Feed*	Amount of Non-Medicated Feed
g/ton (mg/lb)	g/ton	lb/ton	lb/ton
25 (12.5)	500	100	1900
400 (200)	500	1600	400
25 (12.5)	1000	50	1950
400 (200)	1000	800	1200
25 (12.5)	5000	10	1990
400 (200)	5000	160	1840

* Amount of Type B needed to produce the Type C medicated feed with desired level of monensin is as follows:

$$\frac{\text{Desired level of monensin in Type C medicated feed} \times 2000}{\text{monensin g/ton in Type B}}$$

Caution

Do not allow horses or other equines access to feeds containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Manufactured By
Blue Bird Feed Mill
Anytown, USA 12345

Net Weight lb on Bag or Bulk