

N-095-735-U-0310  
~~0309~~

Monensin Medicated Growing Cattle (Pasture) Feed - WF/Cocci

Growing cattle on pasture or dry lot

Type C Med. Feed

page 1 of 2

**Monensin Medicated Growing Cattle Feed**

**Type C Medicated Feed**

**For Growing Cattle on Pasture or in Dry Lot**

For increased rate of weight gain; For prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).

**Active Drug Ingredient**

Monensin, USP .....25 to 400 g/ton\*

**Guaranteed Analysis**

Crude Protein, not less than .....	_____	%
Non-Protein Nitrogen (NPN) <sup>1</sup> , not more than .....	_____	%
Crude Fat, not less than .....	_____	%
Crude Fiber, not more than .....	_____	%
Calcium, not less than .....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than .....	_____	%
Salt <sup>2</sup> , not less than .....	_____	%
Salt <sup>2</sup> , not more than .....	_____	%
Sodium <sup>3</sup> , not less than .....	_____	%
Sodium <sup>3</sup> , not more than .....	_____	%
Potassium, not less than .....	_____	%
Vitamin A <sup>2,4</sup> , not less than .....	_____	I.U./lb.

<sup>1</sup>When added

<sup>2</sup>If added.

<sup>3</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

<sup>4</sup>Other than precursors of Vitamin A.

**Ingredients**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

\* Final printed label on formulated Type C medicated feed must bear a single drug concentration.

N-095735-U-~~0204~~<sup>0210</sup>

Monensin Medicated Growing Cattle Feed - W&Cocci

Growing cattle on pasture or dry lot

Type C med. Feed.

Page 2 of 2

### Feeding Directions

Feed the appropriate amount of Type C Medicated Feed to cattle to provide 50 to 200 mg monensin/hd/day. During the first 5 days, cattle should receive no more than 100 mg monensin/day contained in not less than 1 lb. of feed.

**For increased rate of weight gain:** Feed at a rate of not less than 50 nor more than 200 mg monensin/hd/day in not less than 1 lb. of medicated feed;

-OR-

After the 5<sup>th</sup> day, feed at the rate of 400 mg monensin/hd/day every other day in not less than 2 lbs. of feed.

**For coccidiosis prevention and control:** Feed at the rate of 0.14 to 0.42 mg monensin/lb/body weight per day, depending upon severity of challenge, up to 200 mg monensin/hd/day.

Desired Intake of Monensin	Concentration of Monensin In Type C Medicated Feed	Recommended Feeding Level
mg/hd/day	g/ton (mg/lb)	(lbs/hd/day)
50	25 (12.5)	4
60	30 (15)	4
50	50 (25)	2
60	60 (30)	2
100	100 (50)	2
200	200 (100)	2
150	300 (150)	1
200*	400* (200)	1

\*Maximum Approved Level

### Caution

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

### Warning

A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

Manufactured By  
Blue Bird Feed Mill  
Anytown, USA 12345

Net Weight lb on bag or bulk