

**Oxytetracycline Swine Supplement  
Type B Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

For use in manufacturing medicated swine feed.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to \*2-20 grams oxytetracycline hydrochloride/lb.

GUARANTEED ANALYSIS

Crude Protein, (Min) . . . . .	XX.XX %
Lysine, (Min) . . . . .	X.XX%
Crude Fat, (Min) . . . . .	X.XX%
Crude Fiber, (Max) . . . . .	X.XX%
Calcium, (Min) . . . . .	X.XX%
Calcium, (Max) . . . . .	X.XX%
Phosphorus, (Min) . . . . .	X.XX%
Salt, (Min) <sup>1</sup> . . . . .	X.XX%
Salt, (Max) <sup>1</sup> . . . . .	X.XX%
Sodium, (Min) <sup>2</sup> . . . . .	X.XX%
Sodium, (Max) <sup>2</sup> . . . . .	X.XX%
Selenium, (Min) . . . . .	ppm
Zinc, (Min) . . . . .	ppm

<sup>1</sup>If added

<sup>2</sup>Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below. To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing as a complete feed.

Indications for Use

Oxytetracycline in  
Type C medicated feed

SWINE

For Swine: Mix an amount of Type B feed to provide approximately 400 g/ton varying with body weight and feed consumption to provide 10 mg/lb body weight/day with non-medicated feed ingredients to make a ton of swine feed for treatment of bacterial enteritis caused by Escherichia coli and Salmonella choleraesuis and bacterial pneumonia

Feed approximately 400 g/ton, varying with body weight and feed consumption, to provide 10 mg/lb body weight/day

caused by Pasteurella multocida susceptible to oxytetracycline to be fed continuously for 7 to 14 days.

For Breeding Swine: Mix an amount of Type B feed to provide approximately 400 g/ton varying with body weight and feed consumption to provide 10 mg/lb body weight/day with non-medicated feed ingredients to make a ton of swine feed for control and treatment of leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by Leptospira pomona susceptible to oxytetracycline to be fed continuously for 14 days.

Feed approximately 400 g/ton, varying with body weight and feed consumption, to provide 10 mg/lb body weight/day

RESIDUE WARNING: Zero-day withdrawal period.

\*The medicated feed label must state a single drug concentration.

Net Weight \_\_\_\_\_ lb (\_\_\_\_\_ kg)

Manufactured by  
Blue Bird Feed Company  
Blue Bird, MD 00000