

19 June 2007 (supercedes 12 July 2006)

Bulk or Bag Label - Page 1 of 2

NET WEIGHT: Bulk or 100 lb (45.4 kg) Bag

Blue Bird

Type C Medicated Swine Feed

STARTER-GROWER-FINISHER FEEDS

Tiamulin
(as hydrogen fumarate)*

Indications

35 g

For control of porcine proliferative enteropathies (ileitis) associated with *Lawsonia intracellularis* and for control of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema*) *hyodysenteriae* susceptible to Tiamulin

200 g

For treatment of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema*) *hyodysenteriae* susceptible to Tiamulin

ACTIVE INGREDIENT

Tiamulin (as hydrogen fumarate) *

GUARANTEED ANALYSIS

Crude protein, not less than	_____%
Lysine, not less than	_____%
Crude fat, not less than	_____%
Crude fiber, not more than	_____%
Calcium	Min. ____% Max. ____%
Phosphorus, not less than	_____%
Salt (if added)	Min. ____% Max. ____%
Sodium (if exceeding salt guarantee)	_____%
Selenium	_____ ppm
Zinc	_____ ppm

INGREDIENTS

Each ingredient must be specifically named in accordance with the names and definitions adopted by AAFCO.

Bulk or Bag Label - Page 2 of 2

FEEDING DIRECTIONS

[sample label text below if for control of porcine proliferative enteropathies (ileitis) associated with *Lawsonia intracellularis*.]

Feed continuously as sole ration for not less than 10 days.

[sample label text below if for control of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema) hyodysenteriae* susceptible to Tiamulin]

Feed continuously as sole ration on premises with a history of swine dysentery but where signs of disease have not yet occurred or following approved treatment of disease.

[sample label text below if for treatment of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema) hyodysenteriae* susceptible to Tiamulin]

Feed continuously as sole ration for 14 consecutive days.

WARNING

Withdraw medicated feed 2 days before slaughter following use at 35 g/ton and 7 days before slaughter following use at 200 g/ton.

CAUTION

1. Use as the only source of tiamulin.
2. For use in swine only. Not for use in swine weighing over 250 lbs.
3. Swine being treated with tiamulin should not have access to feeds containing polyether ionophores (e.g., lasalocid, monensin, narasin, salinomycin and semduramicin) as adverse reactions may occur.
4. If signs of toxicity occur, discontinue use of medicated feed.

PROTECT FROM EXCESSIVE TEMPERATURES (TEMPERATURES THAT EXCEED 40°C OR 104°F). DO NOT STORE AT WARM TEMPERATURES UP TO 104°F (40°C), FOR MORE THAN 7 DAYS.

BLUE BIRD FEED COMPANY
Robin, IN 99999