

RECOMBINATE

Antihemophilic Factor (Recombinant)

Lyophilized Powder for Reconstitution for injection

Reconstitute with 5 mL of Sterile Water for Injection using double-ended needles

DESCRIPTION

RECOMBINATE [Antihemophilic Factor (Recombinant)] is a glycoprotein synthesized by a genetically engineered Chinese Hamster Ovary (CHO) cell line. In culture, the CHO cell line secretes recombinant Factor VIII (rFVIII) into the cell culture medium. The rFVIII is purified from the culture medium utilizing a series of chromatography columns. A key step in the purification process is an immunoaffinity chromatography methodology in which a purification matrix, prepared by immobilization of a monoclonal antibody directed to Factor VIII, is utilized to selectively isolate the rFVIII in the medium. The synthesized rFVIII produced by the CHO cells has the same biological effects as human Factor VIII. Structurally the protein has a similar combination of heterogenous heavy and light chains as found in human Factor VIII.

RECOMBINATE is formulated as a sterile, nonpyrogenic, off-white to faint yellow, lyophilized powder preparation of concentrated recombinant Factor VIII for intravenous injection. RECOMBINATE is available in single-dose vials, which contain nominally 250, 500, 1000, 1500, and 2000 International Units per vial. When reconstituted with the appropriate volume of diluent, the product contains the following stabilizers in maximum amounts: For 5 mL reconstitution volume: 25 mg/mL Albumin (Human), 0.40 mg/mL calcium, 3 mg/mL polyethylene glycol (3350), 360 mEq/L sodium, 110 mM histidine, 1.5 µg/Factor VIII International Unit (IU) polysorbate-80. Recombinant Von Willebrand Factor (rVWF) is coexpressed with the rFVIII and helps to stabilize it. The final product contains not more than 2 ng rVWF/IU rFVIII, which will not have any clinically relevant effect in patients with von Willebrand's disease. The product contains no preservative.

Manufacturing of RECOMBINATE is shared by Baxter Healthcare Corporation and Wyeth BioPharma. The recombinant Antihemophilic Factor Concentrate (For Further Manufacturing Use), is produced by Baxter Healthcare Corporation and Wyeth BioPharma (For Further Manufacturing Use) and subsequently formulated and packaged at Baxter Healthcare Corporation.

Each vial of RECOMBINATE is labeled with the Factor VIII activity expressed in IU per vial. Biological potency is determined by an *in vitro* assay which is referenced to the World Health Organization (WHO) International Standard for Factor VIII:C Concentrate.

CLINICAL PHARMACOLOGY

Factor VIII is the specific clotting factor deficient in patients with hemophilia A (classical hemophilia). Hemophilia A is a genetic bleeding disorder characterized by hemorrhages, which may occur spontaneously or after minor trauma. The administration of RECOMBINATE provides an increase in plasma levels of Factor VIII and can temporarily correct the coagulation defect in these patients. Pharmacokinetic studies on sixty-nine (69) patients revealed the circulating mean half-life for RECOMBINATE to be 14.6 ± 4.9 hours (n=67), which was not statistically significantly different from plasma-derived HEMOFIL M, [Antihemophilic Factor (Human), Method M, Monoclonal Purified]. The mean half-life of HEMOFIL M was 14.7 ± 5.1 hours (n=61). The actual baseline recovery observed with RECOMBINATE was 123.9 ± 47.7 IU/dL (n=23), which is significantly higher than the actual HEMOFIL M baseline recovery of 101.7 ± 31.6 IU/dL (n=61). However, the calculated ratio of actual to expected recovery with RECOMBINATE ($121.2 \pm 48.9\%$) is not different on average from HEMOFIL M ($123.4 \pm 16.4\%$).

The clinical study of RECOMBINATE in previously treated patients (individuals with hemophilia A who had been treated with plasma derived Factor VIII) was based on observations made on a study group of 69 patients. These individuals received cumulative amounts of Factor VIII ranging from 20,914 to 1,383,063 IU over the 48 month study. Patients were given a total of 17,700 infusions totaling 28,090,769 IU RECOMBINATE.

These patients were successfully treated for bleeding episodes on a demand basis and also for the prevention of bleeds (prophylaxis). Spontaneous bleeding episodes successfully managed include hemarthroses, soft tissue and muscle bleeds. Management of hemostasis was also evaluated in surgeries. A total of 24 procedures on 13 patients were performed during this study. These included minor (e.g. tooth extraction) and major (e.g. bilateral osteotomies, thoracotomy and liver transplant) procedures. Hemostasis was maintained perioperatively and postoperatively with individualized Factor VIII replacement.

A study of RECOMBINATE in previously untreated patients was also performed as part of an ongoing study. The study group was comprised of seventy-nine (79)¹ patients, of whom seventy-six (76) had received at least one infusion of RECOMBINATE. To date, this cohort has been given 12,209 infusions totaling over 11,277,043 IU of RECOMBINATE. Hemostasis was appropriately managed in spontaneous bleeding episodes, intracranial hemorrhage and surgical procedures.

INDICATIONS and USAGE

The use of RECOMBINATE [Antihemophilic Factor (Recombinant)] is indicated in hemophilia A (classical hemophilia) for the prevention and control of hemorrhagic episodes.² RECOMBINATE is also indicated in the perioperative management of patients with hemophilia A (classical hemophilia).

RECOMBINATE can be of therapeutic value in patients with acquired Factor VIII inhibitors not exceeding 10 Bethesda Units per mL.³ In clinical studies with RECOMBINATE, patients with inhibitors who were entered into the previously treated patient trial and those previously untreated children who have developed inhibitor activity on study, showed clinical hemostatic response when the titer of inhibitor was less than 10 Bethesda Units per mL. However, in such uses, the dosage of RECOMBINATE should be controlled by frequent laboratory determinations of circulating Factor VIII levels as well as the clinical status of the patient.

RECOMBINATE is not indicated in von Willebrand's disease.

CONTRAINDICATIONS

RECOMBINATE is contraindicated in patients who have manifested life-threatening immediate hypersensitivity reactions, including anaphylaxis, to the product or its components, including bovine, mouse or hamster proteins.

WARNINGS

General

The clinical response to RECOMBINATE may vary. If bleeding is not controlled with the recommended dose, the plasma level of factor VIII should be determined and a sufficient dose of RECOMBINATE should be administered to achieve a satisfactory clinical response. If the patient's plasma factor VIII level fails to increase as expected or if bleeding is not controlled after the expected dose, the presence of an inhibitor (neutralizing antibodies) should be suspected and appropriate testing performed. (see **PRECAUTIONS - Monitoring Laboratory Tests**).

Anaphylaxis and Severe Hypersensitivity Reactions

Allergic type hypersensitivity reactions, including anaphylaxis, have been reported with RECOMBINATE and have been manifested as dizziness pruritus, rash, urticaria, flushing, angioedema/face swelling, laryngeal edema, dyspnea, pallor, pyrexia, nausea, paresthesia, hypotension, and loss of consciousness. Discontinue RECOMBINATE if symptoms occur and seek immediate emergency treatment. RECOMBINATE contains trace amounts of bovine proteins, mouse immunoglobulin G (MuIgG), and hamster (CHO) proteins. Patients treated with this product may develop hypersensitivity to these non-human mammalian proteins.

Neutralizing Antibodies

Patients treated with antihemophilic factor (AHF) products should be carefully monitored for the development of factor VIII inhibitors by appropriate clinical observations and laboratory tests. Inhibitors have been reported following administration of RECOMBINATE predominantly in previously untreated and minimally treated patients. The risk of developing inhibitors is highest during the

first 20 exposure days. If expected plasma factor VIII activity levels are not attained, or if bleeding is not controlled with an expected dose, an assay that measures factor VIII inhibitor concentration should be performed (*see PRECAUTIONS - Monitoring Laboratory Tests*).

PRECAUTIONS

General

Certain components used in the packaging of this product contain natural rubber latex.

Identification of the clotting defect as a Factor VIII deficiency is essential before the administration of RECOMBINATE [Antihemophilic Factor (Recombinant)] is initiated. No benefit may be expected from this product in treating other deficiencies.

Formation of Antibodies to Mouse, Hamster or Bovine Protein

As RECOMBINATE contains trace amounts of mouse protein (maximum of 0.1 ng/IU RECOMBINATE), hamster protein (maximum of 1.5 ng CHO protein/IU RECOMBINATE), and bovine protein (maximum of 1 ng BSA/IU RECOMBINATE) the remote possibility exists that patients treated with this product may develop hypersensitivity to these non-human mammalian proteins.

Information for Patients

The patient and physician should discuss the risks and benefits of this product.

Allergic type hypersensitivity reactions have been observed with RECOMBINATE. Patients should be informed of the early signs of hypersensitivity reactions including hives, generalized urticaria, tightness of the chest, wheezing, hypotension, symptoms of laryngeal edema, and anaphylaxis. Patients should be advised to discontinue use of the product and contact their physician if these symptoms occur. Additionally, patients should be informed that local tissue irritation may occur when infusing RECOMBINATE reconstituted with 5 mL sWFI.

Monitoring Laboratory Tests

- Monitor plasma factor VIII activity levels by the one-stage clotting assay to confirm the adequate factor VIII levels have been achieved and maintained, when clinically indicated. (see *DOSAGE AND ADMINISTRATION*).
- Monitor for development of factor VIII inhibitors. Perform assay to determine if factor VIII inhibitor is present if expected factor VIII activity plasma levels are not attained, or if bleeding is not controlled with the expected dose of RECOMBINATE. Use Bethesda Units (BU) to titer inhibitors.
 - If the inhibitor is less than 10 BU per mL, the administration of additional RECOMBINATE concentrate may neutralize the inhibitor, and may permit an appropriate hemostatic response.
 - Adequate hemostasis may not be achieved if Inhibitor titers are above 10 BU per mL. The inhibitor titer may rise following RECOMBINATE infusion as a result of an anamnestic response to factor VIII. The treatment or prevention of bleeding in such patients requires the use of alternative therapeutic approaches and agents.

Carcinogenesis, Mutagenesis, Impairment of Fertility

RECOMBINATE was tested for mutagenicity at doses considerably exceeding plasma concentrations of Factor VIII *in vitro* and at doses up to ten times the expected maximum clinical dose *in vivo*, and did not cause reverse mutations, chromosomal aberrations, or an increase in micronuclei in bone marrow polychromatic erythrocytes. Long-term studies in animals have not been performed to evaluate carcinogenic potential.

Pediatric Use

RECOMBINATE is appropriate for use in children of all ages, including the newborn. Safety and efficacy studies have been performed in both previously treated (n=23) and previously untreated (n=75) children. (See **Clinical Pharmacology** and **Precautions**).

Pregnancy

Pregnancy Category C. Animal reproduction studies have not been conducted with RECOMBINATE. The safety of RECOMBINATE for use in pregnant women has not been established. It is not known whether RECOMBINATE can cause fetal harm when administered to a pregnant woman or can affect reproductive capacity. Physicians should carefully consider the potential risks and benefits for each specific patient before prescribing RECOMBINATE. RECOMBINATE should be given to a pregnant woman only if clearly needed.

Nursing Mothers

It is not known whether this drug is excreted into human milk. Because many drugs are excreted into human milk, caution should be exercised if RECOMBINATE is administered to nursing mothers. RECOMBINATE should be given to nursing mothers only if clinically needed.

ADVERSE REACTIONS

Adverse Reactions from Clinical Trials

During controlled clinical studies with RECOMBINATE enrolling 210 subjects, the most commonly reported adverse drug reactions were chills, flushing, rash and epistaxis.

System Organ Class (SOC)	Preferred MedDRA Term	Number of Subjects	Percent of Evaluable Subjects*
GASTROINTESTINAL DISORDERS	Nausea	1	0.48
GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS	Chills	3	1.43
	Fatigue	1	0.48
	Pyrexia	1	0.48
INFECTIONS AND INFESTATIONS	Ear infections	1	0.48

INVESTIGATIONS	Acoustic stimulation tests abnormal	1	0.48
MUSCULOSKELETAL AND CONNECTIVE TISSUES DISORDERS	Pain in extremity	1	0.48
NERVOUS SYSTEM DISORDERS	Dizziness	1	0.48
	Tremors	1	0.48
RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS	Pharyngolaryngeal pain	1	0.48
SKIN AND SUBCUTANEOUS TISSUE DISORDERS	Hyperhidrosis	1	0.48
	Pruritus	1	0.48
	Rash	2	0.95
	Rash maculopapular	1	0.48
VASCULAR DISORDERS	Epistaxis	1 [†]	0.48
	Flushing	2	0.95
	Hematoma	1	0.48
	Hypotension	1	0.48
	Pallor	1	0.48
	Peripheral coldness	1	0.48

* Number of evaluable subjects experiencing the event/total number of evaluable subjects [% relative to 210, the total number of unique subjects who received at least 1 infusion of RECOMBINATE].

†One subject experienced 11 events of epistaxis.

During the Previously Treated Patients (PTP) study, none of the 71 subjects developed *de novo* evidence of Factor VIII inhibitor. However, during the phase II/III portion of the study, 1 subject with a history of inhibitors exhibited inhibitor activity at 6 months (0.8 Bethesda Units [BU]), which resolved by 9 months. One other subject in this study had detectable Factor VIII inhibitor at baseline (1.26 BU) and exhibited an anamnestic response at 6 months (10.3 BU). During a prospective pharmaco-surveillance study of subjects who received batches of RECOMBINATE containing modestly increased Chinese Hamster Ovary (CHO) cell protein levels, none of the 34 treated subjects developed a Factor VIII inhibitor.

During the Previously Untreated Patients (PUP) study, 22 of the 73 evaluable subjects developed inhibitors to Factor VIII. Of these, 13 subjects displayed no detectable Factor VIII inhibitors at study exit.

Post-Marketing Adverse Reactions

In addition to the adverse reactions noted in clinical trials, the following adverse reactions have been reported in the post-marketing experience. These adverse reactions are listed by MedDRA (version 12.1) System Organ Class (SOC), then by MedDRA coding system Preferred Term in order of severity.

BLOOD AND LYMPHATIC SYSTEM DISORDERS: Factor VIII inhibition

CARDIAC DISORDERS: Tachycardia, Cyanosis

GASTROINTESTINAL DISORDERS: Vomiting, Abdominal pain

GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS:

Malaise, Injection site reactions, Chest pain, Chest discomfort

IMMUNE SYSTEM DISORDERS: Anaphylactic reaction, Hypersensitivity

NERVOUS SYSTEM DISORDERS: Loss of consciousness, Headache, Paresthesia

RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS:

Dyspnea, Cough, Laryngeal edema

SKIN AND SUBCUTANEOUS TISSUE DISORDERS: Angioedema, Urticaria, Erythema

DOSAGE and ADMINISTRATION

Each vial of RECOMBINATE is labeled with the Factor VIII activity expressed in IU per vial. This potency assignment is referenced to the World Health Organization International Standard for Factor VIII:C Concentrate and is evaluated by appropriate methodology to ensure accuracy of the results.

The expected *in vivo* peak increase in Factor VIII level expressed as IU/dL of plasma or % (percent) of normal can be estimated by multiplying the dose administered per kg body weight (IU/kg) by two. This calculation is based on the clinical findings of Abildgaard *et al*⁴ and is supported by the data generated by 419 clinical pharmacokinetic studies with RECOMBINATE in 67 patients over time. This pharmacokinetic data demonstrated a peak recovery point above the pre-infusion baseline of approximately 2.0 IU/dL per IU/kg body weight.

Examples (Assuming patient's baseline Factor VIII level is at <1%):

- (1) A dose of 1750 IU RECOMBINATE administered to a 70 kg patient, *i.e.* 25 IU/kg (1750 IU/70 kg), should be expected to cause a peak post-infusion Factor VIII increase of $25 \text{ IU/kg} \times 2 \text{ (IU/dL)/(IU/kg)} = 50 \text{ IU/dL}$ (50% of normal).
- (2) A peak level of 70% is required in a 40 kg child. In this situation the dose would be $70 \text{ IU/dL} / [2 \text{ (IU/dL)/(IU/kg)}] \times 40 \text{ kg} = 1400 \text{ IU}$.

Recommended Dosage Schedule

Physician supervision of the dosage is required. The following dosage schedule may be used as a guide.

Hemorrhage		
Degree of hemorrhage	Required peak post-infusion Factor VIII activity in the blood (as % of normal or IU/dL plasma)	Frequency of Infusion
Early hemarthrosis or muscle bleed or oral bleed	20-40	Begin infusion every 12 to 24 hours for one-three days until the bleeding episode is resolved (as indicated by pain), or healing is achieved.
More extensive hemarthrosis, muscle bleed, or hematoma	30-60	Repeat infusion every 12 to 24 hours for (usually) three days or more until pain and disability are resolved.
Life threatening bleeds such as head injury, throat bleed, severe abdominal pain	60-100	Repeat infusion every 8 to 24 hours until threat is resolved.
Surgery		
Type of operation		
Minor surgery, including tooth extraction	60-80	A single infusion plus oral antifibrinolytic therapy within one hour is sufficient in approximately 70% of cases.
Major surgery	80-100 (pre- and post-operative)	Repeat infusion every 8 to 24 hours depending on state of healing

If bleeding is not controlled with the recommended dose, the plasma level of Factor VIII should be determined and a sufficient dose of RECOMBINATE should be administered to achieve a satisfactory clinical response

The careful control of the substitution therapy is especially important in cases of major surgery or life threatening hemorrhages. In presence of a low titer

inhibitor, doses larger than those recommended may be necessary as per standard care.

Although dosage can be estimated by the calculations above, it is strongly recommended that whenever possible, appropriate laboratory tests including serial Factor VIII assays be performed on the patient's plasma at suitable intervals to assure that adequate Factor VIII levels have been reached and are maintained. Patients should be evaluated for the development of Factor VIII inhibitors, if the expected plasma Factor VIII activity levels are not attained, or if bleeding is not controlled with an appropriate dose.

Reconstitution: Use Aseptic Technique

1. Bring RECOMBINATE, (dry factor concentrate) and Sterile Water for Injection, USP, (diluent) to room temperature.
2. Remove caps from concentrate and diluent vials.
3. Cleanse stoppers with germicidal solution and allow to dry prior to use. Place vials on a flat surface
4. Remove protective covering from one end of double-ended needle and insert exposed needle through the center of the stopper.
5. Remove protective covering from other end of double-ended needle. Invert diluent vial over the upright RECOMBINATE vial, then rapidly insert free end of the needle through the RECOMBINATE vial stopper at its center. The vacuum in the vial will draw in the diluent.
6. Disconnect the two vials by removing needle from diluent vial stopper, then remove needle from RECOMBINATE vial. Swirl gently until RECOMBINATE is completely dissolved, otherwise active material will be removed by the filter needle. After reconstitution, the solution should be colorless to faint yellow, and substantially free from foreign particles.

NOTE: Do not refrigerate after reconstitution. (See **Administration**)

Administration: Use Aseptic Technique

RECOMBINATE is administered by intravenous (IV) injection after reconstitution.

Administer at room temperature.

RECOMBINATE should be administered not more than 3 hours after reconstitution.

Intravenous Syringe Injection

Parenteral drug products should be inspected for particulate matter and discoloration prior to administration, whenever solution and container permit. The solution should be colorless to faint yellow in appearance. If not, do not use the solution and notify Baxter immediately.

Plastic syringes are recommended for use with this product since proteins such as RECOMBINATE tend to stick to the surface of glass syringes.

1. Attach filter needle to a disposable syringe and draw back plunger to admit air into the syringe.
2. Insert the needle into reconstituted RECOMBINATE.
3. Inject air into vial and then withdraw the reconstituted material into the syringe.
4. Remove and discard the filter needle from the syringe; attach a suitable needle and inject intravenously as instructed under **Rate of Administration**.
5. If a patient is to receive more than one vial of RECOMBINATE, the contents of multiple vials may be drawn into the same syringe by drawing up each vial through a separate unused filter needle. Filter needles are intended to filter the contents of a single vial of RECOMBINATE only.

Rate of Administration

The rate of administration should be a rate that ensures the comfort of the patient. Preparations of RECOMBINATE can be administered at a rate of up to 5 mL per minute when reconstituted with 5 mL of sWFI.

The pulse rate should be determined before and during administration of RECOMBINATE. Should a significant increase in pulse rate occur, reducing the rate of administration or temporarily halting the injection usually allows the symptoms to disappear promptly.

HOW SUPPLIED

RECOMBINATE is available in five different strengths in single-dose vials. The strength is designated on the outer box and on the vial label using the following color codes:

Color Code	Dosage Strength	RECOMBINATE Supplied with 5 mL sWFI
Light blue bar	220 – 400 IU per vial	NDC 0944-2841-15
Pink bar	401-800 IU per vial	NDC 0944-2842-15
Green bar	801-1240 IU per vial	NDC 0944-2843-15
Purple bar	1241-1800 IU per vial	NDC 0944-2844-15
Orange bar	1801-2400 IU per vial	NDC 0944-2845-15

RECOMBINATE is packaged with 5 mL of Sterile Water for Injection, USP, a double-ended needle, a filter needle, one physician insert and one patient insert.

Storage

RECOMBINATE can be refrigerated [2° - 8°C (36° - 46°F)] or stored at room temperature, not to exceed 30°C (86°F). Avoid freezing to prevent damage to the diluent vial. Do not use beyond the expiration date printed on the box.

CLINICAL STUDIES

Over the investigational period of the original safety and efficacy study of RECOMBINATE, none of the 69 subjects without an inhibitor at entry into the study, developed an inhibitor. In the previously untreated patient group there were 73 eligible subjects with Factor VIII levels less than or equal to 2% who received at least one RECOMBINATE treatment (median days 100, range 3-821) and who were tested for an inhibitor after treatment with RECOMBINATE. Of this group, 23 individuals (32%) developed a detectable inhibitor (median days on treatment at time of detection 10, range 3-69) and of these, 8 subjects (11%) showed a titer greater than 10 B.U.

REFERENCES

1. Bray GL, Gomperts ED, Courter S, Gruppo R, *et al*: A Multicenter Study of Recombinant Factor VIII (Recombinate): Safety, Efficacy, and Inhibitor Risk in Previously Untreated Patients with Hemophilia A **Blood** **83**:2428-2435, 1994
2. White GC, McMillan CW, Kingdon HS, *et al*: Use of recombinant antihemophilic factor in the treatment of two patients with classic hemophilia. **New Eng J Med** **320**:166-170, 1989
3. Kessler CM: An Introduction to Factor VIII Inhibitors: The Detection and Quantitation. **Am J Med** **91 (Suppl 5A)**:1S-5S, 1991
4. Abildgaard CF, Simone JV, Corrigan JJ, *et al*: Treatment of hemophilia with glycine-precipitated Factor VIII. **New Eng J Med** **275**:471-475, 1966

To enroll in the confidential, industry-wide Patient Notification System, call 1-888-UPDATE U (1-888-873-2838).

Baxter, RECOMBINATE, and HEMOFIL, are trademarks of Baxter International Inc.

Manufactured by:

Baxter Healthcare Corporation
Westlake Village, CA 91362 USA
U.S. License No. 140

Printed in USA
Issued: March 2010

Patient Information
RECOMBINATE
[Antihemophilic Factor (Recombinant)]

This leaflet summarizes important information about RECOMBINATE. Please read it carefully before using this medicine. This information does not take the place of talking with your healthcare provider, and it does not include all of the important information about RECOMBINATE. If you have any questions after reading this, ask your healthcare provider.

Do not attempt to self-infuse unless you have been taught how by your doctor or hemophilia center.

What is RECOMBINATE [Antihemophilic Factor (Recombinant)]?

RECOMBINATE is a medicine used to replace clotting factor (Factor VIII or antihemophilic factor) that is missing in people with hemophilia A (also called “classic” hemophilia). Hemophilia A is an inherited bleeding disorder that prevents blood from clotting normally.

RECOMBINATE is used to prevent and control bleeding in people with hemophilia A.

RECOMBINATE is not used to treat von Willebrand’s Disease.

Who should not use RECOMBINATE [Antihemophilic Factor (Recombinant)]?

You should not use RECOMBINATE if you

- are allergic to mouse, hamster or bovine proteins.
- are allergic to any ingredients in RECOMBINATE (such as calcium, histidine, human albumin, polyethylene glycol, polysorbate-80 and sodium).

Tell your healthcare provider if you are pregnant or breast-feeding because RECOMBINATE may not be right for you.

How should I use RECOMBINATE [Antihemophilic Factor (Recombinant)]?

RECOMBINATE is given directly into the blood stream.

You may infuse RECOMBINATE at a hemophilia treatment center, at your healthcare provider’s office or in your home. You should be trained on how to do

infusions by your hemophilia treatment center or healthcare provider. Many people with hemophilia A learn to infuse their RECOMBINATE by themselves or with the help of a family member.

You must carefully follow your doctor's or other healthcare provider's instructions regarding the dose and schedule for infusing RECOMBINATE so that your treatment will work best for you.

Your healthcare provider will tell you how much RECOMBINATE to use based on your weight, the severity of your hemophilia A, and where you are bleeding.

You may have to have blood tests done after getting RECOMBINATE to be sure that your blood level of Factor VIII is high enough to clot your blood.

Call your healthcare provider right away if your bleeding does not stop after taking RECOMBINATE.

What should I tell my healthcare provider before I use RECOMBINATE [Antihemophilic Factor (Recombinant)]?

You should tell your healthcare provider if you

- have or have had any medical problems.
- take any medicines, including non-prescription medicines, dietary supplements and herbal remedies.
- have any allergies, including allergies to mouse, hamster or bovine proteins.
- are nursing.
- are pregnant or planning to become pregnant.
- have been told that you have inhibitors to Factor VIII (because Factor VIII may not work for you).

What are the possible side effects of RECOMBINATE [Antihemophilic Factor (Recombinant)]?

You could have an allergic reaction to RECOMBINATE.

Call your healthcare provider right away and stop treatment if you get a:

- Rash or hives
- itching
- tightness of the throat
- chest pain or tightness
- difficulty breathing
- light-headed, dizziness

- fainting.

The most common side effects are chills, flushing, rash and nose bleeds. These are not all possible side effects with RECOMBINATE. You can ask your healthcare provider for information that is written for healthcare professionals. Tell your doctor about any side effect that bothers you or that does not go away.

What are the RECOMBINATE [Antihemophilic Factor (Recombinant)] dosage strengths?

RECOMBINATE comes in five different dosage strengths. The actual strength will be imprinted on the label and on the box. The five different strengths are coded, as follows:

Light-blue	Nominal dosage strength of approximately 250 IU per vial (220 – 400 IU/vial).
Pink	Nominal dosage strength of approximately 500 IU per vial (401 – 800 IU/vial).
Green	Nominal dosage strength of approximately 1000 IU per vial (801 – 1240 IU/vial).
Purple	Nominal dosage strength of approximately 1500 IU per vial (1241-1800 IU/vial)
Orange	Nominal dosage strength of approximately 2000 IU per vial (1801-2400 IU/vial)

Always check the potency printed on the label to make sure you are using the strength prescribed by your doctor. Always check the expiration date printed on the box. You should not use the product after the expiration date printed on the box.

How do I store RECOMBINATE [Antihemophilic Factor (Recombinant)]?

RECOMBINATE vials containing powdered product (without sterile diluent added) should be stored in a refrigerator (2° to 8°C [36° to 46°F]) or at room temperature (up to 30°C [86°F]).

If you choose to store RECOMBINATE at room temperature:

- it should remain at room temperature until infused.
- do not put room temperature product back in the refrigerator..

Store vials in their original box and protect them from extreme exposure to light.

Do not freeze.

Reconstituted product (after mixing dry product with wet diluent) must be used within 3 hours and cannot be stored or refrigerated. Any RECOMBINATE left in the vial at the end of your infusion should be discarded.

What else should I know about RECOMBINATE [Antihemophilic Factor (Recombinant)] and hemophilia A?

Your body may form inhibitors to Factor VIII. An inhibitor is part of the body's normal defense system. If you form inhibitors, it may stop RECOMBINATE from working properly. Call your healthcare provider right away if your bleeding does not stop after taking RECOMBINATE. Consult with your healthcare provider to make sure you are carefully monitored with blood tests for the development of inhibitors to Factor VIII.

Resources at Baxter available to the patients:

Contact Baxter to receive more product information:

Baxter Customer Service 1-800-423-2090

Baxter, RECOMBINATE, and HEMOFIL are trademarks of Baxter International Inc.

Manufactured by:

Baxter Healthcare Corporation
Westlake Village, CA 91362 USA
U.S. License No. 140
Printed in USA

Issued: March 2010

RECOMBINATE

Antihemophilic Factor (Recombinant)

(For intravenous use only)

1. In a quiet place, prepare a clean surface. Let the vial with the FVIII concentrate and the Sterile Water for Injection (diluent) warm up to room temperature.



2. After washing your hands and putting on sterile gloves, remove caps from the concentrate and diluent vial to expose the centers of the rubber stoppers.



3. Disinfect the stoppers with an alcohol swab or other suitable solution suggested by your doctor or hemophilia center.



4. Remove the protective covering from one end of the double-ended transfer needle.



5. Insert the exposed short part of the needle through the diluent stopper.



6. Remove the protective covering from the other end of the double-ended transfer needle.



7. Insert the free end of the needle through the concentrate vial stopper at its center. The vacuum in the vial will draw in the diluent. Invert the diluent vial over the upright concentrate vial.



8. Disconnect the two vials by removing the needle from the diluent vials stopper, then remove the needle from the concentrate vials. Do not recap the needle! Place the needle in a hard-walled Sharps container for proper disposal.



9. Gently roll the vial between palms until all material is dissolved. **Do not shake.** Check to make sure the product is completely dissolved.




10. Attach filter needle to a disposable syringe and draw back the plunger to allow air into the syringe. Insert the needle into the reconstituted FVIII concentrate. Inject air into the vial.




11. Withdraw the solution into the syringe.



12. Prepare the injection site by wiping with an alcohol swab or other suitable solution suggested by your doctor or hemophilia center.



13. Remove and discard the filter needle from the syringe; attach a suitable needle. Use a winged infusion set if available. Do not infuse the product any faster than 5 mL per minute for RECOMBINATE dissolved with 5 mL of sWFI. After infusion, apply pressure with sterile gauze to the infusion site for 3 minutes. Do not recap the needle after infusion. Place it with the used syringe in a hard-walled Sharps container for proper disposal.



Redness of the skin or irritation may be seen when infusing RECOMBINATE dissolved with 5 mL diluent.

14. After infusion, remove peel-off label from the concentrate vial and place it in your factor log. Clean up any spilled blood with freshly prepared 10% bleach solution or soap and water.



IMPORTANT: Contact your doctor or local Hemophilia Treatment Center if you experience any problems.